

Stephane Moyses

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3601113/publications.pdf>

Version: 2024-02-01

16
papers

149
citations

1307594

7
h-index

1199594

12
g-index

16
all docs

16
docs citations

16
times ranked

133
citing authors

#	ARTICLE	IF	CITATIONS
1	Conformationally Sensitive Infrared Vibrations of the Syndiotactic Polystyrene/Ethylbenzene Complex. <i>Macromolecules</i> , 1999, 32, 2684-2689.	4.8	31
2	Structural and morphological features of the syndiotactic polystyrene/ethylbenzene complex. <i>Polymer</i> , 1998, 39, 3537-3544.	3.8	20
3	Characterization of Randomly Branched Poly(vinylidene fluoride). <i>Macromolecules</i> , 2008, 41, 2011-2021.	4.8	18
4	Neutron scattering studies of syndiotactic polystyrene/ethylbenzene. <i>Polymer</i> , 1998, 39, 3665-3674.	3.8	14
5	Separation of Branched Poly(bisphenol A carbonate) Structures by Solvent Gradient at Near-Critical Conditions and Two-Dimensional Liquid Chromatography. <i>Analytical Chemistry</i> , 2018, 90, 5422-5429.	6.5	10
6	Two-dimensional chromatography applied to the study of the thermo-oxidative degradation of poly(styrene- <i>b</i> -butadiene) star block copolymers. <i>Journal of Separation Science</i> , 2012, 35, 1741-1747.	2.5	9
7	Effect of Rheological Strain Hardening on Extrusion Blown Film of Polyvinylidene Fluoride. <i>Journal of Plastic Film and Sheeting</i> , 2007, 23, 203-219.	2.2	8
8	Solution Properties of Poly(Methyl Methacrylate) in Dimethylsulfoxide. <i>International Journal of Polymer Analysis and Characterization</i> , 2008, 13, 413-427.	1.9	8
9	2D chromatography applied to the study of block copolymers synthesized by nitroxide-mediated controlled free radical seeded emulsion polymerization. <i>Journal of Separation Science</i> , 2010, 33, 3511-3518.	2.5	6
10	Correlation between Comprehensive 2D Liquid Chromatography and Monte Carlo Simulations for Branched Polymers. <i>Macromolecules</i> , 2018, 51, 4829-4839.	4.8	6
11	Molecular dimensions in the $\hat{1}$ and $\hat{1}^3$ phases of syndiotactic polystyrene, using small angle neutron scattering. <i>Polymer</i> , 1999, 40, 3269-3274.	3.8	5
12	2D chromatography with optimized size exclusion chromatography resolution and multi-angle light scattering coupling. <i>Journal of Separation Science</i> , 2010, 33, 1480-1486.	2.5	5
13	The chromatography of poly(phenylene ether) on a porous graphitic carbon sorbent. <i>Journal of Chromatography A</i> , 2016, 1468, 136-142.	3.7	5
14	Neutron scattering studies of syndiotactic polystyrene crystals. <i>Macromolecular Symposia</i> , 1997, 114, 145-150.	0.7	2
15	Separation and Simultaneous Molecular Weight Determination of PVDF in PVDF/PMMA Blends Based on Adsorption/Size Exclusion. <i>International Journal of Polymer Analysis and Characterization</i> , 2004, 9, 29-37.	1.9	1
16	The effect of in situ-formed copolymers on the morphology of reactive poly(phenylene) Tj ETQq0 0 0 rgBT /Overlap 10 Tf 50 142 Td (3.8	1