

Concetta De Stefano

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3602441/publications.pdf>

Version: 2024-02-01

234
papers

4,708
citations

117453

34
h-index

197535

49
g-index

235
all docs

235
docs citations

235
times ranked

2563
citing authors

#	ARTICLE	IF	CITATIONS
1	Formation and stability of phytate complexes in solution. <i>Coordination Chemistry Reviews</i> , 2008, 252, 1108-1120.	9.5	180
2	On the possibility of determining the thermodynamic parameters for the formation of weak complexes using a simple model for the dependence on ionic strength of activity coefficients: Na ⁺ , K ⁺ , and Ca ²⁺ complexes of low molecular weight ligands in aqueous solution. <i>Journal of the Chemical Society Dalton Transactions</i> , 1985, , 2353.	1.1	136
3	The PAH composition of surface sediments from Stagnone coastal lagoon, Marsala (Italy). <i>Marine Chemistry</i> , 2006, 99, 117-127.	0.9	90
4	The determination of formation constants of weak complexes by potentiometric measurements: experimental procedures and calculation methods. <i>Talanta</i> , 1987, 34, 933-938.	2.9	86
5	Chelating Agents for the Sequestration of Mercury(II) and Monomethyl Mercury(II). <i>Current Medicinal Chemistry</i> , 2014, 21, 3819-3836.	1.2	74
6	Speciation of phytate ion in aqueous solution. Alkali metal complex formation in different ionic media. <i>Analytical and Bioanalytical Chemistry</i> , 2003, 376, 1030-1040.	1.9	64
7	The inorganic speciation of tin(II) in aqueous solution. <i>Geochimica Et Cosmochimica Acta</i> , 2012, 87, 1-20.	1.6	63
8	Copper(II) complexes of N-(phosphonomethyl)glycine in aqueous solution: a thermodynamic and spectrophotometric study. <i>Talanta</i> , 1997, 45, 425-431.	2.9	61
9	Hydrolysis of (CH ₃) ₂ Sn ²⁺ in Different Ionic Media: Salt Effects and Complex Formation. <i>Journal of Chemical & Engineering Data</i> , 1996, 41, 511-515.	1.0	60
10	Polyacrylate Protonation in Various Aqueous Ionic Media at Different Temperatures and Ionic Strengths. <i>Journal of Chemical & Engineering Data</i> , 2000, 45, 876-881.	1.0	60
11	Protonation of carbonate in aqueous tetraalkylammonium salts at 25°C. <i>Talanta</i> , 2006, 68, 1102-1112.	2.9	57
12	The interaction of amino acids with the major constituents of natural waters at different ionic strengths. <i>Marine Chemistry</i> , 2000, 72, 61-76.	0.9	54
13	Salt effects on the protonation of ortho-phosphate between 10 and 50°C in aqueous solution. A complex formation model. <i>Journal of Solution Chemistry</i> , 1991, 20, 495-515.	0.6	52
14	Study and characterization of the ancient bricks of monastery of "San Filippo di Fragalà" in Frazzanò (Sicily). <i>Analytica Chimica Acta</i> , 2004, 519, 103-111.	2.6	52
15	Dependence on Ionic Strength of Protonation Enthalpies of Polycarboxylate Anions in NaCl Aqueous Solution. <i>Journal of Chemical & Engineering Data</i> , 2001, 46, 1417-1424.	1.0	51
16	Thermodynamic parameters for the protonation of carboxylic acids in aqueous tetraethylammonium iodide solutions. <i>Journal of Solution Chemistry</i> , 1990, 19, 569-587.	0.6	45
17	Speciation of Phytate Ion in Aqueous Solution. Protonation Constants in Tetraethylammonium Iodide and Sodium Chloride. <i>Journal of Chemical & Engineering Data</i> , 2003, 48, 114-119.	1.0	45
18	Equilibrium studies in natural fluids: a chemical speciation model for the major constituents of sea water. <i>Chemical Speciation and Bioavailability</i> , 1994, 6, 65-84.	2.0	44

#	ARTICLE	IF	CITATIONS
19	Polyacrylates in aqueous solution. The dependence of protonation on molecular weight, ionic medium and ionic strength. <i>Reactive and Functional Polymers</i> , 2003, 55, 9-20.	2.0	44
20	Thermodynamic parameters for the binding of inorganic and organic anions by biogenic polyammonium cations. <i>Talanta</i> , 2001, 54, 1135-1152.	2.9	42
21	Protonation Constants of Ethylenediamine, Diethylenetriamine, and Spermine in NaCl(aq), NaI(aq), (CH ₃) ₄ NCl(aq), and (C ₂ H ₅) ₄ NI(aq) at Different Ionic Strengths and t = 25 °C. <i>Journal of Chemical & Engineering Data</i> , 2005, 50, 1917-1923.	1.0	42
22	Acid-Base Properties of Synthetic and Natural Polyelectrolytes: Experimental Results and Models for the Dependence on Different Aqueous Media. <i>Journal of Chemical & Engineering Data</i> , 2009, 54, 589-605.	1.0	42
23	Thermodynamic Properties of Dopamine in Aqueous Solution. Acid-Base Properties, Distribution, and Activity Coefficients in NaCl Aqueous Solutions at Different Ionic Strengths and Temperatures. <i>Journal of Chemical & Engineering Data</i> , 2013, 58, 2835-2847.	1.0	41
24	Formation and stability of zinc(II) and cadmium(II) citrate complexes in aqueous solution at various temperatures. <i>Talanta</i> , 1986, 33, 763-767.	2.9	40
25	The calculation of equilibrium concentrations in large multimetal/multiligand systems. <i>Analytica Chimica Acta</i> , 1986, 191, 385-398.	2.6	40
26	Speciation of phytate ion in aqueous solution. Sequestration of magnesium and calcium by phytate at different temperatures and ionic strengths, in NaCl aq. <i>Biophysical Chemistry</i> , 2006, 124, 18-26.	1.5	40
27	Electrochemical Study on the Stability of Phytate Complexes with Cu ²⁺ , Pb ²⁺ , Zn ²⁺ , and Ni ²⁺ : A Comparison of Different Techniques. <i>Journal of Chemical & Engineering Data</i> , 2010, 55, 4757-4767.	1.0	40
28	Modeling the acid-base properties of glutathione in different ionic media, with particular reference to natural waters and biological fluids. <i>Amino Acids</i> , 2012, 43, 629-648.	1.2	40
29	Speciation of phytate ion in aqueous solution. <i>Thermochimica Acta</i> , 2004, 423, 63-69.	1.2	37
30	Salt effects on the protonation and on alkali and alkaline earth metal complex formation of 1,2,3-propanetricarboxylate in aqueous solution. <i>Talanta</i> , 1994, 41, 1715-1722.	2.9	35
31	Thermodynamic parameters for the formation of glycine complexes with magnesium(II), calcium(II), lead(II), manganese(II), cobalt(II), nickel(II), zinc(II) and cadmium(II) at different temperatures and ionic strengths, with particular reference to natural fluid conditions. <i>Thermochimica Acta</i> , 1995, 255, 109-141.	1.2	35
32	Dependence on Ionic Strength of the Hydrolysis Constants for Dioxouranium(VI) in NaCl(aq) and NaNO ₃ (aq), at pH < 6 and t = 25 °C. <i>Journal of Chemical & Engineering Data</i> , 2002, 47, 533-538.	1.0	35
33	Sequestering ability of polyaminopolycarboxylic ligands towards dioxouranium(VI) cation. <i>Journal of Alloys and Compounds</i> , 2006, 424, 93-104.	2.8	35
34	Speciation of phytate ion in aqueous solution. Protonation constants and copper(II) interactions in NaNO ₃ aq at different ionic strengths. <i>Biophysical Chemistry</i> , 2007, 128, 176-184.	1.5	34
35	Ionic-strength dependence of formation constants—XII A model for the effect of background on the protonation constants of amines and amino-acids. <i>Talanta</i> , 1989, 36, 903-907.	2.9	33
36	Effects of salt on the protonation in aqueous solution of triethylenetetramine and tetraethylenepentamine. <i>Journal of Solution Chemistry</i> , 1993, 22, 927-940.	0.6	33

#	ARTICLE	IF	CITATIONS
37	Speciation of Phytate Ion in Aqueous Solution. Sequestering Ability toward Mercury(II) Cation in NaCl(aq) at Different Ionic Strengths. <i>Journal of Agricultural and Food Chemistry</i> , 2006, 54, 1459-1466.	2.4	33
38	Modeling solubility, acid-base properties and activity coefficients of amoxicillin, ampicillin and (+)6-aminopenicillanic acid, in NaCl(aq) at different ionic strengths and temperatures. <i>European Journal of Pharmaceutical Sciences</i> , 2012, 47, 661-677.	1.9	33
39	Thermodynamics of formation of magnesium, calcium, strontium and barium complexes with 2,2'-bipyridyl and 1,10-phenanthroline, at different ionic strengths in aqueous solution. <i>Talanta</i> , 1985, 32, 675-677.	2.9	32
40	Speciation of phytate ion in aqueous solution. Non covalent interactions with biogenic polyamines. <i>Chemical Speciation and Bioavailability</i> , 2003, 15, 29-36.	2.0	32
41	Modeling ATP protonation and activity coefficients in NaCl(aq) and KCl(aq) by SIT and Pitzer equations. <i>Biophysical Chemistry</i> , 2006, 121, 121-130.	1.5	32
42	Some thermodynamic properties of dl-Tyrosine and dl-Tryptophan. Effect of the ionic medium, ionic strength and temperature on the solubility and acid-base properties. <i>Fluid Phase Equilibria</i> , 2012, 314, 185-197.	1.4	32
43	Binding of carboxylic ligands by protonated amines. <i>Journal of the Chemical Society, Faraday Transactions</i> , 1996, 92, 4219-4226.	1.7	31
44	Binding of polyanions by biogenic amines. I. Formation and stability of protonated putrescine and cadaverine complexes with inorganic anions. <i>Talanta</i> , 1998, 46, 1085-1093.	2.9	31
45	Quantitative parameters for the sequestering capacity of polyacrylates towards alkaline earth metal ions. <i>Talanta</i> , 2003, 61, 181-194.	2.9	30
46	Dioxouranium(VI) carboxylate complexes. A calorimetric and potentiometric investigation of interaction with oxalate at infinite dilution and in NaCl aqueous solution at $I=1.0\text{ mol L}^{-1}$ and $T=25^\circ\text{C}$. <i>Talanta</i> , 2007, 71, 948-963.	2.9	30
47	Speciation of Phytate Ion in Aqueous Solution. Thermodynamic Parameters for Zinc(II) Sequestration at Different Ionic Strengths and Temperatures. <i>Journal of Solution Chemistry</i> , 2009, 38, 115-134.	0.6	30
48	Ionic Strength Dependence of Protonation Constants of N-Alkyl Substituted Open Chain Diamines in NaCl(aq). <i>Journal of Chemical & Engineering Data</i> , 2004, 49, 109-115.	1.0	29
49	Formation and Stability of Cadmium(II)/Phytate Complexes by Different Electrochemical Techniques. Critical Analysis of Results. <i>Journal of Solution Chemistry</i> , 2010, 39, 179-195.	0.6	29
50	Acid-base and UV behavior of 3-(3,4-dihydroxyphenyl)-propenoic acid (caffeic acid) and complexing ability towards different divalent metal cations in aqueous solution. <i>Journal of Molecular Liquids</i> , 2014, 195, 9-16.	2.3	29
51	Ionic strength dependence of formation constants and proton activity coefficients at various temperatures and ionic strengths and their use in the study of complex equilibria. <i>Talanta</i> , 1987, 34, 593-598.	2.9	28
52	Acid-Base Properties and Alkali and Alkaline Earth Metal Complex Formation in Aqueous Solution of Diethylenetriamine- N^2 , N^3 -pentakis(methylenephosphonic acid) Obtained by an Efficient Synthetic Procedure. <i>Industrial & Engineering Chemistry Research</i> , 2014, 53, 9544-9553.	1.8	28
53	Salt effects on the protonation of l-histidine and l-aspartic acid: a complex formation model. <i>Thermochimica Acta</i> , 1991, 177, 39-57.	1.2	27
54	Chemical speciation of amino acids in electrolyte solutions containing major components of natural fluids. <i>Chemical Speciation and Bioavailability</i> , 1995, 7, 1-8.	2.0	27

#	ARTICLE	IF	CITATIONS
55	Hydrolysis of (CH ₃) ₃ Sn ⁺ in Various Salt Media. <i>Journal of Solution Chemistry</i> , 1999, 28, 959-972.	0.6	27
56	Thermodynamic and spectroscopic study for the interaction of dimethyltin(IV) with L-cysteine in aqueous solution. <i>Biophysical Chemistry</i> , 2008, 133, 19-27.	1.5	27
57	Composition, Distribution, and Sources of Polycyclic Aromatic Hydrocarbons in Sediments of the Gulf of Milazzo (Mediterranean Sea, Italy). <i>Polycyclic Aromatic Compounds</i> , 2014, 34, 397-424.	1.4	27
58	Complexation of Hg ²⁺ , CH ₃ Hg ⁺ , Sn ²⁺ and (CH ₃) ₂ Sn ²⁺ with phosphonic NTA derivatives. <i>New Journal of Chemistry</i> , 2016, 40, 1443-1453.	1.4	27
59	The single salt approximation for the major components of seawater: association and acid-base properties. <i>Chemical Speciation and Bioavailability</i> , 1998, 10, 27-30.	2.0	26
60	Modelling of proton and metal exchange in the alginate biopolymer. <i>Analytical and Bioanalytical Chemistry</i> , 2005, 383, 587-596.	1.9	26
61	Thermodynamic Protonation Parameters of some Sulfur-Containing Anions in NaCl(aq) and (CH ₃) ₄ NCl(aq) at T=25°C. <i>Journal of Solution Chemistry</i> , 2009, 38, 1225-1245.	0.6	26
62	Hydrolysis of methyltin(IV) trichloride in aqueous NaCl and NaNO ₃ solutions at different ionic strengths and temperatures. <i>Applied Organometallic Chemistry</i> , 1999, 13, 805-811.	1.7	25
63	Speciation of phytate ion in aqueous solution. Cadmium(II) interactions in aqueous NaCl at different ionic strengths. <i>Analytical and Bioanalytical Chemistry</i> , 2006, 386, 346-356.	1.9	25
64	Activity coefficients, acid-base properties and weak Na ⁺ ion pair formation of some resorcinol derivatives. <i>Fluid Phase Equilibria</i> , 2010, 292, 71-79.	1.4	25
65	Total and Specific Solubility and Activity Coefficients of Neutral Species of (CH ₂) ₂ COOH ⁺ Complexons in Aqueous NaCl Solutions at Different Ionic Strengths, (0 to 5) mol·L ⁻¹ , and 298.15 K. <i>Journal of Chemical & Engineering Data</i> , 2011, 56, 437-443.	1.0	25
66	Speciation of low molecular weight carboxylic ligands in natural fluids: protonation constants and association with major components of seawater of oxydiacetic and citric acids. <i>Analytica Chimica Acta</i> , 1999, 398, 103-110.	2.6	24
67	Thermodynamic Parameters for the Protonation of Poly(allylamine) in Concentrated LiCl(aq) and NaCl(aq). <i>Journal of Chemical & Engineering Data</i> , 2004, 49, 658-663.	1.0	24
68	Solubility and activity coefficients of 2,2'-bipyridyl, 1,10-phenanthroline and 2,2',6',2''-terpyridine in NaCl(aq) at different ionic strengths and T=298.15K. <i>Fluid Phase Equilibria</i> , 2008, 272, 47-52.	1.4	24
69	Quantitative study on the interaction of Sn ²⁺ and Zn ²⁺ with some phosphate ligands, in aqueous solution at different ionic strengths. <i>Journal of Molecular Liquids</i> , 2012, 165, 143-153.	2.3	24
70	Speciation of tin(II) in aqueous solution: thermodynamic and spectroscopic study of simple and mixed hydroxocarboxylate complexes. <i>Monatshefte für Chemie</i> , 2013, 144, 761-772.	0.9	24
71	Acid-Base Properties, Solubility, Activity Coefficients and Na ⁺ Ion Pair Formation of Complexons in NaCl(aq) at Different Ionic Strengths. <i>Journal of Solution Chemistry</i> , 2013, 42, 1452-1471.	0.6	24
72	The calculation of equilibrium concentrations. ES4EC1: A FORTRAN program for computing distribution diagrams and titration curves. <i>Computers & Chemistry</i> , 1989, 13, 343-359.	1.2	23

#	ARTICLE	IF	CITATIONS
73	Thermodynamic parameters for the binding of ATP by protonated open-chain polyamines. Journal of the Chemical Society, Faraday Transactions, 1998, 94, 1091-1095.	1.7	23
74	Salt effects on the protonation of imidazole in aqueous solution at different ionic strengths: A tentative explanation by a complex formation model. Journal of Solution Chemistry, 1989, 18, 23-36.	0.6	22
75	Modeling the Dependence on Medium and Ionic Strength of Glutathione Acid-Base Behavior in LiClaq, NaClaq, KClaq, RbClaq, CsClaq, (CH ₃) ₄ NClaq, and (C ₂ H ₅) ₄ NClaq. Journal of Chemical & Engineering Data, 2007, 52, 1028-1036.	1.0	22
76	Thermodynamics of binary and ternary interactions in the tin(II)/phytate system in aqueous solutions, in the presence of Cl ⁻ or F ⁻ . Journal of Chemical Thermodynamics, 2012, 51, 88-96.	1.0	22
77	Thermodynamic properties of melamine (2,4,6-triamino-1,3,5-triazine) in aqueous solution. Effect of ionic medium, ionic strength and temperature on the solubility and acid-base properties. Fluid Phase Equilibria, 2013, 355, 104-113.	1.4	22
78	Understanding the bioavailability and sequestration of different metal cations in the presence of a biodegradable chelant S,S-EDDS in biological fluids and natural waters. Chemosphere, 2016, 150, 341-356.	4.2	22
79	Thermodynamics of formation of magnesium(II), calcium(II), strontium(II) and barium(II) succinate complexes in aqueous solution. Thermochemica Acta, 1984, 80, 197-208.	1.2	21
80	Ionic strength dependence of formation constants. Part 7. Protonation constants of low molecular weight carboxylic acids at 10, 25 and 45°C. Thermochemica Acta, 1985, 86, 273-280.	1.2	21
81	Quantitative study of the interactions of ATP with amines and amino acids. Journal of the Chemical Society, Faraday Transactions, 1996, 92, 1511-1518.	1.7	21
82	Speciation of polyelectrolytes in natural fluids Protonation and interaction of polymethacrylates with major components of seawater. Talanta, 2002, 58, 405-417.	2.9	21
83	Speciation of phytate ion in aqueous solution. Dimethyltin(IV) interactions in NaClaq at different ionic strengths. Biophysical Chemistry, 2005, 116, 111-120.	1.5	21
84	Interaction of Phytate with Ag ⁺ , CH ₃ Hg ⁺ , Mn ²⁺ , Fe ²⁺ , Co ²⁺ , and VO ²⁺ : Stability Constants and Sequestering Ability. Journal of Chemical & Engineering Data, 2012, 57, 2838-2847.	1.0	21
85	Binding of polyanions by biogenic amines. II. Formation and stability of protonated putrescine and cadaverine complexes with carboxylic ligands. Talanta, 1998, 46, 1079-1084.	2.9	20
86	Speciation of organic matter in natural waters interaction of polyacrylates and polymethacrylates with major cation components of seawater. Marine Chemistry, 2004, 86, 33-44.	0.9	20
87	Sit Parameters for 1:2 Electrolytes and Correlation with Pitzer Coefficients. Annali Di Chimica, 2007, 97, 85-95.	0.6	20
88	Potentiometric and spectrophotometric characterization of the UO ₂ ²⁺ -citrate complexes in aqueous solution, at different concentrations, ionic strengths and supporting electrolytes. Radiochimica Acta, 2012, 100, 13-28.	0.5	20
89	Interaction of Alkyltin(IV) Compounds with Ligands of Interest in the Speciation of Natural fluids: Complexes of (CH ₃) ₂ Sn ²⁺ with Carboxylates. Applied Organometallic Chemistry, 1997, 11, 683-691.	1.7	19
90	pK _a and T _{1/2} ^o charge relationships for the binding of carboxylic anions by open-chain polyammonium cations. Journal of the Chemical Society, Faraday Transactions, 1998, 94, 2395-2398.	1.7	19

#	ARTICLE	IF	CITATIONS
91	Binding of polyanions by biogenic amines. III. Formation and stability of protonated spermidine and spermine complexes with carboxylic ligands. <i>Talanta</i> , 1999, 48, 119-126.	2.9	19
92	Speciation of low molecular weight ligands in natural fluids: protonation constants and association of open chain polyamines with the major components of seawater. <i>Analytica Chimica Acta</i> , 2000, 418, 43-51.	2.6	19
93	Dissociation Constants of Protonated Cysteine Species in NaCl Media. <i>Journal of Solution Chemistry</i> , 2002, 31, 783-792.	0.6	19
94	Speciation of organotin compounds in NaCl aqueous solution: interaction of mono-, di- and tri-organotin(IV) cations with nucleotide 5' monophosphates. <i>Applied Organometallic Chemistry</i> , 2004, 18, 653-661.	1.7	19
95	Dioxouranium(VI) " Carboxylate Complexes. Interaction with Dicarboxylic Acids in Aqueous Solution: Speciation and Structure. <i>Annali Di Chimica</i> , 2006, 96, 399-420.	0.6	19
96	Modeling of Protonation Constants of Linear Aliphatic Dicarboxylates Containing -S-Groups in Aqueous Chloride Salt Solutions, at Different Ionic Strengths, Using the SIT and Pitzer Equations and Empirical Relationships. <i>Journal of Solution Chemistry</i> , 2008, 37, 763-784.	0.6	19
97	Formation and stability of mixed Mg ²⁺ /Ca ²⁺ /phytate species in synthetic seawater media. <i>Marine Chemistry</i> , 2008, 112, 142-148.	0.9	19
98	Modeling the acid-base properties of molybdate(VI) in different ionic media, ionic strengths and temperatures, by EDH, SIT and Pitzer equations. <i>Journal of Molecular Liquids</i> , 2017, 229, 15-26.	2.3	19
99	Speciation of poly-amino carboxylic compounds in seawater. <i>Chemical Speciation and Bioavailability</i> , 2003, 15, 75-86.	2.0	18
100	Interaction of UO ₂ ²⁺ with ATP in aqueous ionic media. <i>Biophysical Chemistry</i> , 2005, 117, 147-153.	1.5	18
101	Speciation of phytate ion in aqueous solution. Protonation in CsClaq at different ionic strengths and mixing effects in LiClaq+CsClaq. <i>Journal of Molecular Liquids</i> , 2008, 138, 76-83.	2.3	18
102	Alkali Metal Ion Complexes with Phosphates, Nucleotides, Amino Acids, and Related Ligands of Biological Relevance. Their Properties in Solution. <i>Metal Ions in Life Sciences</i> , 2016, 16, 133-166.	2.8	18
103	Protonation Constants and Association of Polycarboxylic Ligands with the Major Components of Seawater. <i>Journal of Chemical & Engineering Data</i> , 2000, 45, 996-1000.	1.0	17
104	Critical Evaluation of Protonation Constants. Literature Analysis and Experimental Potentiometric and Calorimetric Data for the Thermodynamics of Phthalate Protonation in Different Ionic Media. <i>Journal of Solution Chemistry</i> , 2006, 35, 1227-1244.	0.6	17
105	Thermodynamic data for lanthanoid(III) sequestration by phytate at different temperatures. <i>Monatshfte für Chemie</i> , 2010, 141, 511-520.	0.9	17
106	On the complexation of metal cations with "pure" diethylenetriamine-N,N,N',N'-tetraacetate-pentakis(methylenephosphonic) acid. <i>New Journal of Chemistry</i> , 2017, 41, 4065-4075.		17
107	Interactions of diethylenetriaminepentaacetic acid (dtpa) and triethylenetetraaminehexaacetic acid (ttha) with major components of natural waters. <i>Analytical and Bioanalytical Chemistry</i> , 2003, 375, 956-967.	1.9	16
108	SIT Parameters for the Dependence of (Poly)carboxylate Activity Coefficients on Ionic Strength in (C ₂ H ₄) ₄ NI _{aq} (0.12 mol·kg ⁻¹) and (CH ₃) ₄ NCl _{aq} (0.39 mol·kg ⁻¹) in the Temperature Range 278 K to 328 K and Correlation with Pitzer Parameters. <i>Journal of Chemical & Engineering Data</i> , 2007, 52, 2195-2203.	1.0	16

#	ARTICLE	IF	CITATIONS
109	On the Complexation of Cu(II) and Cd(II) With Polycarboxyl Ligands. Potentiometric Studies With ISE-H ⁺ , ISE-Cu ²⁺ , and ISE-Cd ²⁺ . Journal of Chemical & Engineering Data, 2010, 55, 714-722.	1.0	16
110	Solubility, activity coefficients and acid-base properties of three naphthol derivatives in NaCl(aq) at different ionic strengths and at T=298.15K. Journal of Molecular Liquids, 2011, 158, 50-56.	2.3	16
111	SALMO and S ₃ M: A Saliva Model and a Single Saliva Salt Model for Equilibrium Studies. Bioinorganic Chemistry and Applications, 2015, 2015, 1-12.	1.8	16
112	Thermodynamic solution properties of a biodegradable chelant (MGDA) and its interaction with the major constituents of natural fluids. Fluid Phase Equilibria, 2017, 434, 63-73.	1.4	16
113	Equilibrium studies in natural fluids: interactions of -PO ₄ ³⁻ , -P ₂ O ₇ ⁴⁻ and -P ₃ O ₁₀ ⁵⁻ with the major constituents of sea water. Chemical Speciation and Bioavailability, 1998, 10, 19-26.	2.0	15
114	Investigations on ancient mortars from the Basilian monastery of Fragal. Journal of Thermal Analysis and Calorimetry, 2008, 91, 477-485.	2.0	15
115	Sequestering ability of phytate towards protonated BPEI and other polyammonium cations in aqueous solution. Biophysical Chemistry, 2008, 136, 108-114.	1.5	15
116	Zinc(II) complexes with hydroxocarboxylates and mixed metal species with tin(II) in different salts aqueous solutions at different ionic strengths: formation, stability, and weak interactions with supporting electrolytes. Monatshefte für Chemie, 2015, 146, 527-540.	0.9	15
117	Complexation of environmentally and biologically relevant metals with bifunctional 3-hydroxy-4-pyridinones. Journal of Molecular Liquids, 2020, 319, 114349.	2.3	15
118	EQUILIBRIUM STUDIES IN NATURAL WATERS: SPECIATION OF PHENOLIC COMPOUNDS IN SYNTHETIC SEAWATER AT DIFFERENT SALINITIES. Environmental Toxicology and Chemistry, 1995, 14, 767.	2.2	15
119	Mixed proton complexes of aminoacids and carboxylic ligands in aqueous solution. Talanta, 1993, 40, 629-635.	2.9	14
120	Formation and stability of proton-amine-inorganic anion complexes in aqueous solution. Journal of Solution Chemistry, 1995, 24, 325-341.	0.6	14
121	Salt effects on the protonation of polymethacrylate and Na ⁺ , K ⁺ , Ca ²⁺ complex formation. Fluid Phase Equilibria, 1999, 163, 127-137.	1.4	14
122	Interaction of Polyamines with Mg ²⁺ and Ca ²⁺ . Journal of Chemical & Engineering Data, 1999, 44, 744-749.	1.0	14
123	Complexes of Azelaic and Diethylenetrioxydiacetic Acids with Na ⁺ , Mg ²⁺ , and Ca ²⁺ in NaCl Aqueous Solutions, at 25 °C. Journal of Chemical & Engineering Data, 2000, 45, 15-19.	1.0	14
124	Binding of acrylic and sulphonic polyanions by open-chain polyammonium cations. Talanta, 2001, 53, 1241-1248.	2.9	14
125	Dissociation constants of protonated cysteine species in seawater media. Marine Chemistry, 2006, 99, 52-61.	0.9	14
126	Uranium(VI) sequestration by polyacrylic and fulvic acids in aqueous solution. Journal of Radioanalytical and Nuclear Chemistry, 2011, 289, 689-697.	0.7	14

#	ARTICLE	IF	CITATIONS
127	Solubility, Activity Coefficients, and Protonation Sequence of Risedronic Acid. Journal of Chemical & Engineering Data, 2014, 59, 3728-3740.	1.0	14
128	Thermodynamics for Proton Binding of Pyridine in Different Ionic Media at Different Temperatures. Journal of Chemical & Engineering Data, 2014, 59, 143-156.	1.0	14
129	Solubility and modeling acid-base properties of adrenaline in NaCl aqueous solutions at different ionic strengths and temperatures. European Journal of Pharmaceutical Sciences, 2015, 78, 37-46.	1.9	14
130	Thermodynamic Properties of O-Donor Polyelectrolytes: Determination of the Acid-Base and Complexing Parameters in Different Ionic Media at Different Temperatures. Journal of Chemical & Engineering Data, 2017, 62, 2676-2688.	1.0	14
131	A new bis-(3-hydroxy-4-pyridinone)-DTPA-derivative: Synthesis, complexation of di-/tri-valent metal cations and in vivo M3+ sequestering ability. Journal of Molecular Liquids, 2019, 281, 280-294.	2.3	14
132	Speciation Studies of Bifunctional 3-Hydroxy-4-Pyridinone Ligands in the Presence of Zn ²⁺ at Different Ionic Strengths and Temperatures. Molecules, 2019, 24, 4084.	1.7	14
133	Understanding the Solution Behavior of Epinephrine in the Presence of Toxic Cations: A Thermodynamic Investigation in Different Experimental Conditions. Molecules, 2020, 25, 511.	1.7	14
134	Caffeic Acid/Eu(III) Complexes: Solution Equilibrium Studies, Structure Characterization and Biological Activity. International Journal of Molecular Sciences, 2022, 23, 888.	1.8	14
135	Protonation thermodynamics of 2,2'-bipyridyl in aqueous solution. Salt effects and weak complex formation. Thermochimica Acta, 1993, 214, 325-338.	1.2	13
136	Dissociation constants of protonated methionine species in NaCl media. Biophysical Chemistry, 2003, 105, 79-87.	1.5	13
137	Dissociation Constants for Citric Acid in NaCl and KCl Solutions and their Mixtures at 25 °C. Journal of Solution Chemistry, 2004, 33, 1349-1366.	0.6	13
138	Speciation of Al ³⁺ in fairly concentrated solutions (20–200 mmol L ⁻¹) at I=1 mol L ⁻¹ (NaNO ₃), in the acidic pH range, at different temperatures. Chemical Speciation and Bioavailability, 2011, 23, 33-37.	2.0	13
139	Protonation Constants, Activity Coefficients, and Chloride Ion Pair Formation of Some Aromatic Amino-Compounds in NaCl (0 mol·kg ⁻¹ to 3 mol·kg ⁻¹) at T = 298.15 K. Journal of Chemical & Engineering Data, 2012, 57, 1851-1859.		13
140	Formation, stability and empirical relationships for the binding of Sn ²⁺ by O-, N- and S-donor ligands. Journal of Molecular Liquids, 2014, 200, 329-339.	2.3	13
141	Thermodynamic Data for the Modeling of Lanthanoid(III) Sequestration by Reduced Glutathione in Aqueous Solution. Journal of Chemical & Engineering Data, 2015, 60, 192-201.	1.0	13
142	Sequestration of HEDPA, NTA and phosphonic NTA derivatives towards Al ³⁺ in aqueous solution. Journal of Molecular Liquids, 2018, 261, 96-106.	2.3	13
143	Bifunctional 3-hydroxy-4-pyridinones as effective aluminium chelators: synthesis, solution equilibrium studies and in vivo evaluation. Journal of Inorganic Biochemistry, 2018, 186, 116-129.	1.5	13
144	Thermodynamic parameters for the formation of calcium α -alaninate complexes in aqueous solution. Thermochimica Acta, 1989, 140, 59-66.	1.2	12

#	ARTICLE	IF	CITATIONS
145	Interaction of methyltin(IV) compounds with carboxylate ligands. Part 2: formation thermodynamic parameters, predictive relationships and sequestering ability. Applied Organometallic Chemistry, 2008, 22, 30-38.	1.7	12
146	Sequestering Ability of Dicarboxylic Ligands Towards Dioxouranium(VI) in NaCl and KNO ₃ Aqueous Solutions at T=298.15 K. Journal of Solution Chemistry, 2009, 38, 1343-1356.	0.6	12
147	Dissociation Constants of Protonated Oxidized Glutathione in Seawater Media at Different Salinities. Aquatic Geochemistry, 2010, 16, 447-466.	1.5	12
148	Thermodynamic Study on the Protonation and Complexation of GLDA with Ca ²⁺ and Mg ²⁺ at Different Ionic Strengths and Ionic Media at 298.15 K. Journal of Chemical & Engineering Data, 2016, 61, 1895-1903.	1.0	12
149	Potentiometric, UV and ¹ H NMR study on the interaction of penicillin derivatives with Zn(II) in aqueous solution. Biophysical Chemistry, 2017, 223, 1-10.	1.5	12
150	New bis-(3-hydroxy-4-pyridinone)-NTA-derivative: Synthesis, binding ability towards Ca ²⁺ , Cu ²⁺ , Zn ²⁺ , Al ³⁺ , Fe ³⁺ and biological assays. Journal of Molecular Liquids, 2018, 272, 609-624.	2.3	12
151	Hydrolysis of dioxouranium(VI): a calorimetric study in NaCl _{aq} and NaClO _{4aq} , at 25°C. Thermochimica Acta, 2004, 414, 185-189.	1.2	11
152	Dioxouranium(VI)-Carboxylate Complexes. Interaction of UO_2^{2+} with 1,2,3,4,5,6-Benzenehexacarboxylate (Mellitate) in 0.1 mol/L (NaCl aq) at 1.0 mol/L. Journal of Solution Chemistry, 2007, 36, 479-496.	0.6	11
153	Sequestration of Alkyltin(IV) Compounds in Aqueous Solution: Formation, Stability, and Empirical Relationships for the Binding of Dimethyltin(IV) Cation by N- and O-Donor Ligands. Bioinorganic Chemistry and Applications, 2009, 2009, 1-17.	1.8	11
154	Palladium(II) Complexes of Aminopolycarboxylic Ligands in Aqueous Solution. Journal of Chemical & Engineering Data, 2011, 56, 4759-4771.	1.0	11
155	Hydrolysis of Monomethyl-, Dimethyl-, and Trimethyltin(IV) Cations in Fairly Concentrated Aqueous Solutions at $I = 1 \text{ mol}\cdot\text{L}^{-1}$ (NaNO ₃) and $T = 298.15 \text{ K}$. Evidence for the Predominance of Polynuclear Species. Journal of Chemical & Engineering Data, 2011, 56, 1108-1115.	1.0	11
156	Studies on polyfunctional O-ligands. Solubility and thermal stability of 1,2,4,5-benzenetetracarboxylate complexes with alkali and alkaline earth metal ions. Thermochimica Acta, 1990, 173, 25-41.	1.2	10
157	Equilibrium studies in natural waters: Speciation of phenolic compounds in synthetic seawater at different salinities. Environmental Toxicology and Chemistry, 1995, 14, 767-773.	2.2	10
158	Binding of glyphosate by open-chain polyammonium cations. Environmental Toxicology and Chemistry, 1999, 18, 2131-2137.	2.2	10
159	The calculation of equilibrium concentrations. Estime: A computer program for comparing execution times on different machines. Computers & Chemistry, 1988, 12, 305-315.	1.2	9
160	Salt effects on the protonation of oxalate in aqueous NaCl, KCl and tetraethylammonium iodide solution at $5 \leq T \leq 50^\circ\text{C}$ and $0.1 \leq I \leq 1$. 1 mol dm ⁻³ . Thermochimica Acta, 1992, 202, 133-149.	1.2	9
161	Stability-Charge and Stability-Structure Relationships in the Binding of Dicarboxylic Ligands by Open-Chain Polyammonium Cations. Journal of Chemical & Engineering Data, 2000, 45, 717-723.	1.0	9
162	Protonation thermodynamics of some aminophenol derivatives in NaCl(aq) ($0.1 \leq I \leq 3 \text{ mol}\cdot\text{kg}^{-1}$) at T=298.15K. Journal of Chemical Thermodynamics, 2012, 44, 154-162.	1.0	9

#	ARTICLE	IF	CITATIONS
163	Thermodynamic Parameters for the Interaction of Amoxicillin and Ampicillin with Magnesium in NaCl Aqueous Solution, at Different Ionic Strengths and Temperatures. <i>Journal of Chemical & Engineering Data</i> , 2017, 62, 1018-1027.	1.0	9
164	Complexing ability of pesticides and related compounds. thermodynamic parameters for the formation of H ⁺ , Na ⁺ and Ca ²⁺ complexes with 2,4-dichlorophenoxyacetate in aqueous solution at different temperatures and ionic strengths. <i>Thermochimica Acta</i> , 1988, 128, 261-266.	1.2	8
165	Mixed aminocarboxylic ligand complexes. <i>Journal of the Chemical Society, Faraday Transactions</i> , 1995, 91, 1619.	1.7	8
166	Medium effects on the protonation enthalpies of linear diamines in NaCl aqueous solutions at 25°C. <i>Thermochimica Acta</i> , 2000, 363, 29-35.	1.2	8
167	Thermodynamics of Proton Association of Polyacrylates and Polymethacrylates in NaCl(aq). <i>Journal of Solution Chemistry</i> , 2003, 32, 967-976.	0.6	8
168	Sequestration of (CH ₃)Hg ⁺ by amino-polycarboxylic chelating agents. <i>Journal of Molecular Liquids</i> , 2012, 172, 46-52.	2.3	8
169	The effect of the tetraalkylammonium salts on the protonation thermodynamics of the phytate anion. <i>Fluid Phase Equilibria</i> , 2014, 383, 126-133.	1.4	8
170	On the interaction of phytate with proton and monocharged inorganic cations in different ionic media, and modeling of acid-base properties at low ionic strength. <i>Journal of Chemical Thermodynamics</i> , 2015, 90, 51-58.	1.0	8
171	Thermodynamics (Solubility and Protonation Constants) of Risedronic Acid in Different Media and Temperatures (283.15–318.15 K). <i>Journal of Solution Chemistry</i> , 2017, 46, 1903-1927.	0.6	8
172	Solubility, acid-base properties and thermodynamics of interaction between three NTA-phosphonate derivatives and the main cationic components (H ⁺ , Na ⁺ , Mg ²⁺ and Ca ²⁺) of natural fluids. <i>Journal of Chemical Thermodynamics</i> , 2018, 123, 117-127.	1.0	8
173	Chemical speciation of organic matter in natural waters. Interaction of nucleotide 5' mono-, di- and triphosphates with major components of seawater. <i>Chemical Speciation and Bioavailability</i> , 2004, 16, 1-8.	2.0	7
174	Dioxouranium(VI)-Carboxylate Complexes. Speciation of UO ₂ ²⁺ -1,2,3-Propanetricarboxylate System in NaCl(aq) at Different Ionic Strengths and at T = 25 °C. <i>Annali Di Chimica</i> , 2007, 97, 163-175.	0.6	7
175	Effect of Ionic Strength and Temperature on the Protonation of Oxidized Glutathione. <i>Journal of Solution Chemistry</i> , 2008, 37, 1245-1259.	0.6	7
176	Speciation of chitosan-phosphate and chitosan-nucleotide systems in NaCl aqueous solution. <i>Chemical Speciation and Bioavailability</i> , 2010, 22, 99-107.	2.0	7
177	Sequestration of some biogenic amines and poly(allyl)amine by high molecular weight polycarboxylic ligands in aqueous solution. <i>Journal of Molecular Liquids</i> , 2010, 151, 138-144.	2.3	7
178	Understanding the bioavailability and sequestration of different metal cations in the presence of a biodegradable chelant MGDA in biological fluids and natural waters. <i>Chemosphere</i> , 2017, 183, 107-118.	4.2	7
179	Complexing ability of pesticides and related compounds. Thermodynamic parameters for the formation of copper(II) complexes of phenoxyacetate, 2-phenoxy. <i>Thermochimica Acta</i> , 1989, 149, 199-204.	1.2	6
180	Mixed-metal complexes in solution. A potentiometric and spectrophotometric study of copper(II)-nickel(II)-L-hydroxylysine complexes in aqueous solution. <i>Journal of the Chemical Society Dalton Transactions</i> , 1989, , 1745-1749.	1.1	6

#	ARTICLE	IF	CITATIONS
181	Binding of hexacyanoferrate(II) by aliphatic amines in aqueous solution. <i>Journal of Solution Chemistry</i> , 1996, 25, 155-165.	0.6	6
182	Interaction of l-malic acid with alkaline metals and open chain polyammonium cations in aqueous solution. <i>Talanta</i> , 2001, 54, 25-36.	2.9	6
183	Chemical speciation of nucleotide 5â€²-monophosphates in the presence of biogenic amines. <i>Chemical Speciation and Bioavailability</i> , 2001, 13, 113-119.	2.0	6
184	Binding of fluoride and carbonate by open chain polyammonium cations. <i>Talanta</i> , 2004, 64, 510-517.	2.9	6
185	Additivity Factors in the Binding of the Diethyltin(IV) Cation by Ligands Containing Amino and Carboxylic Groups at Different Ionic Strengths. <i>Journal of Solution Chemistry</i> , 2005, 34, 1211-1226.	0.6	6
186	Sequestration of biogenic amines by alginic and fulvic acids. <i>Biophysical Chemistry</i> , 2006, 122, 221-231.	1.5	6
187	Sequestration of organometallic compounds by synthetic and naturally occurring polycarboxylate ligands. Binding of monomethylmercury(II) by polyacrylic and alginic acids. <i>Chemical Speciation and Bioavailability</i> , 2007, 19, 129-140.	2.0	6
188	Speciation of chitosan with low and high molecular weight carboxylates in aqueous solution. <i>Chemical Speciation and Bioavailability</i> , 2009, 21, 81-91.	2.0	6
189	Medium Effect on the Acid-Base Properties of Branched Polyethylenimine in Different Aqueous Electrolyte Solutions. <i>Journal of Chemical & Engineering Data</i> , 2009, 54, 502-510.	1.0	6
190	Thermodynamic study of the non covalent interactions of phytate with xanthine derivatives and histamine in aqueous solution. <i>Journal of Molecular Liquids</i> , 2013, 178, 37-43.	2.3	6
191	Thermodynamics for proton binding of phytate in KNO ₃ (aq) at different temperatures and ionic strengths. <i>Thermochimica Acta</i> , 2013, 566, 193-202.	1.2	6
192	Enhancement of Hydrolysis through the Formation of Mixed Heterometal Species: Al ³⁺ /CH ₃ Sn ³⁺ Mixtures. <i>Journal of Chemical & Engineering Data</i> , 2013, 58, 821-826.	1.0	6
193	Evaluation of the sequestering ability of different complexones towards Ag ⁺ ion. <i>Journal of Molecular Liquids</i> , 2014, 199, 432-439.	2.3	6
194	Sequestering Ability of Aminopolycarboxylic (APCs) and Aminopolyphosphonic (APPs) Ligands Toward Palladium(II) in Aqueous Solution. <i>Journal of Chemical & Engineering Data</i> , 2014, 59, 1970-1983.	1.0	6
195	Acid-Base and Thermodynamic Properties of α -Gluconic Acid and Its Interaction with Sn ²⁺ and Zn ²⁺ . <i>Journal of Chemical & Engineering Data</i> , 2016, 61, 2040-2051.	1.0	6
196	Prediction of water solubility and Setschenow coefficients by tree-based regression strategies. <i>Journal of Molecular Liquids</i> , 2019, 282, 401-406.	2.3	6
197	Thermodynamic study on polyaspartic acid biopolymer in solution and prediction of its chemical speciation and bioavailability in natural fluids. <i>Journal of Molecular Liquids</i> , 2019, 274, 68-76.	2.3	6
198	Complexing ability of pesticides and related compounds. Thermodynamic parameters for the formation of zinc(ii) complexes (and other divalent cations) of phenoxyacetate, 2-phenoxypropionate and 2-phenoxybutyrate in aqueous solution at different temperatures and ionic strengths. <i>Thermochimica Acta</i> , 1990, 162, 449-451.	1.2	5

#	ARTICLE	IF	CITATIONS
199	Speciation of N-(phosphonomethyl)glycine (glyphosate) in aqueous solutions containing major components of natural fluids. <i>Chemical Speciation and Bioavailability</i> , 1996, 8, 59-65.	2.0	5
200	Binding of Phosphate, Pyrophosphate, and Hexacyanoferrate(II) by Fully N-Methyl Substituted Polyammonium Cations in Aqueous Solution. <i>Journal of Chemical & Engineering Data</i> , 2004, 49, 133-137.	1.0	5
201	Mixing Effects on the Protonation of Polycarboxylates. Protonation of Benzenehexacarboxylate in LiCl ⁺ KCl, NaCl ⁺ KCl, NaCl ⁺ LiCl, and LiCl ⁺ CsCl Aqueous Solutions at $I = 1 \text{ mol}\cdot\text{L}^{-1}$ and $T = 298.15 \text{ K}$. <i>Journal of Chemical & Engineering Data</i> , 2009, 54, 2137-2139.	1.0	5
202	Risedronate complexes with Mg ²⁺ , Zn ²⁺ , Pb ²⁺ , and Cu ²⁺ : Species thermodynamics and sequestering ability in NaCl(aq) at different ionic strengths and at $T = 298.15 \text{ K}$. <i>Journal of Molecular Liquids</i> , 2021, 343, 117699.	2.3	5
203	Formation and stability of magnesium-oxodiacetate complexes at different temperatures and ionic strengths in aqueous solution. <i>Thermochimica Acta</i> , 1987, 117, 317-323.	1.2	4
204	Dissociation constants of protonated methionine species in seawater media. <i>Marine Chemistry</i> , 2007, 106, 463-470.	0.9	4
205	Sequestration of alkyltin(IV) cations by complexation with amino-polycarboxylic chelating agents. <i>Journal of Molecular Liquids</i> , 2013, 187, 74-82.	2.3	4
206	Phytate-molybdate interactions in NaCl(aq) at different ionic strengths: unusual behaviour of the protonated species. <i>New Journal of Chemistry</i> , 2018, 42, 7671-7679.	1.4	4
207	Use of Gantrez Copolymers as Potential Chelating Agent for the Selective Sequestration of Metal Ions. Studies of the Interactions in Aqueous Solution at Different Ionic Strengths and Temperatures. <i>Journal of Chemical & Engineering Data</i> , 2018, 63, 4193-4204.	1.0	4
208	Thermodynamic Study on the Interaction of Nicotinic Acid with H ⁺ , Na ⁺ , Ca ²⁺ and Mg ²⁺ at Different Temperatures and Ionic Strengths. <i>Journal of Solution Chemistry</i> , 2019, 48, 1671-1684.	0.6	4
209	8-Hydroxyquinoline-2-Carboxylic Acid as Possible Molybdophore: A Multi-Technique Approach to Define Its Chemical Speciation, Coordination and Sequestering Ability in Aqueous Solution. <i>Biomolecules</i> , 2020, 10, 930.	1.8	4
210	The Solution Behavior of Dopamine in the Presence of Mono and Divalent Cations: A Thermodynamic Investigation in Different Experimental Conditions. <i>Biomolecules</i> , 2021, 11, 1312.	1.8	4
211	SCALPO, a computer program for simulating calorimetric and potentiometric titration curves. <i>Analytica Chimica Acta</i> , 1989, 222, 385-390.	2.6	3
212	Thermodynamic Parameters for the Formation of Pyrophosphate-protonated Polyamine Complexes. <i>Journal of Chemical Research Synopses</i> , 1998, , 480-481.	0.3	3
213	Title is missing!. <i>Journal of Solution Chemistry</i> , 2000, 29, 1101-1114.	0.6	3
214	Stability-Charge and Stability-Structure Relationships in the Binding of Tri- and Tetracarboxylic Ligands by Open-Chain Polyammonium Cations. <i>Journal of Chemical & Engineering Data</i> , 2001, 46, 1365-1370.	1.0	3
215	Modelling the separation of amines by high performance liquid chromatography. <i>Analytica Chimica Acta</i> , 2001, 436, 333-342.	2.6	3
216	Quantitative Study of the Interaction between ATP and Aromatic Amines in Aqueous Solution. <i>Journal of Solution Chemistry</i> , 2012, 41, 1240-1253.	0.6	3

#	ARTICLE	IF	CITATIONS
217	Thermodynamic study on the protonation of glycine in different (water+1-butyl-3-methylimidazolium) Tj ETQq1 1 0.784314 rgBT /Overl 163-169.	1.0	3
218	Thermodynamics of Zn ²⁺ 2-mercaptopyridine-N-oxide and 2-hydroxypyridine-N-oxide interactions: Stability, solubility, activity coefficients and medium effects. Journal of Molecular Liquids, 2015, 211, 876-884.	2.3	3
219	Exploring various ligand classes for the efficient sequestration of stannous cations in the environment. Science of the Total Environment, 2018, 643, 704-714.	3.9	3
220	Thermodynamic Behavior of Polyalcohols and Speciation Studies in the Presence of Divalent Metal Cations. Journal of Chemical & Engineering Data, 2020, 65, 2805-2812.	1.0	3
221	Thermodynamic Solution Properties of a Biodegradable Chelant (L-glutamic-N,N-diacetic Acid, L-GLDA) and Its Sequestering Ability toward Cd ²⁺ . Molecules, 2021, 26, 7087.	1.7	3
222	Bifunctional 3-Hydroxy-4-Pyridinones as Potential Selective Iron(III) Chelators: Solution Studies and Comparison with Other Metals of Biological and Environmental Relevance. Molecules, 2021, 26, 7280.	1.7	3
223	The Effect of Metal Cations on the Aqueous Behavior of Dopamine. Thermodynamic Investigation of the Binary and Ternary Interactions with Cd ²⁺ , Cu ²⁺ and UO ₂ ²⁺ in NaCl at Different Ionic Strengths and Temperatures. Molecules, 2021, 26, 7679.	1.7	3
224	Thermodynamic parameters for the protonation of urea at different temperatures and ionic strengths. Journal of Solution Chemistry, 1992, 21, 397-404.	0.6	2
225	Evaluation of behaviour of linear monoamines CH ₃ (CH ₂) _n NH ₂ (n=1-6) in ion chromatography. Analytica Chimica Acta, 2003, 477, 41-48.	2.6	2
226	Thermodynamic Study on the Protonation and Complexation of the Neuroleptic Drug, Gabapentin with Na ⁺ , Ca ²⁺ and Mg ²⁺ at Various Temperatures and Ionic Strengths. Journal of Solution Chemistry, 2020, 49, 1225-1236.	0.6	2
227	Modelling the Hydrolysis of Mixed Mono-, Di- and Trimethyltin(IV) Complexes in Aqueous Solutions. Journal of Solution Chemistry, 2015, 44, 1611-1625.	0.6	1
228	Polycarboxylic acids in sea water: acid-base properties, solubilities, activity coefficients, and complex formation constants at different salinities. Monatshefte für Chemie, 2016, 147, 1481-1505.	0.9	1
229	Thermodynamic Study on the Protonation and Na ⁺ , Ca ²⁺ , Mg ²⁺ -Complexation of a Biodegradable Chelant (HEIDA) at Different Ionic Strengths and Temperatures. Journal of Solution Chemistry, 2018, 47, 528-543.	0.6	1
230	Complexation of Molybdenum(VI) with GLDA at Different Ionic Strengths. Journal of Solution Chemistry, 2018, 47, 1965-1979.	0.6	1
231	Characterization of the thermodynamic properties of some benzenepolycarboxylic acids: Acid-base properties, weak complexes, total and neutral species solubility, solubility products in NaCl _{aq} , (CH ₃) ₄ NCl _{aq} and Synthetic Sea Water (SSW). Fluid Phase Equilibria, 2019, 480, 41-52.	1.4	1
232	Nature as Resource. Thermodynamic characterization of natural and synthetic polymers and their sequestering ability towards some bivalent metal cations. Journal of Chemical Thermodynamics, 2020, 150, 106205.	1.0	1
233	Towards a rational design of materials for the removal of environmentally relevant cations: polymer inclusion membranes (PIMs) and surface-modified PIMs for Sn ²⁺ sequestration in aqueous solution. Environmental Science and Pollution Research, 2021, 28, 51072-51087.	2.7	1
234	Behavior of Antibacterial Ofloxacin; Hydration Constants and Solubility in Aqueous Solutions of Sodium Chloride at Different Temperatures. Journal of Solution Chemistry, 2021, 50, 1236-1257.	0.6	1