Gabriela F De Larrañaga

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3605990/publications.pdf

Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Clinical and therapeutic value of the adjusted Global Antiphospholipid Syndrome Score in primary obstetric antiphospholipid syndrome. Lupus, 2022, 31, 354-362. | 1.6 | 5 |
| 2 | A focus on the roles of histones in health and diseases. Clinical Biochemistry, 2021, 94, 12-19. | 1.9 | 11 |
| 3 | Risk factors for early severe preeclampsia in obstetric antiphospholipid syndrome with conventional treatment. The impact of hydroxychloroquine. Lupus, 2020, 29, 1736-1742. | 1.6 | 9 |
| 4 | A high-risk laboratory profile of antiphospholipid antibodies and thrombosis is associated with a large number of extra-criteria manifestations in obstetric antiphospholipid syndrome. Immunologic Research, 2019, 67, 478-485. | 2.9 | 11 |
| 5 | Trombofilia hereditaria y pérdidas de embarazo. Estudio de una cohorte de Argentina. Medicina ClÃnica, 2019, 152, 249-254. | 0.6 | 5 |
| 6 | Maternal carriers of the ANXA5 M2 haplotype are exposed to a greater risk for placenta-mediated pregnancy complications. Journal of Assisted Reproduction and Genetics, 2018, 35, 921-928. | 2.5 | 11 |
| 7 | What is the best time to assess the antiphospholipid antibodies (aPL) profile to better predict the obstetric outcome in antiphospholipid syndrome (APS) patients?. Immunologic Research, 2018, 66, 577-583. | 2.9 | 6 |
| 8 | Pregnancy failure in patients with obstetric antiphospholipid syndrome with conventional treatment: the influence of a triple positive antibody profile. Lupus, 2017, 26, 983-988. | 1.6 | 48 |
| 9 | Decreased Levels of Circulatig Protein S in Patients with Active Crohn's Disease. Gastroenterology, 2017, 152, S612. | 1.3 | 0 |
| 10 | Effect of gene–gene and gene–environment interactions associated with antituberculosis drug-induced hepatotoxicity. Pharmacogenetics and Genomics, 2017, 27, 363-371. | 1.5 | 12 |
| 11 | The â^'308 G/A polymorphism in the tumor necrosis factor-α gene is not associated with development and progression of rheumatoid arthritis in Argentinean patients. International Journal of Rheumatic Diseases, 2016, 19, 476-481. | 1.9 | 7 |
| 12 | tagSNP rs1495741 as a useful molecular marker to predict antituberculosis drug-induced hepatotoxicity. Pharmacogenetics and Genomics, 2016, 26, 357-361. | 1.5 | 7 |
| 13 | Q222R polymorphism in the DNAse I gene is not associated with susceptibility to rheumatoid arthritis or to disease course in an Argentine patient cohort. Egyptian Rheumatologist, 2016, 38, 289-293. | 1.0 | 1 |
| 14 | D-Dimer as a prognostic marker of morbidity and mortality among HIV patients: a call for attention. Infectious Diseases, 2016, 48, 860-861. | 2.8 | 3 |
| 15 | The â^'2518ÂA/G polymorphism in the monocyte chemoattractant protein 1 gene (MCP-1) is associated with an increased risk of rheumatoid arthritis in Argentine patients. Clinical Rheumatology, 2016, 35, 3057-3061. | 2.2 | 8 |
| 16 | Paternal factor V Leiden and recurrent pregnancy loss: a new concept behind fetal genetics?: reply. Journal of Thrombosis and Haemostasis, 2014, 12, 1759-1760. | 3.8 | 2 |
| 17 | Paternal factor V Leiden and recurrent pregnancy loss: a new concept behind fetal genetics?. Journal of Thrombosis and Haemostasis, 2014, 12, 666-669. | 3.8 | 15 |
| 18 | Neutrophil Extracellular Traps in Sepsis. Shock, 2014, 42, 286-294. | 2.1 | 134 |

2

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | 4G/5G plasminogen activator inhibitor-1 and â^'308 A/G tumor necrosis factor-α promoter gene polymorphisms in Argentinean lupus patients: focus on lupus nephritis. Clinical and Experimental Medicine, 2014, 14, 83-89. | 3.6 | 7 |
| 20 | Sex, ethnicity, and slow acetylator profile are the major causes of hepatotoxicity induced by antituberculosis drugs. Journal of Gastroenterology and Hepatology (Australia), 2013, 28, 323-328. | 2.8 | 57 |
| 21 | Role of -675 4G/5G in the Plasminogen Activator Inhibitor-1 Gene and -308G/A Tumor Necrosis Factor-α Gene Polymorphisms in Obese Argentinean Patients. Genetic Testing and Molecular Biomarkers, 2012, 16, 372-375. | 0.7 | 3 |
| 22 | The -2518 A/G polymorphism in the monocyte chemoattractant protein 1 gene is associated with the risk of developing systemic lupus erythematosus in Argentinean patients: a multicenter study. European Cytokine Network, 2012, 23, 7-11. | 2.0 | 9 |
| 23 | Biomarkers in sepsis at time zero: intensive care unit scores, plasma measurements and polymorphisms in Argentina. Journal of Infection in Developing Countries, 2012, 6, 555-562. | 1.2 | 17 |
| 24 | The distribution of allelic and genotypic frequencies of N-Acetyltransferase-2 variants in an Argentine population. Journal of Infection in Developing Countries, 2012, 6, 671-674. | 1.2 | 10 |
| 25 | Antiâ€Î²2glycoprotein I antibodies from leprosy patients do not show thrombogenic effects in an in vivo animal model. Journal of Thrombosis and Haemostasis, 2011, 9, 859-861. | 3.8 | 13 |
| 26 | Relationship between 4G/5G polymorphism in the plasminogen activator inhibitor-1 gene and obesity in Argentinian Hispanic adults. Blood Coagulation and Fibrinolysis, 2010, 21, 196-198. | 1.0 | 1 |
| 27 | Role of 4G/5G promoter polymorphism of Plasminogen Activator Inhibitor-1 (PAI-1) gene in outcome of sepsis. Thrombosis Research, 2010, 125, 367-369. | 1.7 | 11 |
| 28 | Thrombophilia in Human Immunodeficiency Virus—Infected Patients with Osteonecrosis: Is There a Real Connection? The First Case-Control Study. Clinical and Applied Thrombosis/Hemostasis, 2009, 15, 340-347. | 1.7 | 13 |
| 29 | Plasma Plasminogen Activator Inhibitor-1 Levels and Nonalcoholic Fatty Liver in Individuals With Features of Metabolic Syndrome. Clinical and Applied Thrombosis/Hemostasis, 2008, 14, 319-324. | 1.7 | 11 |
| 30 | The range of normal values of liver enzymes in the era of metabolic syndrome: the need for a redefinition. European Journal of Gastroenterology and Hepatology, 2008, 20, 589-591. | 1.6 | 2 |
| 31 | Relationship between hepatitis C virus (HCV) and insulin resistance, endothelial perturbation, and platelet activation in HIV–HCV–coinfected patients under highly active antiretroviral treatment. European Journal of Clinical Microbiology and Infectious Diseases, 2006, 25, 98-103. | 2.9 | 32 |
| 32 | Soluble thrombomodulin levels among women with a history of recurrent pregnancy loss, with or without antiphospholipid antibodies. Blood Coagulation and Fibrinolysis, 2005, 16, 31-35. | 1.0 | 4 |
| 33 | Catastrophic antiphospholipid syndrome and Kikuchi-Fujimoto disease: the first case reported. Lupus, 2005, 14, 967-969. | 1.6 | 10 |
| 34 | Insulin resistance status is an important determinant of PAI-1 levels in HIV-infected patients, independently of the lipid profile. Journal of Thrombosis and Haemostasis, 2004, 2, 532-534. | 3.8 | 9 |
| 35 | Association between the acquired free protein S deficiency in HIV-infected patients with the lipid profile levels. Journal of Thrombosis and Haemostasis, 2004, 2, 1195-1197. | 3.8 | 3 |
| 36 | Relationship between hepatic enzymes and insulin resistance syndrome markers in HIV with lipodystrophy. Hepatology, 2004, 40, 1475-1476. | 7.3 | 2 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Antibodies to tissue factor pathway inhibitor are uncommonly detected in patients with infection-related antiphospholipid antibodies. Journal of Thrombosis and Haemostasis, 2003, 1, 2250-2251. | 3.8 | 8 |
| 38 | Endothelial markers and HIV infection in the era of highly active antiretroviral treatment. Thrombosis Research, 2003, 110, 93-98. | 1.7 | 44 |
| 39 | Viral load and disease progression as responsible for endothelial activation and/or injury in human immunodeficiency virus-1-infected patients. Blood Coagulation and Fibrinolysis, 2003, 14, 15-18. | 1.0 | 60 |
| 40 | Activated protein C resistance in patients with arterial ischemic stroke. Journal of Stroke and Cerebrovascular Diseases, 2001, 10, 128-131. | 1.6 | 7 |
| 41 | High prevalence of antiphospholipid antibodies in leprosy: evaluation of antigen reactivity. Lupus, 2000, 9, 594-600. | 1.6 | 65 |
| 42 | Acquired Hypoprothrombinemia Related to High Titres of Antiprotein-phospholipid Antibodies. Thrombosis and Haemostasis, 1999, 81, 317-318. | 3.4 | 5 |
| 43 | Different Types of Antiphospholipid Antibodies in AIDS. Thrombosis Research, 1999, 96, 19-25. | 1.7 | 68 |