

Gabriela F De Larrañaga

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3605990/publications.pdf>

Version: 2024-02-01

43
papers

774
citations

758635

12
h-index

525886

27
g-index

44
all docs

44
docs citations

44
times ranked

1217
citing authors

#	ARTICLE	IF	CITATIONS
1	Clinical and therapeutic value of the adjusted Global Antiphospholipid Syndrome Score in primary obstetric antiphospholipid syndrome. <i>Lupus</i> , 2022, 31, 354-362.	0.8	5
2	A focus on the roles of histones in health and diseases. <i>Clinical Biochemistry</i> , 2021, 94, 12-19.	0.8	11
3	Risk factors for early severe preeclampsia in obstetric antiphospholipid syndrome with conventional treatment. The impact of hydroxychloroquine. <i>Lupus</i> , 2020, 29, 1736-1742.	0.8	9
4	A high-risk laboratory profile of antiphospholipid antibodies and thrombosis is associated with a large number of extra-criteria manifestations in obstetric antiphospholipid syndrome. <i>Immunologic Research</i> , 2019, 67, 478-485.	1.3	11
5	Trombofilia hereditaria y pÃ©rdidas de embarazo. Estudio de una cohorte de Argentina. <i>Medicina ClÃ©nica</i> , 2019, 152, 249-254.	0.3	5
6	Maternal carriers of the ANXA5 M2 haplotype are exposed to a greater risk for placenta-mediated pregnancy complications. <i>Journal of Assisted Reproduction and Genetics</i> , 2018, 35, 921-928.	1.2	11
7	What is the best time to assess the antiphospholipid antibodies (aPL) profile to better predict the obstetric outcome in antiphospholipid syndrome (APS) patients?. <i>Immunologic Research</i> , 2018, 66, 577-583.	1.3	6
8	Pregnancy failure in patients with obstetric antiphospholipid syndrome with conventional treatment: the influence of a triple positive antibody profile. <i>Lupus</i> , 2017, 26, 983-988.	0.8	48
9	Decreased Levels of Circulating Protein S in Patients with Active Crohn's Disease. <i>Gastroenterology</i> , 2017, 152, S612.	0.6	0
10	Effect of gene-gene and gene-environment interactions associated with antituberculosis drug-induced hepatotoxicity. <i>Pharmacogenetics and Genomics</i> , 2017, 27, 363-371.	0.7	12
11	The -308 G/A polymorphism in the tumor necrosis factor- α gene is not associated with development and progression of rheumatoid arthritis in Argentinean patients. <i>International Journal of Rheumatic Diseases</i> , 2016, 19, 476-481.	0.9	7
12	tagSNP rs1495741 as a useful molecular marker to predict antituberculosis drug-induced hepatotoxicity. <i>Pharmacogenetics and Genomics</i> , 2016, 26, 357-361.	0.7	7
13	Q222R polymorphism in the DNase I gene is not associated with susceptibility to rheumatoid arthritis or to disease course in an Argentine patient cohort. <i>Egyptian Rheumatologist</i> , 2016, 38, 289-293.	0.5	1
14	D-Dimer as a prognostic marker of morbidity and mortality among HIV patients: a call for attention. <i>Infectious Diseases</i> , 2016, 48, 860-861.	1.4	3
15	The -2518A/G polymorphism in the monocyte chemoattractant protein 1 gene (MCP-1) is associated with an increased risk of rheumatoid arthritis in Argentine patients. <i>Clinical Rheumatology</i> , 2016, 35, 3057-3061.	1.0	8
16	Paternal factor V Leiden and recurrent pregnancy loss: a new concept behind fetal genetics?: reply. <i>Journal of Thrombosis and Haemostasis</i> , 2014, 12, 1759-1760.	1.9	2
17	Paternal factor V Leiden and recurrent pregnancy loss: a new concept behind fetal genetics?. <i>Journal of Thrombosis and Haemostasis</i> , 2014, 12, 666-669.	1.9	15
18	Neutrophil Extracellular Traps in Sepsis. <i>Shock</i> , 2014, 42, 286-294.	1.0	134

#	ARTICLE	IF	CITATIONS
19	4G/5G plasminogen activator inhibitor-1 and -308 A/G tumor necrosis factor- α promoter gene polymorphisms in Argentinean lupus patients: focus on lupus nephritis. <i>Clinical and Experimental Medicine</i> , 2014, 14, 83-89.	1.9	7
20	Sex, ethnicity, and slow acetylator profile are the major causes of hepatotoxicity induced by antituberculosis drugs. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2013, 28, 323-328.	1.4	57
21	Role of -675 4G/5G in the Plasminogen Activator Inhibitor-1 Gene and -308G/A Tumor Necrosis Factor- α Gene Polymorphisms in Obese Argentinean Patients. <i>Genetic Testing and Molecular Biomarkers</i> , 2012, 16, 372-375.	0.3	3
22	The -2518 A/G polymorphism in the monocyte chemoattractant protein 1 gene is associated with the risk of developing systemic lupus erythematosus in Argentinean patients: a multicenter study. <i>European Cytokine Network</i> , 2012, 23, 7-11.	1.1	9
23	Biomarkers in sepsis at time zero: intensive care unit scores, plasma measurements and polymorphisms in Argentina. <i>Journal of Infection in Developing Countries</i> , 2012, 6, 555-562.	0.5	17
24	The distribution of allelic and genotypic frequencies of N-Acetyltransferase-2 variants in an Argentine population. <i>Journal of Infection in Developing Countries</i> , 2012, 6, 671-674.	0.5	10
25	Anti- β 2glycoprotein I antibodies from leprosy patients do not show thrombogenic effects in an in vivo animal model. <i>Journal of Thrombosis and Haemostasis</i> , 2011, 9, 859-861.	1.9	13
26	Relationship between 4G/5G polymorphism in the plasminogen activator inhibitor-1 gene and obesity in Argentinian Hispanic adults. <i>Blood Coagulation and Fibrinolysis</i> , 2010, 21, 196-198.	0.5	1
27	Role of 4G/5G promoter polymorphism of Plasminogen Activator Inhibitor-1 (PAI-1) gene in outcome of sepsis. <i>Thrombosis Research</i> , 2010, 125, 367-369.	0.8	11
28	Thrombophilia in Human Immunodeficiency Virus-Infected Patients with Osteonecrosis: Is There a Real Connection? The First Case-Control Study. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2009, 15, 340-347.	0.7	13
29	Plasma Plasminogen Activator Inhibitor-1 Levels and Nonalcoholic Fatty Liver in Individuals With Features of Metabolic Syndrome. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2008, 14, 319-324.	0.7	11
30	The range of normal values of liver enzymes in the era of metabolic syndrome: the need for a redefinition. <i>European Journal of Gastroenterology and Hepatology</i> , 2008, 20, 589-591.	0.8	2
31	Relationship between hepatitis C virus (HCV) and insulin resistance, endothelial perturbation, and platelet activation in HIV-HCV coinfecting patients under highly active antiretroviral treatment. <i>European Journal of Clinical Microbiology and Infectious Diseases</i> , 2006, 25, 98-103.	1.3	32
32	Soluble thrombomodulin levels among women with a history of recurrent pregnancy loss, with or without antiphospholipid antibodies. <i>Blood Coagulation and Fibrinolysis</i> , 2005, 16, 31-35.	0.5	4
33	Catastrophic antiphospholipid syndrome and Kikuchi-Fujimoto disease: the first case reported. <i>Lupus</i> , 2005, 14, 967-969.	0.8	10
34	Insulin resistance status is an important determinant of PAI-1 levels in HIV-infected patients, independently of the lipid profile. <i>Journal of Thrombosis and Haemostasis</i> , 2004, 2, 532-534.	1.9	9
35	Association between the acquired free protein S deficiency in HIV-infected patients with the lipid profile levels. <i>Journal of Thrombosis and Haemostasis</i> , 2004, 2, 1195-1197.	1.9	3
36	Relationship between hepatic enzymes and insulin resistance syndrome markers in HIV with lipodystrophy. <i>Hepatology</i> , 2004, 40, 1475-1476.	3.6	2

#	ARTICLE	IF	CITATIONS
37	Antibodies to tissue factor pathway inhibitor are uncommonly detected in patients with infection-related antiphospholipid antibodies. <i>Journal of Thrombosis and Haemostasis</i> , 2003, 1, 2250-2251.	1.9	8
38	Endothelial markers and HIV infection in the era of highly active antiretroviral treatment. <i>Thrombosis Research</i> , 2003, 110, 93-98.	0.8	44
39	Viral load and disease progression as responsible for endothelial activation and/or injury in human immunodeficiency virus-1-infected patients. <i>Blood Coagulation and Fibrinolysis</i> , 2003, 14, 15-18.	0.5	60
40	Activated protein C resistance in patients with arterial ischemic stroke. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2001, 10, 128-131.	0.7	7
41	High prevalence of antiphospholipid antibodies in leprosy: evaluation of antigen reactivity. <i>Lupus</i> , 2000, 9, 594-600.	0.8	65
42	Acquired Hypoprothrombinemia Related to High Titres of Antiprotein-phospholipid Antibodies. <i>Thrombosis and Haemostasis</i> , 1999, 81, 317-318.	1.8	5
43	Different Types of Antiphospholipid Antibodies in AIDS. <i>Thrombosis Research</i> , 1999, 96, 19-25.	0.8	68