Sung Woo Kwon

List of Publications by Citations

Source: https://exaly.com/author-pdf/361537/sung-woo-kwon-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

40 444 12 20 h-index g-index citations papers 2.76 47 3.2 533 avg, IF L-index ext. citations ext. papers

#	Paper	IF	Citations
40	Coronary artery calcium scoring does not add prognostic value to standard 64-section CT angiography protocol in low-risk patients suspected of having coronary artery disease. <i>Radiology</i> , 2011 , 259, 92-9	20.5	50
39	Lipoprotein(a) and LDL particle size are related to the severity of coronary artery disease. <i>Cardiology</i> , 2007 , 108, 282-9	1.6	34
38	Significance of small dense low-density lipoprotein as a risk factor for coronary artery disease and acute coronary syndrome. <i>Yonsei Medical Journal</i> , 2006 , 47, 405-14	3	29
37	Diverse left ventricular morphology and predictors of short-term outcome in patients with stress-induced cardiomyopathy. <i>International Journal of Cardiology</i> , 2013 , 168, 331-7	3.2	27
36	Serum levels of advanced glycation end products are associated with in-stent restenosis in diabetic patients. <i>Yonsei Medical Journal</i> , 2005 , 46, 78-85	3	21
35	Qualitative assessment of neointimal tissue after drug-eluting stent implantation: comparison between follow-up optical coherence tomography and intravascular ultrasound. <i>American Heart Journal</i> , 2011 , 161, 367-72	4.9	18
34	Prognostic significance of elevated lipoprotein(a) in coronary artery revascularization patients. <i>International Journal of Cardiology</i> , 2013 , 167, 1990-4	3.2	16
33	Comprehensive assessment of microcirculation after primary percutaneous intervention in ST-segment elevation myocardial infarction: insight from thermodilution-derived index of microcirculatory resistance and coronary flow reserve. <i>Coronary Artery Disease</i> , 2016 , 27, 34-9	1.4	16
32	Prognostic impact of alkaline phosphatase measured at time of presentation in patients undergoing primary percutaneous coronary intervention for ST-segment elevation myocardial infarction. <i>PLoS ONE</i> , 2017 , 12, e0171914	3.7	15
31	Outcomes of cardiac involvement in patients with late-stage Duchenne muscular dystrophy under management in the pulmonary rehabilitation center of a tertiary referral hospital. <i>Cardiology</i> , 2012 , 121, 186-93	1.6	14
30	Numbness after Transradial Cardiac Catheterization: the Results from a Nerve Conduction Study of the Superficial Radial Nerve. <i>Korean Circulation Journal</i> , 2016 , 46, 161-8	2.2	13
29	Relation Between Neutrophil-to-Lymphocyte Ratio and Index of Microcirculatory Resistance in Patients With ST-Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. <i>American Journal of Cardiology</i> , 2016 , 118, 1323-1328	3	12
28	Large false lumen area is a predictor of failed false lumen volume reduction after stent-graft repair in type B aortic dissection. <i>Journal of Endovascular Therapy</i> , 2014 , 21, 697-706	2.5	11
27	Prognostic Impact of Combined Contrast-Induced Acute Kidney Injury and Hypoxic Liver Injury in Patients with ST Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention: Results from INTERSTELLAR Registry. <i>PLoS ONE</i> , 2016 , 11, e0159416	3.7	11
26	Impact of final kissing balloon inflation after simple stent implantation for the treatment of non-left main true coronary bifurcation lesions in patients with acute coronary syndrome. International Journal of Cardiology, 2014, 177, 907-11	3.2	8
25	Synephrine-containing dietary supplement precipitating apical ballooning syndrome in a young female. <i>Korean Journal of Internal Medicine</i> , 2013 , 28, 356-60	2.5	8
24	Prognostic Impact of Combined Dysglycemia and Hypoxic Liver Injury on Admission in Patients With ST-Segment Elevation Myocardial Infarction Who Underwent Primary Percutaneous Coronary Intervention (from the INTERSTELLARICohort). <i>American Journal of Cardiology</i> , 2017 , 119, 1179-1185	3	7

(2020-2015)

23	Elevated Lipoprotein(a) has Incremental Prognostic Value in Type 2 Diabetic Patients with Symptomatic Coronary Artery Disease. <i>Journal of Atherosclerosis and Thrombosis</i> , 2015 , 22, 527-34	4	7
22	Addition of routine blood biomarkers to TIMI risk score improves predictive performance of 1-year mortality in patients with ST-segment elevation myocardial infarction. <i>BMC Cardiovascular Disorders</i> , 2020 , 20, 486	2.3	7
21	Relation of Stature to Outcomes in Korean Patients Undergoing Primary Percutaneous Coronary Intervention for Acute ST-Elevation Myocardial Infarction (from the INTERSTELLAR Registry). <i>American Journal of Cardiology</i> , 2016 , 118, 177-82	3	7
20	Predictors of stent fracture in patients treated with closed-cell design stents: sirolimus-eluting stent and its bare-metal counterpart, the BX velocity stent. <i>Coronary Artery Disease</i> , 2011 , 22, 40-4	1.4	4
19	Prognostic impact of the combination of serum transaminase and alkaline phosphatase determined in the emergency room in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. <i>PLoS ONE</i> , 2020 , 15, e0233286	3.7	4
18	Prognostic Value of Elevated Homocysteine Levels in Korean Patients with Coronary Artery Disease: A Propensity Score Matched Analysis. <i>Korean Circulation Journal</i> , 2016 , 46, 154-60	2.2	4
17	Impact of gender on heart failure presentation in non-obstructive hypertrophic cardiomyopathy. <i>Heart and Vessels</i> , 2020 , 35, 214-222	2.1	4
16	Prognostic Implications of Newly Developed T-Wave Inversion After Primary Percutaneous Coronary Intervention in Patients With ST-Segment Elevation Myocardial Infarction. <i>American Journal of Cardiology</i> , 2017 , 119, 515-519	3	3
15	Multidetector computed tomography for the evaluation of coronary artery disease; the diagnostic accuracy in calcified coronary arteries, comparing with IVUS imaging. <i>Yonsei Medical Journal</i> , 2014 , 55, 599-605	3	3
14	Prone position coronary angiography due to intractable back pain: another merit of transradial approach compared to transfemoral approach. <i>Journal of Invasive Cardiology</i> , 2012 , 24, 605-7	0.7	3
13	Outcome of Triple Antiplatelet Therapy Including Cilostazol in Elderly Patients with ST-Elevation Myocardial Infarction who Underwent Primary Percutaneous Coronary Intervention: Results from the INTERSTELLAR Registry. <i>Drugs and Aging</i> , 2017 , 34, 467-477	4.7	2
12	Kilt Technique as an Angle Modification Method for Endovascular Repair of Abdominal Aortic Aneurysm with Severe Neck Angle. <i>Annals of Thoracic and Cardiovascular Surgery</i> , 2017 , 23, 96-103	1.8	2
11	Ivabradine-Induced Torsade de Pointes in Patients with Heart Failure Reduced Ejection Fraction. <i>International Heart Journal</i> , 2020 , 61, 1044-1048	1.8	2
10	Complete Versus Culprit-Only Revascularization for ST-Segment Elevation Myocardial Infarction and Multivessel Disease in the 2 Generation Drug-Eluting Stent Era: Data from the INTERSTELLAR Registry. <i>Korean Circulation Journal</i> , 2018 , 48, 989-999	2.2	2
9	Neutrophil-to-Lymphocyte Ratio at Emergency Room Predicts Mechanical Complications of ST-segment Elevation Myocardial Infarction. <i>Journal of Korean Medical Science</i> , 2021 , 36, e131	4.7	1
8	mHealth Interventions for Lifestyle and Risk Factor Modification in Coronary Heart Disease: Randomized Controlled Trial. <i>JMIR MHealth and UHealth</i> , 2021 , 9, e29928	5.5	1
7	Assessment of optimal renin-angiotensin-system inhibition strategy in Asian patients with STEMI after primary myocardial revascularization <i>Reviews in Cardiovascular Medicine</i> , 2022 , 23, 18	3.9	
6	The selection of Eblocker after successful reperfusion in patients with ST-elevation myocardial infarction. <i>Perfusion (United Kingdom)</i> , 2020 , 35, 338-347	1.9	

- Clinical Implication of Hypoxic Liver Injury for Predicting Hypoxic Hepatitis and In-Hospital Mortality in ST Elevation Myocardial Infarction Patients. *Yonsei Medical Journal*, **2021**, 62, 877-884
- 3
- Prognostic impact of the combination of serum transaminase and alkaline phosphatase determined in the emergency room in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention **2020**, 15, e0233286
- Prognostic impact of the combination of serum transaminase and alkaline phosphatase determined in the emergency room in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention **2020**, 15, e0233286
- Prognostic impact of the combination of serum transaminase and alkaline phosphatase determined in the emergency room in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention **2020**, 15, e0233286
- Prognostic impact of the combination of serum transaminase and alkaline phosphatase determined in the emergency room in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention **2020**, 15, e0233286