

# Alessandro Nini

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/3615941/alessandro-nini-publications-by-year.pdf>

**Version:** 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

46  
papers

818  
citations

15  
h-index

27  
g-index

105  
ext. papers

1,041  
ext. citations

3.6  
avg, IF

3.63  
L-index

| #  | Paper   | IF   | Citations |
|----|---|------|-----------|
| 46 | Preoperative clinical and radiographic predictors of major vascular surgery in patients with testicular cancer undergoing post-chemotherapy residual tumor resection (PC-RPLND). <i>World Journal of Urology</i> , <b>2021</b> , 1  | 4    | 1         |
| 45 | Can local treatment prolong the sensitivity of metastatic prostate cancer to androgen deprivation or even prevent castration resistance?. <i>World Journal of Urology</i> , <b>2021</b> , 39, 3231-3237   | 4    | 1         |
| 44 | Robotic Salvage Lymph Node Dissection in Recurrent Prostate Cancer: Lessons Learned from 68 Cases and Implications for Future Clinical Management. <i>Journal of Urology</i> , <b>2021</b> , 206, 88-96   | 2.5  | 1         |
| 43 | Evaluation of HER2 expression in urothelial carcinoma cells as a biomarker for circulating tumor cells. <i>Cytometry Part B - Clinical Cytometry</i> , <b>2020</b> , 98, 355-367  | 3.4  | 5         |
| 42 | Long-term Outcomes of Salvage Lymph Node Dissection for Nodal Recurrence of Prostate Cancer After Radical Prostatectomy: Not as Good as Previously Thought. <i>European Urology</i> , <b>2020</b> , 78, 661-669   | 10.2 | 30        |
| 41 | Assessing the Best Surgical Template at Salvage Pelvic Lymph Node Dissection for Nodal Recurrence of Prostate Cancer After Radical Prostatectomy: When Can Bilateral Dissection be Omitted? Results from a Multi-institutional Series. <i>European Urology</i> , <b>2020</b> , 78, 779-782  | 10.2 | 6         |
| 40 | Evaluation of the oncologic benefit of adjunctive surgery at time of postchemotherapy retroperitoneal lymph node dissection (PC-RPLND).. <i>Journal of Clinical Oncology</i> , <b>2020</b> , 38, 388-388  | 2.2  | 1         |
| 39 | Perioperative and oncologic outcomes of open radical nephrectomy and inferior vena cava thrombectomy with liver mobilization and Pringle maneuver for Mayo III level tumor thrombus: single institution experience. <i>Minerva Urologica E Nefrologica = the Italian Journal of Urology and Nephrology</i> , <b>2020</b>          | 4.4  | 2         |
| 38 | Validation of the two best models to predict Benign Pathohistology in patients with advanced nonseminomatous germ cell tumors (NSGCT) undergoing postchemotherapy retroperitoneal lymph node dissection (PC-RPLND).. <i>Journal of Clinical Oncology</i> , <b>2020</b> , 38, 389-389  | 2.2  |           |
| 37 | Reply by Authors. <i>Journal of Urology</i> , <b>2020</b> , 204, 302  | 2.5  |           |
| 36 | Underestimation of Positron Emission Tomography/Computerized Tomography in Assessing Tumor Burden in Prostate Cancer Nodal Recurrence: Head-to-Head Comparison of Ga-PSMA and C-Choline in a Large, Multi-Institutional Series of Extended Salvage Lymph Node Dissections. <i>Journal of Urology</i> , <b>2020</b> , 204, 296-302 | 2.5  | 17        |
| 35 | Robotic Assisted Retroperitoneal Lymph Node Dissection for Small Volume Metastatic Testicular Cancer. <i>Journal of Urology</i> , <b>2020</b> , 204, 1242-1248  | 2.5  | 6         |
| 34 | Hypertension and Cardiovascular Morbidity Following Surgery for Kidney Cancer. <i>European Urology Oncology</i> , <b>2020</b> , 3, 209-215  | 6.7  | 21        |
| 33 | The critical role of lymph node dissection in selecting high-risk nonmetastatic renal cancer candidates for adjuvant therapy after nephrectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2019</b> , 37, 293.e25-293.e30  | 2.8  | 5         |
| 32 | Predictive and prognostic effect of inflammatory lymphadenopathies in renal cell carcinoma. <i>World Journal of Urology</i> , <b>2019</b> , 37, 701-708   | 4    | 0         |
| 31 | On-clamp versus off-clamp partial nephrectomy: Propensity score-matched comparison of long-term functional outcomes. <i>International Journal of Urology</i> , <b>2019</b> , 26, 985-991  | 2.3  | 14        |
| 30 | Management, Treatment, and Molecular Background of the Growing Teratoma Syndrome. <i>Urologic Clinics of North America</i> , <b>2019</b> , 46, 419-427  | 2.9  | 9         |

|    |  |      |    |
|----|--|------|----|
| 29 | Postchemotherapy retroperitoneal lymph node dissection (PC-RPLND) in patients with testis cancer in the salvage setting.. <i>Journal of Clinical Oncology</i> , <b>2019</b> , 37, 524-524  | 2.2  | 2  |
| 28 | Analysis of three models to predict pathohistology in patients undergoing postchemotherapy RPLND for (pcRPLND) advanced nonseminomatous germ cell tumors (NSGCT).. <i>Journal of Clinical Oncology</i> , <b>2019</b> , 37, e16053-e16053   | 2.2  |    |
| 27 | The side and the location of the primary tumor does not affect the probability of lymph node invasion in patients with renal cell carcinoma. <i>World Journal of Urology</i> , <b>2019</b> , 37, 1623-1629   | 4    | 2  |
| 26 | Oncological outcome of patients treated with spot-specific salvage lymphnode dissection (sLND) for positron-emission tomography (PET)-positive prostate cancer (PCa) relapse. <i>World Journal of Urology</i> , <b>2019</b> , 37, 2081-2090  | 4    | 4  |
| 25 | Preservation of Ejaculatory Function After Postchemotherapy Retroperitoneal Lymph Node Dissection (PC-RPLND) in Patients With Testicular Cancer: Template vs. Bilateral Resection. <i>Frontiers in Surgery</i> , <b>2018</b> , 5, 80   | 2.3  | 13 |
| 24 | Perioperative and Oncologic Outcomes of Nephrectomy and Caval Thrombectomy Using Extracorporeal Circulation and Deep Hypothermic Circulatory Arrest for Renal Cell Carcinoma Invading the Supradiaphragmatic Inferior Vena Cava and/or Right Atrium. <i>European Urology</i> , <b>2018</b> , 73, 700-700 | 10.2 | 19 |
| 23 | The Effect of Anatomical Location of Lymph Node Metastases on Cancer Specific Survival in Patients with Clear Cell Renal Cell Carcinoma. <i>Frontiers in Surgery</i> , <b>2018</b> , 5, 26   | 2.3  | 7  |
| 22 | Comparison of 2-Year Oncological Outcome and Early Recurrence Patterns in Patients with Urothelial Bladder Carcinoma Treated with Open or Robot-Assisted Radical Cystectomy with an Extracorporeal Urinary Diversion. <i>Urologia Internationalis</i> , <b>2018</b> , 101, 224-231                       | 1.9  | 6  |
| 21 | Development of the First Model of Radical Prostatectomy in the Mouse: A Feasibility Study. <i>European Urology</i> , <b>2018</b> , 73, 482-484   | 10.2 | 2  |
| 20 | Surgical management and outcomes of patients with bone metastases in germ cell tumors: A case series. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2018</b> , 36, 82.e1-82.e5   | 2.8  | 7  |
| 19 | Retroperitoneal lymph node dissection in the setting of elevated markers. <i>Current Opinion in Urology</i> , <b>2018</b> , 28, 435-439  | 2.8  | 6  |
| 18 | Multiparametric Magnetic Resonance Imaging/Ultrasound Fusion Prostate Biopsy-Are 2 Biopsy Cores per Magnetic Resonance Imaging Lesion Required?. <i>Journal of Urology</i> , <b>2018</b> , 200, 1030-1034  | 2.5  | 18 |
| 17 | Pathological High-risk Renal Cell Carcinoma: Trends in Clinical Characteristics Over 25 Years. <i>Anticancer Research</i> , <b>2018</b> , 38, 4123-4130  | 2.3  | 6  |
| 16 | Clinical and pathological characteristics of a series of patients with newly diagnosed primary renal liposarcoma: natural history and effect on survival. <i>Archivos Espanoles De Urologia</i> , <b>2018</b> , 71, 555-558  | 0.4  | 1  |
| 15 | When to perform preoperative chest computed tomography for renal cancer staging. <i>BJU International</i> , <b>2017</b> , 120, 490-496   | 5.6  | 4  |
| 14 | The Role of Prostate-specific Antigen Persistence After Radical Prostatectomy for the Prediction of Clinical Progression and Cancer-specific Mortality in Node-positive Prostate Cancer Patients. <i>European Urology</i> , <b>2016</b> , 69, 1142-8   | 10.2 | 38 |
| 13 | Follow-up After Treatment for Renal Cell Carcinoma: The Evidence Beyond the Guidelines. <i>European Urology Focus</i> , <b>2016</b> , 1, 272-281   | 5.1  | 18 |
| 12 | Lymphadenopathies in patients with renal cell carcinoma: clinical and pathological predictors of pathologically confirmed lymph node invasion. <i>World Journal of Urology</i> , <b>2016</b> , 34, 1139-45   | 4    | 12 |

|    |  |      |     |
|----|--|------|-----|
| 11 | Patterns of Clinical Recurrence of Node-positive Prostate Cancer and Impact on Long-term Survival. <i>European Urology</i> , <b>2015</b> , 68, 777-84  | 10.2 | 41  |
| 10 | More extensive pelvic lymph node dissection improves survival in patients with node-positive prostate cancer. <i>European Urology</i> , <b>2015</b> , 67, 212-9  | 10.2 | 134 |
| 9  | Extended pelvic lymph node dissection in patients with prostate cancer previously treated with surgery for lower urinary tract symptoms. <i>BJU International</i> , <b>2015</b> , 116, 366-72  | 5.6  | 3   |
| 8  | Impact of adjuvant radiation therapy on urinary continence recovery after radical prostatectomy. <i>European Urology</i> , <b>2014</b> , 65, 546-51  | 10.2 | 68  |
| 7  | Naftopidil for the treatment of benign prostate hyperplasia: a systematic review. <i>Current Medical Research and Opinion</i> , <b>2014</b> , 30, 719-32   | 2.5  | 17  |
| 6  | Preoperative erectile function is the only predictor of the use of a high number of phosphodiesterase type-5 inhibitors after bilateral nerve-sparing radical prostatectomy. <i>International Journal of Impotence Research</i> , <b>2014</b> , 26, 201-4                | 2.3  | 8   |
| 5  | Clinical signature and pathogenetic factors of diabetes associated with pancreas disease (T3cDM): a prospective observational study in surgical patients. <i>Acta Diabetologica</i> , <b>2014</b> , 51, 801-11   | 3.9  | 23  |
| 4  | Assessing the most accurate formula to predict the risk of lymph node metastases from prostate cancer in contemporary patients treated with radical prostatectomy and extended pelvic lymph node dissection. <i>Radiotherapy and Oncology</i> , <b>2013</b> , 109, 211-6 | 5.3  | 15  |
| 3  | Performance characteristics of computed tomography in detecting lymph node metastases in contemporary patients with prostate cancer treated with extended pelvic lymph node dissection. <i>European Urology</i> , <b>2012</b> , 61, 1132-8                               | 10.2 | 95  |
| 2  | Optimizing postoperative sexual function after radical prostatectomy. <i>Therapeutic Advances in Urology</i> , <b>2012</b> , 4, 347-65   | 3.2  | 12  |
| 1  | Tbr2-positive intermediate (basal) neuronal progenitors safeguard cerebral cortex expansion by controlling amplification of pallial glutamatergic neurons and attraction of subpallial GABAergic interneurons. <i>Genes and Development</i> , <b>2010</b> , 24, 1816-26  | 12.6 | 86  |