Arne Janssen

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/36186/arne-janssen-publications-by-year.pdf

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

 156
 5,639
 43
 68

 papers
 citations
 h-index
 g-index

 166
 6,284
 3.6
 5.56

 ext. papers
 ext. citations
 avg, IF
 L-index

#	Paper	IF	Citations
156	Artificial selection for timing of dispersal in predatory mites yields lines that differ in prey exploitation strategies <i>Ecology and Evolution</i> , 2022 , 12, e8760	2.8	
155	UV light attracts Diaphorina citri and its parasitoid. <i>Biological Control</i> , 2022 , 170, 104928	3.8	О
154	Biodiversity in and around Greenhouses: Benefits and Potential Risks for Pest Management. <i>Insects</i> , 2021 , 12,	2.8	2
153	Pesticides do not significantly reduce arthropod pest densities in the presence of natural enemies. <i>Ecology Letters</i> , 2021 , 24, 2010-2024	10	7
152	Ants affect citrus pests and their natural enemies in contrasting ways. <i>Biological Control</i> , 2021 , 158, 104	· 6 ;181	1
151	A predatory mite as potential biological control agent of Diaphorina citri. <i>BioControl</i> , 2021 , 66, 237-248	2.3	5
150	Predatory mites protect own eggs against predators. <i>Entomologia Experimentalis Et Applicata</i> , 2021 , 169, 501-507	2.1	O
149	Experimental evolution of cowpea mild mottle virus reveals recombination-driven reduction in virulence accompanied by increases in diversity and viral fitness. <i>Virus Research</i> , 2021 , 303, 198389	6.4	O
148	Extrafloral nectary-bearing leguminous trees enhance pest control and increase fruit weight in associated coffee plants. <i>Agriculture, Ecosystems and Environment</i> , 2021 , 319, 107538	5.7	4
147	Compatibility of two predator species for biological control of the two-spotted spider mite. Experimental and Applied Acarology, 2020 , 80, 409-422	2.1	2
146	High-quality alternative food reduces cannibalism in the predatory mite Amblyseius herbicolus (Acari: Phytoseiidae). <i>Experimental and Applied Acarology</i> , 2020 , 81, 189-200	2.1	8
145	Males cannibalise and females disperse in the predatory mite Phytoseiulus persimilis. <i>Experimental and Applied Acarology</i> , 2020 , 82, 185-198	2.1	1
144	The distribution of herbivores between leaves matches their performance only in the absence of competitors. <i>Ecology and Evolution</i> , 2020 , 10, 8405-8415	2.8	4
143	The omnivorous predator Macrolophus pygmaeus, a good candidate for the control of both greenhouse whitefly and poinsettia thrips on gerbera plants. <i>Insect Science</i> , 2020 , 27, 510-518	3.6	5
142	The use of volatile cues in recognition of kin eggs by predatory mites. <i>Ecological Entomology</i> , 2020 , 45, 1220-1223	2.1	2
141	Field distribution patterns of pests are asymmetrically affected by the presence of other herbivores. <i>Bulletin of Entomological Research</i> , 2020 , 110, 611-619	1.7	0
140	Plant feeding by an omnivorous predator affects plant phenology and omnivore performance. <i>Biological Control</i> , 2019 , 135, 66-72	3.8	3

(2016-2019)

139	Associative learning in immature lacewings (Ceraeochrysa cubana). <i>Entomologia Experimentalis Et Applicata</i> , 2019 , 167, 775-783	2.1	O
138	Herbivores avoid host plants previously exposed to their omnivorous predator Macrolophus pygmaeus. <i>Journal of Pest Science</i> , 2019 , 92, 737-745	5.5	14
137	Herbivore performance and plant defense after sequential attacks by inducing and suppressing herbivores. <i>Insect Science</i> , 2019 , 26, 108-118	3.6	6
136	Rock Powder Can Improve Vermicompost Chemical Properties and Plant Nutrition: an On-farm Experiment. <i>Communications in Soil Science and Plant Analysis</i> , 2018 , 49, 1-12	1.5	19
135	Gender-specific differences in cannibalism between a laboratory strain and a field strain of a predatory mite. <i>Experimental and Applied Acarology</i> , 2018 , 74, 239-247	2.1	3
134	Phytophagy of omnivorous predator Macrolophus pygmaeus affects performance of herbivores through induced plant defences. <i>Oecologia</i> , 2018 , 186, 101-113	2.9	28
133	Prey exploitation and dispersal strategies vary among natural populations of a predatory mite. <i>Ecology and Evolution</i> , 2018 , 8, 10384-10394	2.8	4
132	Ontogenetic stage-specific reciprocal intraguild predation. <i>Oecologia</i> , 2018 , 188, 743-751	2.9	1
131	Reciprocal intraguild predation and predator coexistence. <i>Ecology and Evolution</i> , 2018 , 8, 6952-6964	2.8	3
130	Parasitoids follow herbivorous insects to a novel host plant, generalist predators less so. <i>Entomologia Experimentalis Et Applicata</i> , 2017 , 162, 261-271	2.1	4
129	Performance of Orius insidiosus on alternative foods. <i>Journal of Applied Entomology</i> , 2017 , 141, 702-70)7 _{1.7}	8
128	Predator performance is impaired by the presence of a second prey species. <i>Bulletin of Entomological Research</i> , 2017 , 107, 313-321	1.7	6
127	Predatory interactions between prey affect patch selection by predators. <i>Behavioral Ecology and Sociobiology</i> , 2017 , 71, 66	2.5	8
126	How to evaluate the potential occurrence of intraguild predation. <i>Experimental and Applied Acarology</i> , 2017 , 72, 103-114	2.1	10
125	Non-crop plant to attract and conserve an aphid predator (Coleoptera: Coccinellidae) in tomato. <i>Biological Control</i> , 2017 , 115, 129-134	3.8	13
124	Two predatory mite species as potential control agents of broad mites. <i>BioControl</i> , 2017 , 62, 505-513	2.3	6
123	Supplying high-quality alternative prey in the litter increases control of an above-ground plant pest by a generalist predator. <i>Biological Control</i> , 2017 , 105, 19-26	3.8	27
122	Herbivores with similar feeding modes interact through the induction of different plant responses. <i>Oecologia</i> , 2016 , 180, 1-10	2.9	28

121	Size of predatory mites and refuge entrance determine success of biological control of the coconut mite. <i>BioControl</i> , 2016 , 61, 681-689	2.3	9
120	Antipredator behaviours of a spider mite in response to cues of dangerous and harmless predators. <i>Experimental and Applied Acarology</i> , 2016 , 69, 263-76	2.1	12
119	Down-regulation of plant defence in a resident spider mite species and its effect upon con- and heterospecifics. <i>Oecologia</i> , 2016 , 180, 161-7	2.9	32
118	Biological control of mealybugs with lacewing larvae is affected by the presence and type of supplemental prey. <i>BioControl</i> , 2016 , 61, 555-565	2.3	9
117	Breaking and entering: predators invade the shelter of their prey and gain protection. <i>Experimental and Applied Acarology</i> , 2015 , 67, 247-57	2.1	4
116	Distribution and oviposition site selection by predatory mites in the presence of intraguild predators. <i>Experimental and Applied Acarology</i> , 2015 , 67, 477-91	2.1	5
115	Alternative food promotes broad mite control on chilli pepper plants. <i>BioControl</i> , 2015 , 60, 817-825	2.3	25
114	Predators marked with chemical cues from one prey have increased attack success on another prey species. <i>Ecological Entomology</i> , 2015 , 40, 62-68	2.1	6
113	No adaptation of a herbivore to a novel host but loss of adaptation to its native host. <i>Scientific Reports</i> , 2015 , 5, 16211	4.9	6
112	Active prey mixing as an explanation for polyphagy in predatory arthropods: synergistic dietary effects on egg production despite a behavioural cost. <i>Functional Ecology</i> , 2015 , 29, 1317-1324	5.6	23
111	Mechanisms and ecological consequences of plant defence induction and suppression in herbivore communities. <i>Annals of Botany</i> , 2015 , 115, 1015-51	4.1	162
110	Extrafloral nectaries of associated trees can enhance natural pest control. <i>Agriculture, Ecosystems and Environment</i> , 2014 , 188, 198-203	5.7	47
109	Time scales of associating food and odor by predator communities in the field. <i>Behavioral Ecology</i> , 2014 , 25, 1123-1130	2.3	9
108	Increased control of thrips and aphids in greenhouses with two species of generalist predatory bugs involved in intraguild predation. <i>Biological Control</i> , 2014 , 79, 1-7	3.8	44
107	Witnessing predation can affect strength of counterattack in phytoseiids with ontogenetic predator pre	2.8	8
106	Generalist red velvet mite predator (Balaustium sp.) performs better on a mixed diet. <i>Experimental and Applied Acarology</i> , 2014 , 62, 19-32	2.1	17
105	Juvenile prey induce antipredator behaviour in adult predators. <i>Experimental and Applied Acarology</i> , 2013 , 59, 275-82	2.1	9
104	Biological control of aphids in the presence of thrips and their enemies. <i>BioControl</i> , 2013 , 58, 45-55	2.3	37

(2010-2012)

103	Predator-prey role reversals, juvenile experience and adult antipredator behaviour. <i>Scientific Reports</i> , 2012 , 2, 728	4.9	35
102	Prey temporarily escape from predation in the presence of a second prey species. <i>Ecological Entomology</i> , 2012 , 37, 529-535	2.1	22
101	Limited predator-induced dispersal in whiteflies. <i>PLoS ONE</i> , 2012 , 7, e45487	3.7	5
100	Whether ideal free or not, predatory mites distribute so as to maximize reproduction. <i>Oecologia</i> , 2012 , 169, 95-104	2.9	14
99	Intraguild predation among plant pests: western flower thrips larvae feed on whitefly crawlers. <i>BioControl</i> , 2012 , 57, 533-539	2.3	13
98	Invasion success in communities with reciprocal intraguild predation depends on the stage structure of the resident population. <i>Oikos</i> , 2012 , 121, 67-76	4	24
97	A herbivorous mite down-regulates plant defence and produces web to exclude competitors. <i>PLoS ONE</i> , 2011 , 6, e23757	3.7	44
96	A herbivore that manipulates plant defence. <i>Ecology Letters</i> , 2011 , 14, 229-36	10	171
95	Hyperpredation by generalist predatory mites disrupts biological control of aphids by the aphidophagous gall midge Aphidoletes aphidimyza. <i>Biological Control</i> , 2011 , 57, 246-252	3.8	26
94	Leaf domatia reduce intraguild predation among predatory mites. <i>Ecological Entomology</i> , 2011 , 36, 435	-441	28
93	Can plants evolve stable alliances with the enemies Lenemies?. Journal of Plant Interactions, 2011, 6, 71-	- 75 8	15
92	Context-dependent fitness effects of behavioral manipulation by a parasitoid. <i>Behavioral Ecology</i> , 2010 , 21, 33-36	2.3	8
91	Biological control of broad mites (Polyphagotarsonemus latus) with the generalist predator Amblyseius swirskii. <i>Experimental and Applied Acarology</i> , 2010 , 52, 29-34	2.1	66
90	Spider mite web mediates anti-predator behaviour. Experimental and Applied Acarology, 2010, 52, 1-10	2.1	41
89	Pollen subsidies promote whitefly control through the numerical response of predatory mites. <i>BioControl</i> , 2010 , 55, 253-260	2.3	96
88	Pest species diversity enhances control of spider mites and whiteflies by a generalist phytoseiid predator. <i>BioControl</i> , 2010 , 55, 387-398	2.3	68
87	Order of invasion affects the spatial distribution of a reciprocal intraguild predator. <i>Oecologia</i> , 2010 , 163, 79-89	2.9	21
86	Cues of intraguild predators affect the distribution of intraguild prey. <i>Oecologia</i> , 2010 , 163, 335-40	2.9	23

85	Vector and virus induce plant responses that benefit a non-vector herbivore. <i>Basic and Applied Ecology</i> , 2010 , 11, 162-169	3.2	32
84	Leaf domatia do not affect population dynamics of the predatory mite Iphiseiodes zuluagai. <i>Basic and Applied Ecology</i> , 2010 , 11, 144-152	3.2	8
83	Biological control of an acarine pest by single and multiple natural enemies. <i>Biological Control</i> , 2009 , 50, 60-65	3.8	42
82	Patterns of exclusion in an intraguild predator-prey system depend on initial conditions. <i>Journal of Animal Ecology</i> , 2008 , 77, 624-30	4.7	32
81	Domatia reduce larval cannibalism in predatory mites. <i>Ecological Entomology</i> , 2008 , 33, 374-379	2.1	36
80	Biological control of thrips and whiteflies by a shared predator: Two pests are better than one. <i>Biological Control</i> , 2008 , 44, 372-379	3.8	162
79	Herbivore benefits from vectoring plant virus through reduction of period of vulnerability to predation. <i>Oecologia</i> , 2008 , 156, 797-806	2.9	50
78	Parasitoid increases survival of its pupae by inducing hosts to fight predators. <i>PLoS ONE</i> , 2008 , 3, e2276	5 3.7	75
77	Odour-mediated sexual attraction in nabids (Heteroptera: Nabidae). <i>European Journal of Entomology</i> , 2008 , 105, 159-162		6
76	Food Web Interactions and Ecosystem Processes. <i>Ecological Studies</i> , 2008 , 175-191	1.1	
75	Adaptation in a spider mite population after long-term evolution on a single host plant. <i>Journal of Evolutionary Biology</i> , 2007 , 20, 2016-27	2.3	58
74	Use of odours by Cycloneda sanguinea to assess patch quality. <i>Entomologia Experimentalis Et Applicata</i> , 2007 , 124, 313-318	2.1	23
73	Predators induce egg retention in prey. <i>Oecologia</i> , 2007 , 150, 699-705	2.9	42
72	A phytoseiid predator from the tropics as potential biological control agent for the spider mite Tetranychus urticae Koch (Acari: Tetranychidae). <i>Biological Control</i> , 2007 , 42, 105-109	3.8	42
71			
	Habitat structure affects intraguild predation. <i>Ecology</i> , 2007 , 88, 2713-9	4.6	229
70	Habitat structure affects intraguild predation. <i>Ecology</i> , 2007 , 88, 2713-9 To be an intra-guild predator or a cannibal: is prey quality decisive?. <i>Ecological Entomology</i> , 2006 , 31, 430-436	2.1	26
	To be an intra-guild predator or a cannibal: is prey quality decisive?. <i>Ecological Entomology</i> , 2006 ,		

(2003-2006)

67	Can plants betray the presence of multiple herbivore species to predators and parasitoids? The role of learning in phytochemical information networks. <i>Ecological Research</i> , 2006 , 21, 3-8	1.9	62
66	Pheromone-induced priming of a defensive response in Western flower thrips. <i>Journal of Chemical Ecology</i> , 2006 , 32, 1599-603	2.7	27
65	Intraguild interactions between the predatory mites Neoseiulus californicus and Phytoseiulus persimilis. <i>Experimental and Applied Acarology</i> , 2006 , 38, 33-46	2.1	27
64	Previous and present diets of mite predators affect antipredator behaviour of whitefly prey. <i>Experimental and Applied Acarology</i> , 2006 , 38, 113-24	2.1	12
63	Searching behaviour of an omnivorous predator for novel and native host plants of its herbivores: a study on arthropod colonization of eucalyptus in Brazil. <i>Entomologia Experimentalis Et Applicata</i> , 2005 , 116, 135-142	2.1	25
62	Host-plant species modifies the diet of an omnivore feeding on three trophic levels. <i>Oikos</i> , 2005 , 111, 47-56	4	23
61	Adaptation in the asexual false spider mite Brevipalpus phoenicis: evidence for frozen niche variation. <i>Experimental and Applied Acarology</i> , 2005 , 36, 165-76	2.1	24
60	How predatory mites find plants with whitefly prey. Experimental and Applied Acarology, 2005, 36, 263-	7 5 .1	11
59	Can plants use an entomopathogenic virus as a defense against herbivores?. <i>Oecologia</i> , 2005 , 143, 396-	4 <u>0</u> .1 ₉	16
58	Herbivore-induced plant volatiles trigger sporulation in entomopathogenic fungi: the case of Neozygites tanajoae infecting the cassava green mite. <i>Journal of Chemical Ecology</i> , 2005 , 31, 1003-21	2.7	34
57	Fitness consequences of food-for-protection strategies in plants 2005 , 109-134		10
56	Diet of intraguild predators affects antipredator behavior in intraguild prey. <i>Behavioral Ecology</i> , 2005 , 16, 364-370	2.3	55
55	Global Persistence Despite Local Extinction in Acarine Predator-Prey Systems: Lessons From Experimental and Mathematical Exercises. <i>Advances in Ecological Research</i> , 2005 , 183-220	4.6	16
54	Prey attack and predators defend: counterattacking prey trigger parental care in predators. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2005 , 272, 1929-33	4.4	48
53	Interactions Between Two Neotropical Phytoseiid Predators on Cassava Plants and Consequences for Biological Control of a Shared Spider Mite Prey: a Screenhouse Evaluation. <i>Biocontrol Science and Technology</i> , 2004 , 14, 63-76	1.7	16
52	Herbivore arthropods benefit from vectoring plant viruses. <i>Ecology Letters</i> , 2004 , 8, 70-79	10	185
51	Vulnerability of Bemisia tabaci immatures to phytoseiid predators: Consequences for oviposition and influence of alternative food. <i>Entomologia Experimentalis Et Applicata</i> , 2004 , 110, 95-102	2.1	28
50	Phytoseiid predators of whiteflies feed and reproduce on non-prey food sources. <i>Experimental and Applied Acarology</i> , 2003 , 31, 15-26	2.1	95

49	Phytoseiid predator of whitefly feeds on plant tissue. Experimental and Applied Acarology, 2003, 31, 27-3	3<u>6</u>1	21
48	Herbivore host plant selection: whitefly learns to avoid host plants that harbour predators of her offspring. <i>Oecologia</i> , 2003 , 136, 484-8	2.9	81
47	Poor host plant quality causes omnivore to consume predator eggs. <i>Journal of Animal Ecology</i> , 2003 , 72, 478-483	4.7	66
46	Olfactory orientation of the truffle beetle, Leiodes cinnamomea. <i>Entomologia Experimentalis Et Applicata</i> , 2003 , 109, 147-153	2.1	20
45	Evolution of Exploitation and Defense in Tritrophic Interactions 2002 , 297-322		15
44	Oviposition patterns in a predatory mite reduce the risk of egg predation caused by prey. <i>Ecological Entomology</i> , 2002 , 27, 660-664	2.1	60
43	The benefits of clustering eggs: the role of egg predation and larval cannibalism in a predatory mite. <i>Oecologia</i> , 2002 , 131, 20-26	2.9	38
42	Flexible antipredator behaviour in herbivorous mites through vertical migration in a plant. <i>Oecologia</i> , 2002 , 132, 143-149	2.9	54
41	Prey preference and reproductive success of the generalist predator Orius laevigatus. <i>Oikos</i> , 2002 , 97, 116-124	4	47
40	Evolution of herbivore-induced plant volatiles. <i>Oikos</i> , 2002 , 97, 134-138	4	28
39	An ecological cost of plant defence: attractiveness of bitter cucumber plants to natural enemies of herbivores. <i>Ecology Letters</i> , 2002 , 5, 377-385	10	87
38			
<i>J</i> -	Interspecific infanticide deters predators. <i>Ecology Letters</i> , 2002 , 5, 490-494	10	60
37	Interspecific infanticide deters predators. <i>Ecology Letters</i> , 2002 , 5, 490-494 Phytoseiid predators suppress populations of Bemisia tabaci on cucumber plants with alternative food. <i>Experimental and Applied Acarology</i> , 2002 , 27, 57-68	10	117
	Phytoseiid predators suppress populations of Bemisia tabaci on cucumber plants with alternative		
37	Phytoseiid predators suppress populations of Bemisia tabaci on cucumber plants with alternative food. <i>Experimental and Applied Acarology</i> , 2002 , 27, 57-68 Prey preference, intraguild predation and population dynamics of an arthropod food web on	2.1	117
37	Phytoseiid predators suppress populations of Bemisia tabaci on cucumber plants with alternative food. <i>Experimental and Applied Acarology</i> , 2002 , 27, 57-68 Prey preference, intraguild predation and population dynamics of an arthropod food web on plants. <i>Experimental and Applied Acarology</i> , 2001 , 25, 785-808 Predatory mites avoid ovipositing near counterattacking prey. <i>Experimental and Applied Acarology</i> ,	2.1	117 34
37 36 35	Phytoseiid predators suppress populations of Bemisia tabaci on cucumber plants with alternative food. <i>Experimental and Applied Acarology</i> , 2002 , 27, 57-68 Prey preference, intraguild predation and population dynamics of an arthropod food web on plants. <i>Experimental and Applied Acarology</i> , 2001 , 25, 785-808 Predatory mites avoid ovipositing near counterattacking prey. <i>Experimental and Applied Acarology</i> , 2001 , 25, 613-23 Phytoseiid predators as potential biological control agents for Bemisia tabaci. <i>Experimental and</i>	2.1 2.1 2.1	1173435

31	Interactions mediated by predators in arthropod food webs. Neotropical Entomology, 2001, 30, 1-9	1.2	17	
30	INFERRING COLONIZATION PROCESSES FROM POPULATION DYNAMICS IN SPATIALLY STRUCTURED PREDATOR PREY SYSTEMS. <i>Ecology</i> , 2000 , 81, 3350-3361	4.6	3	
29	Can plants use entomopathogens as bodyguards?. <i>Ecology Letters</i> , 2000 , 3, 228-235	10	98	
28	Diet of a polyphagous arthropod predator affects refuge seeking of its thrips prey. <i>Animal Behaviour</i> , 2000 , 60, 369-375	2.8	83	
27	Inferring Colonization Processes from Population Dynamics in Spatially Structured Predator-Prey Systems. <i>Ecology</i> , 2000 , 81, 3350	4.6	15	
26	Kin recognition by the predatory mite Iphiseius degenerans: discrimination among own, conspecific, and heterospecific eggs. <i>Ecological Entomology</i> , 2000 , 25, 147-155	2.1	43	
25	Behaviour and indirect interactions in food webs of plant-inhabiting arthropods 1999 , 231-249		5	
24	Plants with spider-mite prey attract more predatory mites than clean plants under greenhouse conditions. <i>Entomologia Experimentalis Et Applicata</i> , 1999 , 90, 191-198	2.1	64	
23	Absence of odour-mediated avoidance of heterospecific competitors by the predatory mite Phytoseiulus persimilis. <i>Entomologia Experimentalis Et Applicata</i> , 1999 , 92, 73-82	2.1	22	
22	Attraction of a generalist predator towards herbivore-infested plants. <i>Entomologia Experimentalis Et Applicata</i> , 1999 , 93, 303-312	2.1	37	
21	Spider Mites Avoid Plants with Predators. Experimental and Applied Acarology, 1999, 23, 803-815	2.1	46	
20	Interactions between arthropod predators and plants: A conspiracy against herbivorous arthropods? 1999 , 207-229		27	
19	Do western flower thrips avoid plants infested with spider mites? Interactions between potential competitors 1999 , 375-380		4	
18	Review Behaviour and indirect interactions in food webs of plant-inhabiting arthropods. <i>Experimental and Applied Acarology</i> , 1998 , 22, 497-521	2.1	109	
17	Predators induce interspecific herbivore competition for food in refuge space. <i>Ecology Letters</i> , 1998 , 1, 171-177	10	67	
16	Predators Use Volatiles to Avoid Prey Patches with Conspecifics. <i>Journal of Animal Ecology</i> , 1997 , 66, 223	4.7	75	
15	Improved control capacity of the mite predator Phytoseiulus persimilis (Acari: Phytoseiidae) on tomato. <i>Experimental and Applied Acarology</i> , 1997 , 21, 507-518	2.1	39	
14	Metapopulation dynamics of a persisting predatorprey system in the laboratory: time series analysis. <i>Experimental and Applied Acarology</i> , 1997 , 21, 415-430	2.1	43	

13	Modelling Fungal (Neozygites cf. Floridana) Epizootics in Local Populations of Cassava Green Mites (Mononychellus Tanajoa). <i>Experimental and Applied Acarology</i> , 1997 , 21, 485-506	2.1	16
12	Odour-mediated responses of phytophagous mites to conspecific and heterospecific competitors. <i>Oecologia</i> , 1997 , 110, 179-185	2.9	130
11	Specificity of odour-mediated avoidance of competition in Drosophila parasitoids. <i>Behavioral Ecology and Sociobiology</i> , 1995 , 36, 229-235	2.5	43
10	Odour-Mediated Avoidance of Competition in Drosophila parasitoids: The Ghost of Competition. <i>Oikos</i> , 1995 , 73, 356	4	41
9	Specificity of odour-mediated avoidance of competition in Drosophila parasitoids 1995 , 36, 229		5
8	Evolution of Life-History Patterns in the Phytoseiidae 1994 , 70-98		24
7	Clutch Size in a Larval-Pupal Endoparasitoid: Consequences for Fitness. <i>Journal of Animal Ecology</i> , 1994 , 63, 807	4.7	52
6	Phytoseiid life-histories, local predator-prey dynamics, and strategies for control of tetranychid mites. <i>Experimental and Applied Acarology</i> , 1992 , 14, 233-250	2.1	102
5	Preselecting predatory mites for biological control: the use of an olfactometer. <i>Bulletin of Entomological Research</i> , 1990 , 80, 177-181	1.7	28
4	Optimal Host Selection by Drosophila Parasitoids in the Field. <i>Functional Ecology</i> , 1989 , 3, 469	5.6	76
3	Reproductive success of Amblyseius idaeus and A. anonymus on a diet of two-spotted spider mites. <i>Experimental and Applied Acarology</i> , 1988 , 4, 41-51	2.1	27
2	Ecology meets plant physiology: herbivore-induced plant responses and their indirect effects on arthropod communities188-218		36
1	The omnivorous predator Macrolophus pygmaeus induces production of plant volatiles that attract	5.5	О