

# John Browse

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3618642/publications.pdf>

Version: 2024-02-01

106  
papers

14,785  
citations

23567

58  
h-index

27406

106  
g-index

125  
all docs

125  
docs citations

125  
times ranked

11975  
citing authors

#	ARTICLE	IF	CITATIONS
1	Overexpression mutants reveal a role for a chloroplast MPD protein in regulation of reactive oxygen species during chilling in Arabidopsis. <i>Journal of Experimental Botany</i> , 2022, 73, 2666-2681.	4.8	3
2	A multigene approach secures hydroxy fatty acid production in Arabidopsis. <i>Journal of Experimental Botany</i> , 2022, 73, 2875-2888.	4.8	9
3	Expression of Physaria longchain acyl-CoA synthetases and hydroxy fatty acid accumulation in transgenic Arabidopsis. <i>Journal of Plant Physiology</i> , 2022, 274, 153717.	3.5	0
4	Molecular Approaches Reduce Saturates and Eliminate trans Fats in Food Oils. <i>Frontiers in Plant Science</i> , 2022, 13, .	3.6	4
5	Lipid Isolation from Plants. <i>Methods in Molecular Biology</i> , 2021, 2295, 3-13.	0.9	1
6	Phosphatidylglycerol Composition Is Central to Chilling Damage in the Arabidopsis <i>fab1</i> Mutant. <i>Plant Physiology</i> , 2020, 184, 1717-1730.	4.8	7
7	Castor LPCAT and PDAT1A Act in Concert to Promote Transacylation of Hydroxy-Fatty Acid onto Triacylglycerol. <i>Plant Physiology</i> , 2020, 184, 709-719.	4.8	11
8	The biochemistry of headgroup exchange during triacylglycerol synthesis in canola. <i>Plant Journal</i> , 2020, 103, 83-94.	5.7	18
9	Arabidopsis Flowers Unlocked the Mechanism of Jasmonate Signaling. <i>Plants</i> , 2019, 8, 285.	3.5	26
10	Identification, characterization and field testing of Brassica napus mutants producing high oleic oils. <i>Plant Journal</i> , 2019, 98, 33-41.	5.7	30
11	Tri-Hydroxy-Triacylglycerol Is Efficiently Produced by Position-Specific Castor Acyltransferases. <i>Plant Physiology</i> , 2019, 179, 1050-1063.	4.8	39
12	Development Defects of Hydroxy-Fatty Acid-Accumulating Seeds Are Reduced by Castor Acyltransferases. <i>Plant Physiology</i> , 2018, 177, 553-564.	4.8	17
13	Overexpression of Seipin1 Increases Oil in Hydroxy Fatty Acid-Accumulating Seeds. <i>Plant and Cell Physiology</i> , 2018, 59, 205-214.	3.1	18
14	Trimethylguanosine Synthase1 (TGS1) Is Essential for Chilling Tolerance. <i>Plant Physiology</i> , 2017, 174, 1713-1727.	4.8	25
15	Expression of Castor LPAT2 Enhances Ricinoleic Acid Content at the sn-2 Position of Triacylglycerols in Lesquerella Seed. <i>International Journal of Molecular Sciences</i> , 2016, 17, 507.	4.1	32
16	Control of Carbon Assimilation and Partitioning by Jasmonate: An Accounting of Growth-Defense Tradeoffs. <i>Plants</i> , 2016, 5, 7.	3.5	96
17	50 Years of Arabidopsis research: highlights and future directions. <i>New Phytologist</i> , 2016, 209, 921-944.	7.3	186
18	<i>WRINKLED1</i> Rescues Feedback Inhibition of Fatty Acid Synthesis in Hydroxylase-Expressing Seeds. <i>Plant Physiology</i> , 2016, 171, 179-191.	4.8	60

#	ARTICLE	IF	CITATIONS
19	Epidermal jasmonate perception is sufficient for all aspects of jasmonate-mediated male fertility in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2016, 85, 634-647.	5.7	44
20	Directed evolution increases desaturation of a cyanobacterial fatty acid desaturase in eukaryotic expression systems. <i>Biotechnology and Bioengineering</i> , 2016, 113, 1522-1530.	3.3	10
21	Identification of <i>Arabidopsis</i> <i>GPAT9</i> ( <i>At5g60620</i> ) as an Essential Gene Involved in Triacylglycerol Biosynthesis. <i>Plant Physiology</i> , 2016, 170, 163-179.	4.8	150
22	A <i>Caenorhabditis elegans</i> model for ether lipid biosynthesis and function. <i>Journal of Lipid Research</i> , 2016, 57, 265-275.	4.2	49
23	Type 1 diacylglycerol acyltransferases of <i>Brassica napus</i> preferentially incorporate oleic acid into triacylglycerol. <i>Journal of Experimental Botany</i> , 2015, 66, 6497-6506.	4.8	33
24	Male sterility in <i>Arabidopsis</i> induced by overexpression of a <i>MYC5</i> - <i>SRDX</i> chimeric repressor. <i>Plant Journal</i> , 2015, 81, 849-860.	5.7	84
25	A Small Phospholipase A2-1 from Castor Catalyzes the Removal of Hydroxy Fatty Acids from Phosphatidylcholine in Transgenic <i>Arabidopsis</i> Seeds. <i>Plant Physiology</i> , 2015, 167, 1259-1270.	4.8	50
26	Mutations in the Prokaryotic Pathway Rescue the <i>fatty acid biosynthesis1</i> Mutant in the Cold. <i>Plant Physiology</i> , 2015, 169, 442-452.	4.8	22
27	Reducing Isozyme Competition Increases Target Fatty Acid Accumulation in Seed Triacylglycerols of Transgenic <i>Arabidopsis</i> . <i>Plant Physiology</i> , 2015, 168, 36-46.	4.8	51
28	Fatty acid synthesis is inhibited by inefficient utilization of unusual fatty acids for glycerolipid assembly. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 1204-1209.	7.1	118
29	Reducing saturated fatty acids in <i>Arabidopsis</i> seeds by expression of a <i>Caenorhabditis elegans</i> 16:0-specific desaturase. <i>Plant Biotechnology Journal</i> , 2013, 11, 480-489.	8.3	12
30	Rapid separation of developing <i>Arabidopsis</i> seeds from siliques for RNA or metabolite analysis. <i>Plant Methods</i> , 2013, 9, 9.	4.3	15
31	Characterizing Jasmonate Regulation of Male Fertility in <i>Arabidopsis</i> . <i>Methods in Molecular Biology</i> , 2013, 1011, 13-23.	0.9	9
32	Cytochrome b5 Reductase Encoded by <i>CBR1</i> Is Essential for a Functional Male Gametophyte in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2013, 25, 3052-3066.	6.6	50
33	Homologous electron transport components fail to increase fatty acid hydroxylation in transgenic <i>Arabidopsis thaliana</i> . <i>F1000Research</i> , 2013, 2, 203.	1.6	7
34	Homologous electron transport components fail to increase fatty acid hydroxylation in transgenic <i>Arabidopsis thaliana</i> . <i>F1000Research</i> , 2013, 2, 203.	1.6	6
35	JAZ8 Lacks a Canonical Degron and Has an EAR Motif That Mediates Transcriptional Repression of Jasmonate Responses in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2012, 24, 536-550.	6.6	214
36	<i>Arabidopsis</i> mutants reveal that short- and long-term thermotolerance have different requirements for trienoic fatty acids. <i>Journal of Experimental Botany</i> , 2012, 63, 1435-1443.	4.8	51

#	ARTICLE	IF	CITATIONS
37	Acyl Editing and Headgroup Exchange Are the Major Mechanisms That Direct Polyunsaturated Fatty Acid Flux into Triacylglycerols. <i>Plant Physiology</i> , 2012, 160, 1530-1539.	4.8	182
38	The Arabidopsis JAZ2 Promoter Contains a G-Box and Thymidine-Rich Module that are Necessary and Sufficient for Jasmonate-Dependent Activation by MYC Transcription Factors and Repression by JAZ Proteins. <i>Plant and Cell Physiology</i> , 2012, 53, 330-343.	3.1	75
39	The Significance of Different Diacylglycerol Synthesis Pathways on Plant Oil Composition and Bioengineering. <i>Frontiers in Plant Science</i> , 2012, 3, 147.	3.6	238
40	Social Network: JAZ Protein Interactions Expand Our Knowledge of Jasmonate Signaling. <i>Frontiers in Plant Science</i> , 2012, 3, 41.	3.6	120
41	The pathway of triacylglycerol synthesis through phosphatidylcholine in Arabidopsis produces a bottleneck for the accumulation of unusual fatty acids in transgenic seeds. <i>Plant Journal</i> , 2011, 68, 387-399.	5.7	180
42	Genome-wide level and biochemical diversity of the acyl-activating enzyme superfamily in plants. <i>Plant Journal</i> , 2011, 66, 143-160.	5.7	75
43	Characterization of JAZ-interacting bHLH transcription factors that regulate jasmonate responses in Arabidopsis. <i>Journal of Experimental Botany</i> , 2011, 62, 2143-2154.	4.8	291
44	Castor Phospholipid:Diacylglycerol Acyltransferase Facilitates Efficient Metabolism of Hydroxy Fatty Acids in Transgenic Arabidopsis. <i>Plant Physiology</i> , 2011, 155, 683-693.	4.8	157
45	Construction of a Full-Length cDNA Library from Castor Endosperm for High-Throughput Functional Screening. <i>Methods in Molecular Biology</i> , 2011, 729, 37-52.	0.9	1
46	Organ fusion and defective cuticle function in a <i>lacs1 lacs2</i> double mutant of Arabidopsis. <i>Planta</i> , 2010, 231, 1089-1100.	3.2	126
47	Lipid biochemists salute the genome. <i>Plant Journal</i> , 2010, 61, 1092-1106.	5.7	67
48	Jasmonate perception by inositol-phosphate-potentiated COI1-JAZ co-receptor. <i>Nature</i> , 2010, 468, 400-405.	27.8	1,192
49	A Mutation in the <i>LPAT1</i> Gene Suppresses the Sensitivity of <i>fab1</i> Plants to Low Temperature. <i>Plant Physiology</i> , 2010, 153, 1135-1143.	4.8	13
50	Saving the Bilayer. <i>Science</i> , 2010, 330, 185-186.	12.6	12
51	MYB108 Acts Together with MYB24 to Regulate Jasmonate-Mediated Stamen Maturation in Arabidopsis. <i>Plant Physiology</i> , 2009, 149, 851-862.	4.8	222
52	An enzyme regulating triacylglycerol composition is encoded by the <i>ROD1</i> gene of Arabidopsis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 18837-18842.	7.1	275
53	The power of mutants for investigating jasmonate biosynthesis and signaling. <i>Phytochemistry</i> , 2009, 70, 1539-1546.	2.9	122
54	Top hits in contemporary JAZ: An update on jasmonate signaling. <i>Phytochemistry</i> , 2009, 70, 1547-1559.	2.9	158

#	ARTICLE	IF	CITATIONS
55	Jasmonate Passes Muster: A Receptor and Targets for the Defense Hormone. <i>Annual Review of Plant Biology</i> , 2009, 60, 183-205.	18.7	796
56	Jasmonate: Preventing the Maize Tassel from Getting in Touch with His Feminine Side. <i>Science Signaling</i> , 2009, 2, pe9.	3.6	28
57	A critical role of two positively charged amino acids in the Jas motif of Arabidopsis JAZ proteins in mediating coronatine and jasmonoyl isoleucine dependent interactions with the COI1 F-box protein. <i>Plant Journal</i> , 2008, 55, 979-988.	5.7	334
58	Metabolic engineering of hydroxy fatty acid production in plants: RcDGAT2 drives dramatic increases in ricinoleate levels in seed oil. <i>Plant Biotechnology Journal</i> , 2008, 6, 819-831.	8.3	292
59	The <i>AAE14</i> gene encodes the Arabidopsis <i>o</i> -succinylbenzoyl-CoA ligase that is essential for phyloquinone synthesis and photosystem function. <i>Plant Journal</i> , 2008, 54, 272-283.	5.7	61
60	New Weapons and a Rapid Response against Insect Attack. <i>Plant Physiology</i> , 2008, 146, 832-838.	4.8	210
61	Fatty Acid Desaturation and the Regulation of Adiposity in <i>Caenorhabditis elegans</i> . <i>Genetics</i> , 2007, 176, 865-875.	2.9	184
62	Arabidopsis ESK1 encodes a novel regulator of freezing tolerance. <i>Plant Journal</i> , 2007, 49, 786-799.	5.7	142
63	JAZ repressor proteins are targets of the SCFCO11 complex during jasmonate signalling. <i>Nature</i> , 2007, 448, 661-665.	27.8	2,055
64	An analysis of expressed sequence tags of developing castor endosperm using a full-length cDNA library. <i>BMC Plant Biology</i> , 2007, 7, 42.	3.6	51
65	A high-throughput screen for genes from castor that boost hydroxy fatty acid accumulation in seed oils of transgenic Arabidopsis. <i>Plant Journal</i> , 2006, 45, 847-856.	5.7	130
66	Transcriptional regulators of stamen development in Arabidopsis identified by transcriptional profiling. <i>Plant Journal</i> , 2006, 46, 984-1008.	5.7	299
67	A mutation in Arabidopsis cytochrome b5 reductase identified by high-throughput screening differentially affects hydroxylation and desaturation. <i>Plant Journal</i> , 2006, 48, 920-932.	5.7	70
68	Altered rates of protein transport in Arabidopsis mutants deficient in chloroplast membrane unsaturation. <i>Phytochemistry</i> , 2006, 67, 1629-1636.	2.9	19
69	A Suppressor of <i>fab1</i> Challenges Hypotheses on the Role of Thylakoid Unsaturation in Photosynthetic Function. <i>Plant Physiology</i> , 2006, 141, 1012-1020.	4.8	28
70	The role of <i>C. elegans</i> stearyl-CoA desaturases in fat storage and energy homeostasis. <i>FASEB Journal</i> , 2006, 20, A523.	0.5	0
71	Identification of a plastid acyl-acyl carrier protein synthetase in Arabidopsis and its role in the activation and elongation of exogenous fatty acids. <i>Plant Journal</i> , 2005, 44, 620-632.	5.7	60
72	Jasmonate: An Oxylin Signal with Many Roles in Plants. <i>Vitamins and Hormones</i> , 2005, 72, 431-456.	1.7	147

#	ARTICLE	IF	CITATIONS
73	The Acyl-CoA Synthetase Encoded by LACS2 Is Essential for Normal Cuticle Development in Arabidopsis. <i>Plant Cell</i> , 2004, 16, 629-642.	6.6	310
74	Identification of the Arabidopsis Palmitoyl-Monogalactosyldiacylglycerol $\Delta^7$ -Desaturase Gene FAD5, and Effects of Plastidial Retargeting of Arabidopsis Desaturases on the <i>fad5</i> Mutant Phenotype. <i>Plant Physiology</i> , 2004, 136, 4237-4245.	4.8	85
75	Counting the cost of a cold-blooded life: Metabolomics of cold acclimation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004, 101, 14996-14997.	7.1	29
76	Peroxisomal Acyl-CoA Synthetase Activity Is Essential for Seedling Development in Arabidopsis <i>thaliana</i> . <i>Plant Cell</i> , 2004, 16, 394-405.	6.6	231
77	Microarray and differential display identify genes involved in jasmonate-dependent anther development. <i>Plant Molecular Biology</i> , 2003, 52, 775-786.	3.9	65
78	Arabidopsis Contains a Large Superfamily of Acyl-Activating Enzymes. Phylogenetic and Biochemical Analysis Reveals a New Class of Acyl-Coenzyme A Synthetases. <i>Plant Physiology</i> , 2003, 132, 1065-1076.	4.8	168
79	Photoinhibition in Mutants of Arabidopsis Deficient in Thylakoid Unsaturation. <i>Plant Physiology</i> , 2002, 129, 876-885.	4.8	73
80	Mutants of Arabidopsis reveal many roles for membrane lipids. <i>Progress in Lipid Research</i> , 2002, 41, 254-278.	11.6	279
81	Polyunsaturated fatty acid synthesis: what will they think of next?. <i>Trends in Biochemical Sciences</i> , 2002, 27, 467-473.	7.5	308
82	A KAS2 cDNA complements the phenotypes of the Arabidopsis <i>fab1</i> mutant that differs in a single residue bordering the substrate binding pocket. <i>Plant Journal</i> , 2002, 29, 761-770.	5.7	65
83	Production of Polyunsaturated Fatty Acids by Polyketide Synthases in Both Prokaryotes and Eukaryotes. <i>Science</i> , 2001, 293, 290-293.	12.6	647
84	Temperature sensing and cold acclimation. <i>Current Opinion in Plant Biology</i> , 2001, 4, 241-246.	7.1	212
85	Trienoic Fatty Acids Are Required to Maintain Chloroplast Function at Low Temperatures. <i>Plant Physiology</i> , 2000, 124, 1697-1705.	4.8	209
86	Characterization of an acyl-CoA synthetase from Arabidopsis <i>thaliana</i> . <i>Biochemical Society Transactions</i> , 2000, 28, 957-958.	3.4	3
87	Antifungal compounds from idioblast cells isolated from avocado fruits. <i>Phytochemistry</i> , 2000, 54, 183-189.	2.9	70
88	Identification and Characterization of an Animal $\Delta^{12}$ Fatty Acid Desaturase Gene by Heterologous Expression in <i>Saccharomyces cerevisiae</i> . <i>Archives of Biochemistry and Biophysics</i> , 2000, 376, 399-408.	3.0	91
89	A Palmitoyl-CoA-Specific $\Delta^9$ Fatty Acid Desaturase from <i>Caenorhabditis elegans</i> . <i>Biochemical and Biophysical Research Communications</i> , 2000, 272, 263-269.	2.1	128
90	Genetic Engineering of Plant Chilling Tolerance. , 1999, 21, 79-93.		14

#	ARTICLE	IF	CITATIONS
91	Polyunsaturated membranes are required for photosynthetic competence in a mutant of <i>Arabidopsis</i> . <i>Plant Journal</i> , 1998, 15, 521-530.	5.7	71
92	A Determinant of Substrate Specificity Predicted from the Acyl-Acyl Carrier Protein Desaturase of Developing <i>Cat's Claw</i> Seed <sup>1</sup> . <i>Plant Physiology</i> , 1998, 117, 593-598.	4.8	103
93	A New Class of <i>Arabidopsis</i> Mutants with Reduced Hexadecatrienoic Acid Fatty Acid Levels <sup>1</sup> . <i>Plant Physiology</i> , 1998, 117, 923-930.	4.8	59
94	Novel mutations affecting leaf stearate content and plant size in <i>Arabidopsis</i> . <i>Theoretical and Applied Genetics</i> , 1997, 94, 975-981.	3.6	12
95	Dissecting desaturation: plants prove advantageous. <i>Trends in Cell Biology</i> , 1996, 6, 148-153.	7.9	122
96	The Critical Requirement for Linolenic Acid Is Pollen Development, Not Photosynthesis, in an <i>Arabidopsis</i> Mutant. <i>Plant Cell</i> , 1996, 8, 403.	6.6	167
97	An Octadecanoid Pathway Mutant (JL5) of Tomato Is Compromised in Signaling for Defense against Insect Attack. <i>Plant Cell</i> , 1996, 8, 2067.	6.6	81
98	Elevated Levels of High-Melting-Point Phosphatidylglycerols Do Not Induce Chilling Sensitivity in an <i>Arabidopsis</i> Mutant. <i>Plant Cell</i> , 1995, 7, 17.	6.6	12
99	Lipid Biosynthesis. <i>Plant Cell</i> , 1995, 7, 957.	6.6	407
100	Altered body morphology is caused by increased stearate levels in a mutant of <i>Arabidopsis</i> . <i>Plant Journal</i> , 1994, 6, 401-412.	5.7	60
101	Enhanced Thermal Tolerance in a Mutant of <i>Arabidopsis</i> Deficient in Palmitic Acid Unsaturation. <i>Plant Physiology</i> , 1989, 91, 401-408.	4.8	105
102	A Mutant of <i>Arabidopsis</i> Deficient in Desaturation of Palmitic Acid in Leaf Lipids. <i>Plant Physiology</i> , 1989, 90, 943-947.	4.8	131
103	Altered Chloroplast Structure and Function in a Mutant of <i>Arabidopsis</i> Deficient in Plastid Glycerol-3-Phosphate Acyltransferase Activity. <i>Plant Physiology</i> , 1989, 90, 846-853.	4.8	49
104	Enhanced Thermal Tolerance of Photosynthesis and Altered Chloroplast Ultrastructure in a Mutant of <i>Arabidopsis</i> Deficient in Lipid Desaturation. <i>Plant Physiology</i> , 1989, 90, 1134-1142.	4.8	144
105	A Mutant of <i>Arabidopsis</i> Deficient in the Chloroplast 16:1/18:1 Desaturase. <i>Plant Physiology</i> , 1989, 90, 522-529.	4.8	136
106	A Mutant of <i>Arabidopsis</i> Deficient in C <sub>18:3</sub> and C <sub>16:3</sub> Leaf Lipids. <i>Plant Physiology</i> , 1986, 81, 859-864.	4.8	163