

Akahito Sako

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3619038/publications.pdf>

Version: 2024-02-01

14
papers

292
citations

1039406

9
h-index

996533

15
g-index

15
all docs

15
docs citations

15
times ranked

613
citing authors

#	ARTICLE	IF	CITATIONS
1	Association of handgrip strength with hospitalization, cardiovascular events, and mortality in Japanese patients with type 2 diabetes. <i>Scientific Reports</i> , 2017, 7, 7041.	1.6	65
2	The impact of PNPLA3 and JAZF1 on hepatocellular carcinoma in non-viral hepatitis patients with type 2 diabetes mellitus. <i>Journal of Gastroenterology</i> , 2016, 51, 370-379.	2.3	41
3	Acute hepatitis B in Japan: Incidence, clinical practices and health policy. <i>Hepatology Research</i> , 2011, 41, 39-45.	1.8	30
4	Daily Physical Activity Assessed by a Triaxial Accelerometer Is Beneficially Associated with Waist Circumference, Serum Triglycerides, and Insulin Resistance in Japanese Patients with Prediabetes or Untreated Early Type 2 Diabetes. <i>Journal of Diabetes Research</i> , 2015, 2015, 1-6.	1.0	28
5	Hospitalization with hypoglycemia in patients without diabetes mellitus. <i>Medicine (United States)</i> , 2017, 96, e7271.	0.4	25
6	Prevalence and in-hospital mortality of gastrostomy and jejunostomy in Japan: a retrospective study with a national administrative database. <i>Gastrointestinal Endoscopy</i> , 2014, 80, 88-96.	0.5	20
7	Hospitalization for Hypoglycemia in Japanese Diabetic Patients. <i>Medicine (United States)</i> , 2015, 94, e1029.	0.4	18
8	Associations of Low-Intensity Resistance Training with Body Composition and Lipid Profile in Obese Patients with Type 2 Diabetes. <i>PLoS ONE</i> , 2015, 10, e0132959.	1.1	18
9	Associations between lower extremity muscle mass and metabolic parameters related to obesity in Japanese obese patients with type 2 diabetes. <i>PeerJ</i> , 2015, 3, e942.	0.9	15
10	Development of diabetes mellitus associated with quetiapine. <i>Medicine (United States)</i> , 2017, 96, e5900.	0.4	9
11	Hospitalization for urinary tract infections in Japan, 2010–2015: a retrospective study using a national inpatient database. <i>BMC Infectious Diseases</i> , 2021, 21, 1048.	1.3	8
12	Validation of the 3-Day Rule for Stool Bacterial Tests in Japan. <i>Internal Medicine</i> , 2014, 53, 533-539.	0.3	5
13	Short sleep duration is associated with B-type natriuretic peptide levels and predicts the death of Japanese patients with type 2 diabetes. <i>Sleep Medicine</i> , 2017, 36, 1-5.	0.8	4
14	A significant association between physical restraint and the development of venous thromboembolism in psychiatric patients. <i>International Journal of Cardiology</i> , 2012, 157, 442-443.	0.8	3