

# Jaakko LÃ¥ngsjÃ¶

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3619807/publications.pdf>

Version: 2024-02-01

10  
papers

386  
citations

1163117

8  
h-index

1372567

10  
g-index

10  
all docs

10  
docs citations

10  
times ranked

594  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Foundations of Human Consciousness: Imaging the Twilight Zone. <i>Journal of Neuroscience</i> , 2021, 41, 1769-1778.  | 3.6 | 30        |
| 2  | Alpha band frontal connectivity is a state-specific electroencephalographic correlate of unresponsiveness during exposure to dexmedetomidine and propofol. <i>British Journal of Anaesthesia</i> , 2020, 125, 518-528.                        | 3.4 | 17        |
| 3  | Differentiating Drug-related and State-related Effects of Dexmedetomidine and Propofol on the Electroencephalogram. <i>Anesthesiology</i> , 2018, 129, 22-36.   | 2.5 | 45        |
| 4  | Using Positron Emission Tomography in Revealing the Mystery of General Anesthesia: Study Design Challenges and Opportunities. <i>Methods in Enzymology</i> , 2018, 603, 279-303.  | 1.0 | 5         |
| 5  | S100B, NSE and MMP-9 fail to predict neurologic outcome while elevated S100B associates with milder initial clinical presentation after aneurysmal subarachnoid hemorrhage. <i>Journal of the Neurological Sciences</i> , 2018, 390, 129-134. | 0.6 | 9         |
| 6  | Time-courses of plasma IL-6 and HMGB-1 reflect initial severity of clinical presentation but do not predict poor neurologic outcome following subarachnoid hemorrhage. <i>ENeurologicalSci</i> , 2017, 6, 55-62.                              | 1.3 | 14        |
| 7  | Increased plasma UCH-L1 after aneurysmal subarachnoid hemorrhage is associated with unfavorable neurological outcome. <i>Journal of the Neurological Sciences</i> , 2016, 361, 144-149.   | 0.6 | 15        |
| 8  | Directional Connectivity between Frontal and Posterior Brain Regions Is Altered with Increasing Concentrations of Propofol. <i>PLoS ONE</i> , 2014, 9, e113616.   | 2.5 | 31        |
| 9  | High intensity exercise decreases global brain glucose uptake in humans. <i>Journal of Physiology</i> , 2005, 568, 323-332.   | 2.9 | 144       |
| 10 | Imaging of adrenal incidentalomas with PET using (11)C-metomidate and (18)F-FDG. <i>Journal of Nuclear Medicine</i> , 2004, 45, 972-9.  | 5.0 | 76        |