

Ron M C Herings

List of Publications by Citations

Source: <https://exaly.com/author-pdf/3622301/ron-m-c-herings-publications-by-citations.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

119
papers

3,497
citations

32
h-index

55
g-index

126
ext. papers

4,023
ext. citations

3.8
avg, IF

4.86
L-index

#	Paper	IF	Citations
119	Impact of osteoporosis treatment adherence on fracture rates in North America and Europe. <i>American Journal of Medicine</i> , 2009 , 122, S3-13	2.4	205
118	Combining electronic healthcare databases in Europe to allow for large-scale drug safety monitoring: the EU-ADR Project. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 1-11	2.6	198
117	Lower risk of cancer in patients on metformin in comparison with those on sulfonylurea derivatives: results from a large population-based follow-up study. <i>Diabetes Care</i> , 2012 , 35, 119-24	14.6	190
116	Increased risk of achilles tendon rupture with quinolone antibacterial use, especially in elderly patients taking oral corticosteroids. <i>Archives of Internal Medicine</i> , 2003 , 163, 1801-7		180
115	Non-steroidal anti-inflammatory drugs and risk of heart failure in four European countries: nested case-control study. <i>BMJ, The</i> , 2016 , 354, i4857	5.9	138
114	Psoriasis may not be an independent risk factor for acute ischemic heart disease hospitalizations: results of a large population-based Dutch cohort. <i>Journal of Investigative Dermatology</i> , 2010 , 130, 962-7	4.3	117
113	Tendon disorders attributed to fluoroquinolones: a study on 42 spontaneous reports in the period 1988 to 1998. <i>Arthritis and Rheumatism</i> , 2001 , 45, 235-9		108
112	Adherence to evidence-based statin guidelines reduces the risk of hospitalizations for acute myocardial infarction by 40%: a cohort study. <i>European Heart Journal</i> , 2007 , 28, 154-9	9.5	102
111	New opportunities for drug outcomes research in cancer patients: the linkage of the Eindhoven Cancer Registry and the PHARMO Record Linkage System. <i>European Journal of Cancer</i> , 2010 , 46, 395-404	7.5	93
110	Risk of upper gastrointestinal bleeding from different drug combinations. <i>Gastroenterology</i> , 2014 , 147, 784-792.e9; quiz e13-4	13.3	90
109	Half of breast cancer patients discontinue tamoxifen and any endocrine treatment before the end of the recommended treatment period of 5 years: a population-based analysis. <i>Breast Cancer Research and Treatment</i> , 2010 , 122, 843-51	4.4	86
108	Persisting gender differences and attenuating age differences in cardiovascular drug use for prevention and treatment of coronary heart disease, 1998-2010. <i>European Heart Journal</i> , 2013 , 34, 3198-205	9.5	81
107	The effect of discontinuation of antihypertensives on the risk of acute myocardial infarction and stroke. <i>Current Medical Research and Opinion</i> , 2008 , 24, 121-7	2.5	68
106	Differences in antihypertensive drug persistence associated with drug class and gender: a PHARMO study. <i>Pharmacoepidemiology and Drug Safety</i> , 2005 , 14, 795-803	2.6	66
105	Using electronic health care records for drug safety signal detection: a comparative evaluation of statistical methods. <i>Medical Care</i> , 2012 , 50, 890-7	3.1	64
104	Electronic healthcare databases for active drug safety surveillance: is there enough leverage?. <i>Pharmacoepidemiology and Drug Safety</i> , 2012 , 21, 611-21	2.6	63
103	Harmonization process for the identification of medical events in eight European healthcare databases: the experience from the EU-ADR project. <i>Journal of the American Medical Informatics Association: JAMIA</i> , 2013 , 20, 184-92	8.6	61

102	Enhanced persistence with tiotropium compared with other respiratory drugs in COPD. <i>Respiratory Medicine</i> , 2007 , 101, 1398-405	4.6	58
101	Type 2 Diabetes Mellitus Treatment Patterns Across Europe: A Population-based Multi-database Study. <i>Clinical Therapeutics</i> , 2017 , 39, 759-770	3.5	49
100	How representative of a general type 2 diabetes population are patients included in cardiovascular outcome trials with SGLT2 inhibitors? A large European observational study. <i>Diabetes, Obesity and Metabolism</i> , 2019 , 21, 968-974	6.7	46
99	Statin Use After Diagnosis of Colon Cancer and Patient Survival. <i>Gastroenterology</i> , 2017 , 153, 470-479.e433	4.3	44
98	The risk of new onset heart failure associated with dopamine agonist use in Parkinson's disease. <i>Pharmacological Research</i> , 2012 , 65, 358-64	10.2	43
97	Three-year dispensing patterns with long-acting inhaled drugs in COPD: a database analysis. <i>Respiratory Medicine</i> , 2011 , 105, 259-65	4.6	38
96	Risk of cardiac valve regurgitation with dopamine agonist use in Parkinson's disease and hyperprolactinaemia: a multi-country, nested case-control study. <i>Drug Safety</i> , 2012 , 35, 159-71	5.1	37
95	Non-steroidal anti-inflammatory drugs and melanoma risk: large Dutch population-based case-control study. <i>Journal of Investigative Dermatology</i> , 2009 , 129, 2620-7	4.3	37
94	Drug-induced acute myocardial infarction: identifying prime suspects from electronic healthcare records-based surveillance system. <i>PLoS ONE</i> , 2013 , 8, e72148	3.7	35
93	Use of Lipid-modifying Therapy and LDL-C Goal Attainment in a High-Cardiovascular-Risk Population in the Netherlands. <i>Clinical Therapeutics</i> , 2017 , 39, 819-827.e1	3.5	34
92	Useful Interplay Between Spontaneous ADR Reports and Electronic Healthcare Records in Signal Detection. <i>Drug Safety</i> , 2015 , 38, 1201-10	5.1	34
91	Substantial increase in the use of adjuvant systemic treatment for early stage breast cancer reflects changes in guidelines in the period 1990-2006 in the southeastern Netherlands. <i>European Journal of Cancer</i> , 2008 , 44, 1846-54	7.5	33
90	Pharmacy-Based Medical Record Linkage Systems	270-286	33
89	Prescribing of rosiglitazone and pioglitazone following safety signals: analysis of trends in dispensing patterns in the Netherlands from 1998 to 2008. <i>Drug Safety</i> , 2012 , 35, 471-80	5.1	32
88	Risk of fracture with thiazolidinediones: an individual patient data meta-analysis. <i>Frontiers in Endocrinology</i> , 2013 , 4, 11	5.7	32
87	EU-ADR healthcare database network vs. spontaneous reporting system database: preliminary comparison of signal detection. <i>Studies in Health Technology and Informatics</i> , 2011 , 166, 25-30	0.5	32
86	Increasing trends in the incidence and prevalence rates of type 1 diabetes among children and adolescents in the Netherlands. <i>Pediatric Diabetes</i> , 2016 , 17, 44-52	3.6	31
85	Replication of the OMOP experiment in Europe: evaluating methods for risk identification in electronic health record databases. <i>Drug Safety</i> , 2013 , 36 Suppl 1, S159-69	5.1	31

84	Developments of the incidence of osteoporosis in The Netherlands: a PHARMO study. <i>Pharmacoepidemiology and Drug Safety</i> , 2007 , 16, 166-72	2.6	30
83	Population-based analysis of non-steroidal anti-inflammatory drug use among children in four European countries in the SOS project: what size of data platforms and which study designs do we need to assess safety issues?. <i>BMC Pediatrics</i> , 2013 , 13, 192	2.6	29
82	Risk of acute myocardial infarction during use of individual NSAIDs: A nested case-control study from the SOS project. <i>PLoS ONE</i> , 2018 , 13, e0204746	3.7	29
81	Risk of ischemic stroke and the use of individual non-steroidal anti-inflammatory drugs: A multi-country European database study within the SOS Project. <i>PLoS ONE</i> , 2018 , 13, e0203362	3.7	28
80	Risk of dipeptidyl peptidase-4 (DPP-4) inhibitors on site-specific cancer: A systematic review and meta-analysis. <i>Diabetes/Metabolism Research and Reviews</i> , 2018 , 34, e3004	7.5	27
79	Statins are less effective in common daily practice among patients with hypercholesterolemia: the REALITY-PHARMO study. <i>Current Medical Research and Opinion</i> , 2004 , 20, 1025-33	2.5	27
78	First year of life medication use and hospital admission rates: premature compared with term infants. <i>Journal of Pediatrics</i> , 2013 , 163, 61-6.e1	3.6	26
77	Prevalence and incidence of acute and chronic comorbidity in patients with squamous cell carcinoma of the head and neck. <i>Head and Neck</i> , 2012 , 34, 238-44	4.2	25
76	Isotretinoin exposure during pregnancy: a population-based study in The Netherlands. <i>BMJ Open</i> , 2014 , 4, e005602	3	25
75	Effect of non-persistent use of oral glucose-lowering drugs on HbA1c goal attainment. <i>Current Medical Research and Opinion</i> , 2008 , 24, 2523-9	2.5	24
74	Record linkage for pharmacoepidemiological studies in cancer patients. <i>Pharmacoepidemiology and Drug Safety</i> , 2012 , 21, 94-103	2.6	23
73	Prescribing pattern of antipsychotic drugs during the years 1996-2010: a population-based database study in Europe with a focus on torsadogenic drugs. <i>British Journal of Clinical Pharmacology</i> , 2016 , 82, 487-97	3.8	23
72	The role of electronic healthcare record databases in paediatric drug safety surveillance: a retrospective cohort study. <i>British Journal of Clinical Pharmacology</i> , 2015 , 80, 304-14	3.8	21
71	Using real-world healthcare data for pharmacovigilance signal detection - the experience of the EU-ADR project. <i>Expert Review of Clinical Pharmacology</i> , 2015 , 8, 95-102	3.8	21
70	Psoriasis may not be an independent predictor for the use of cardiovascular and anti-diabetic drugs: a 5-year prevalence study. <i>Acta Dermato-Venereologica</i> , 2009 , 89, 476-83	2.2	21
69	Effect of persistent use of antihypertensives on blood pressure goal attainment. <i>Current Medical Research and Opinion</i> , 2008 , 24, 1025-31	2.5	21
68	Evaluating performance of electronic healthcare records and spontaneous reporting data in drug safety signal detection. <i>International Journal of Clinical Pharmacy</i> , 2015 , 37, 94-104	2.3	20
67	Signal detection of potentially drug-induced acute liver injury in children using a multi-country healthcare database network. <i>Drug Safety</i> , 2014 , 37, 99-108	5.1	20

66	Pulmonary embolism, myocardial infarction, and ischemic stroke in lung cancer patients: results from a longitudinal study. <i>Lung</i> , 2013 , 191, 501-9	2.9	19
65	Relation Between Different Measures of Glycemic Exposure and Microvascular and Macrovascular Complications in Patients with Type 2 Diabetes Mellitus: An Observational Cohort Study. <i>Diabetes Therapy</i> , 2017 , 8, 1097-1109	3.6	17
64	Exposure to phototoxic NSAIDs and quinolones is associated with an increased risk of melanoma. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 1437-44	2.8	16
63	Sex- and site-specific differences in colorectal cancer risk among people with type 2 diabetes. <i>International Journal of Colorectal Disease</i> , 2019 , 34, 269-276	3	16
62	A semantic approach for the homogeneous identification of events in eight patient databases: a contribution to the European eu-ADR project. <i>Studies in Health Technology and Informatics</i> , 2009 , 150, 190-4	0.5	16
61	A dynamic perspective on pharmaceutical competition, drug development and cost effectiveness. <i>Health Policy</i> , 2011 , 100, 18-24	3.2	15
60	Identifying Cases of Type 2 Diabetes in Heterogeneous Data Sources: Strategy from the EMIF Project. <i>PLoS ONE</i> , 2016 , 11, e0160648	3.7	15
59	Incident cancer risk after the start of aspirin use: results from a Dutch population-based cohort study of low dose aspirin users. <i>International Journal of Cancer</i> , 2014 , 135, 157-65	7.5	14
58	Quick titration of pergolide in cotreatment with domperidone is safe and effective. <i>Clinical Neuropharmacology</i> , 2001 , 24, 177-80	1.4	14
57	Effectiveness of initiating extrafine-particle versus fine-particle inhaled corticosteroids as asthma therapy in the Netherlands. <i>BMC Pulmonary Medicine</i> , 2016 , 16, 80	3.5	13
56	Use of multiple international healthcare databases for the detection of rare drug-associated outcomes: a pharmacoepidemiological programme comparing rosuvastatin with other marketed statins. <i>Pharmacoepidemiology and Drug Safety</i> , 2010 , 19, 1218-24	2.6	13
55	Design and evaluation of a semantic approach for the homogeneous identification of events in eight patient databases: a contribution to the European EU-ADR project. <i>Studies in Health Technology and Informatics</i> , 2010 , 160, 1085-9	0.5	13
54	Cholesterol level goal attainment with statins: clinical management guideline recommendations versus management in actual clinical practice. <i>Pharmacotherapy</i> , 2012 , 32, 631-41	5.8	12
53	Use of oral contraceptives in three European countries: a population-based multi-database study. <i>European Journal of Contraception and Reproductive Health Care</i> , 2016 , 21, 81-7	1.8	11
52	Can Electronic Health Records Databases Complement Spontaneous Reporting System Databases? A Historical-Reconstruction of the Association of Rofecoxib and Acute Myocardial Infarction. <i>Frontiers in Pharmacology</i> , 2018 , 9, 594	5.6	11
51	Glycemic control and long-acting insulin analog utilization in patients with type 2 diabetes. <i>Advances in Therapy</i> , 2010 , 27, 211-22	4.1	11
50	Rates of prostate surgery and acute urinary retention for benign prostatic hyperplasia in men treated with dutasteride or finasteride. <i>BMC Urology</i> , 2016 , 16, 53	2.2	11
49	Existing Data Sources for Clinical Epidemiology: The PHARMO Database Network. <i>Clinical Epidemiology</i> , 2020 , 12, 415-422	5.9	10

48	Prediction models for development of retinopathy in people with type 2 diabetes: systematic review and external validation in a Dutch primary care setting. <i>Diabetologia</i> , 2020 , 63, 1110-1119	10.3	10
47	Does use of estrogens decrease the Breslow thickness of melanoma of the skin? Oral contraceptives and hormonal replacement therapy. <i>Melanoma Research</i> , 2009 , 19, 327-32	3.3	10
46	LDL-C reductions and goal attainment among naive statin users in the Netherlands: real life results. <i>Current Medical Research and Opinion</i> , 2008 , 24, 2241-50	2.5	10
45	Incidence of cardiovascular events in breast cancer patients receiving chemotherapy in clinical practice. <i>Pharmacoepidemiology and Drug Safety</i> , 2008 , 17, 125-34	2.6	10
44	Clinical effectiveness of liraglutide vs basal insulin in a real-world setting: Evidence of improved glycaemic and weight control in obese people with type 2 diabetes. <i>Diabetes, Obesity and Metabolism</i> , 2018 , 20, 2093-2102	6.7	9
43	Change of initial oral antidiabetic therapy in type 2 diabetic patients. <i>International Journal of Clinical Pharmacy</i> , 2009 , 31, 622-6		9
42	Type 2 Diabetes, but Not Insulin (Analog) Treatment, Is Associated With More Advanced Stages of Breast Cancer: A National Linkage of Cancer and Pharmacy Registries. <i>Diabetes Care</i> , 2019 , 42, 434-442	14.6	8
41	Impact of a Forced Dose-Equivalent Levothyroxine Brand Switch on Plasma Thyrotropin: A Cohort Study. <i>Thyroid</i> , 2020 , 30, 821-828	6.2	8
40	Asthma-Related Outcomes in Patients Initiating Extrafine Ciclesonide or Fine-Particle Inhaled Corticosteroids. <i>Allergy, Asthma and Immunology Research</i> , 2017 , 9, 116-125	5.3	7
39	Major changes in chemotherapy regimens administered to breast cancer patients during 2000-2008 in the Netherlands. <i>Breast Journal</i> , 2013 , 19, 394-401	1.2	7
38	COX-2 inhibitors: complex association with lower risk of hospitalization for gastrointestinal events compared to traditional NSAIDs plus proton pump inhibitors. <i>Pharmacoepidemiology and Drug Safety</i> , 2009 , 18, 880-90	2.6	7
37	Occurrence of Comorbidities before and after Soft Tissue Sarcoma Diagnosis. <i>Sarcoma</i> , 2012 , 2012, 402109	1.0	7
36	Burden of Exacerbations in Patients with Moderate to Very Severe COPD in the Netherlands: A Real-life Study. <i>COPD: Journal of Chronic Obstructive Pulmonary Disease</i> , 2015 , 12, 132-43	2	6
35	Quick assessment of drug-related admissions over time (QUADRAT study). <i>Pharmacoepidemiology and Drug Safety</i> , 2015 , 24, 495-503	2.6	6
34	Antihypertensive drugs: a perspective on pharmaceutical price erosion and its impact on cost-effectiveness. <i>Value in Health</i> , 2012 , 15, 381-8	3.3	6
33	Venous thromboembolism prophylaxis after total knee or hip replacement: treatment pattern and outcomes. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 972-8	2.6	6
32	Determination of non-treatment with statins of high risk patients in The Netherlands. <i>Current Medical Research and Opinion</i> , 2010 , 26, 271-8	2.5	6
31	Costs of hospital events in patients with metastatic colorectal cancer. <i>Journal of Medical Economics</i> , 2011 , 14, 656-61	2.4	6

30	Comparison of incidences of cardiovascular events among new users of different statins: a retrospective observational cohort study. <i>Current Medical Research and Opinion</i> , 2009 , 25, 2621-9	2.5	6
29	Dutch trends in the use of potentially harmful medication during pregnancy. <i>British Journal of Clinical Pharmacology</i> , 2020 , 86, 2377-2392	3.8	5
28	Characterization and cholesterol management in patients with cardiovascular events and/or type 2 diabetes in the Netherlands. <i>Current Medical Research and Opinion</i> , 2017 , 33, 91-100	2.5	5
27	Rosuvastatin safety: a comprehensive, international pharmacoepidemiology programme. <i>Pharmacoepidemiology and Drug Safety</i> , 2006 , 15, 454-61	2.6	5
26	Post authorization safety study comparing quetiapine to risperidone and olanzapine. <i>Human Psychopharmacology</i> , 2016 , 31, 304-12	2.3	5
25	Observational cohort study: safety outcomes in children using proton pump inhibitors or histamine-2 receptor antagonists. <i>Current Medical Research and Opinion</i> , 2018 , 34, 577-583	2.5	5
24	Proton pump inhibitors are not associated with an increased risk of colorectal cancer. <i>GastroHep</i> , 2020 , 2, 165-170	1	4
23	Myocardial infarction, ischaemic stroke and pulmonary embolism before and after breast cancer hospitalisation. A population-based study. <i>Thrombosis and Haemostasis</i> , 2011 , 106, 149-55	7	4
22	Impact of EMA regulatory label changes on systemic diclofenac initiation, discontinuation, and switching to other pain medicines in Scotland, England, Denmark, and The Netherlands. <i>Pharmacoepidemiology and Drug Safety</i> , 2020 , 29, 296-305	2.6	4
21	Thrombotic risk minimization for Diane-35 and generics. <i>Pharmacoepidemiology and Drug Safety</i> , 2017 , 26, 1411-1417	2.6	3
20	Assessing the determinants of the potential for cost-effectiveness over time: the empirical case of COPD. <i>Value in Health</i> , 2013 , 16, 426-33	3.3	3
19	Infants born before 32 weeks of gestation or with respiratory disease are most likely to receive palivizumab in the Netherlands. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2015 , 104, 927-32	3.1	3
18	Characteristics and Absolute Survival of Metastatic Colorectal Cancer Patients Treated With Biologics: A Real-World Data Analysis From Three European Countries. <i>Frontiers in Oncology</i> , 2021 , 11, 630456	5.3	3
17	Prognostic models for predicting the risk of foot ulcer or amputation in people with type 2 diabetes: a systematic review and external validation study. <i>Diabetologia</i> , 2021 , 64, 1550-1562	10.3	3
16	Impact of EU regulatory label changes for diclofenac in people with cardiovascular disease in four countries: Interrupted time series regression analysis. <i>British Journal of Clinical Pharmacology</i> , 2021 , 87, 1129-1140	3.8	3
15	Characteristics of children and adolescents first prescribed proton pump inhibitors or histamine-2-receptor antagonists: an observational cohort study. <i>Current Medical Research and Opinion</i> , 2017 , 33, 2251-2259	2.5	2
14	Adherence and Persistence with Once-Daily vs Twice-Daily Direct Oral Anticoagulants Among Patients with Atrial Fibrillation: Real-World Analyses from the Netherlands, Italy and Germany.. <i>Drugs - Real World Outcomes</i> , 2022 , 1	2.2	2
13	Levothyroxine use and the risk of colorectal cancer: a large population-based case-control study. <i>Endocrine Connections</i> , 2021 ,	3.5	2

12	Rationale and design of a European epidemiological post-authorization safety study (PASS) program: rivaroxaban use in routine clinical practice. <i>Expert Opinion on Drug Safety</i> , 2020 , 19, 1513-1520 ^{4.1}	2
11	Use of intravenous iron and risk of anaphylaxis: A multinational observational post-authorisation safety study in Europe. <i>Pharmacoepidemiology and Drug Safety</i> , 2021 , 30, 1447-1457	2.6 2
10	Impact of cancer diagnosis and treatment on glycaemic control among individuals with colorectal cancer using glucose-lowering drugs. <i>Acta Diabetologica</i> , 2016 , 53, 727-35	3.9 2
9	Performance of prediction models for nephropathy in people with type 2 diabetes: systematic review and external validation study. <i>BMJ, The</i> , 2021 , 374, n2134	5.9 2
8	Tendon disorders attributed to fluoroquinolones: a study on 42 spontaneous reports in the period 1988 to 1998 2001 , 45, 235	2
7	Methylphenidate Treatment Initiated During Childhood Is Continued in Adulthood in Half of the Study Population. <i>Journal of Child and Adolescent Psychopharmacology</i> , 2019 , 29, 426-432	2.9 1
6	No Evidence for an Association Between Renal Function and Serious Bleeding Events in Patients Treated With Coumarins: A Population-Based Study. <i>Annals of Pharmacotherapy</i> , 2018 , 52, 221-234	2.9 1
5	Twenty-year trends in the use of anti-seizure medication among pregnant women in the Netherlands.. <i>Epilepsy and Behavior</i> , 2022 , 127, 108549	3.2 1
4	A population-based linked cohort of cancer and primary care data: A new source to study the management of cancer in primary care. <i>European Journal of Cancer Care</i> , 2021 , 31, e13529	2.4 1
3	Impact of EMA regulatory label changes on hydroxyzine initiation, discontinuation and switching to other medicines in Denmark, Scotland, England and the Netherlands: An interrupted time series regression analysis. <i>Pharmacoepidemiology and Drug Safety</i> , 2021 , 30, 482-491	2.6 1
2	Reduction in use of cyproterone/ethinylestradiol (Diane-35 and generics) after risk minimization measures in the Netherlands, UK and Italy. <i>Contraception</i> , 2020 , 102, 243-245	2.5
1	PHARMO Database Network. <i>Springer Series on Epidemiology and Public Health</i> , 2021 , 109-118	0.4