

Risa Ozaki

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3623336/publications.pdf>

Version: 2024-02-01

38
papers

1,517
citations

394390

19
h-index

330122

37
g-index

38
all docs

38
docs citations

38
times ranked

2306
citing authors

#	ARTICLE	IF	CITATIONS
1	Skin autofluorescence is associated with progression of kidney disease in type 2 diabetes: A prospective cohort study from the Hong Kong diabetes biobank. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2022, 32, 436-446.	2.6	11
2	Nonalbuminuric Diabetic Kidney Disease and Risk of All-Cause Mortality and Cardiovascular and Kidney Outcomes in Type 2 Diabetes: Findings From the Hong Kong Diabetes Biobank. <i>American Journal of Kidney Diseases</i> , 2022, 80, 196-206.e1.	1.9	12
3	Clinical Predictors and Long-term Impact of Acute Kidney Injury on Progression of Diabetic Kidney Disease in Chinese Patients With Type 2 Diabetes. <i>Diabetes</i> , 2022, 71, 520-529.	0.6	6
4	Effect of a Web-Based Management Guide on Risk Factors in Patients With Type 2 Diabetes and Diabetic Kidney Disease. <i>JAMA Network Open</i> , 2022, 5, e223862.	5.9	13
5	Development of genome-wide polygenic risk scores for lipid traits and clinical applications for dyslipidemia, subclinical atherosclerosis, and diabetes cardiovascular complications among East Asians. <i>Genome Medicine</i> , 2021, 13, 29.	8.2	18
6	Glycemic Variability and Time in Range During Self-titration of Once Daily Insulin Glargine 300 U/ml Versus Neutral Protamine Hagedorn Insulin in Insulin-naïve Chinese Type 2 Diabetes Patients. <i>Diabetes Therapy</i> , 2021, 12, 1399-1413.	2.5	5
7	Skin autofluorescence is associated with higher risk of cardiovascular events in Chinese adults with type 2 diabetes: A prospective cohort study from the Hong Kong Diabetes Biobank. <i>Journal of Diabetes and Its Complications</i> , 2021, 35, 108015.	2.3	9
8	Association of technologically assisted integrated care with clinical outcomes in type 2 diabetes in Hong Kong using the prospective JADE Program: A retrospective cohort analysis. <i>PLoS Medicine</i> , 2020, 17, e1003367.	8.4	24
9	Obesity, clinical, and genetic predictors for glycemic progression in Chinese patients with type 2 diabetes: A cohort study using the Hong Kong Diabetes Register and Hong Kong Diabetes Biobank. <i>PLoS Medicine</i> , 2020, 17, e1003209.	8.4	31
10	Progression of glucose intolerance and cardiometabolic risk factors over a decade in Chinese women with polycystic ovary syndrome: A case-control study. <i>PLoS Medicine</i> , 2019, 16, e1002953.	8.4	38
11	From Hong Kong Diabetes Register to JADE Program to RAMP-DM for Data-Driven Actions. <i>Diabetes Care</i> , 2019, 42, 2022-2031.	8.6	79
12	Sudomotor dysfunction independently predicts incident cardiovascular and renal events and all-cause death in type 2 diabetes: the Joint Asia Diabetes Evaluation register. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, 1320-1328.	0.7	6
13	Progression of diabetic kidney disease and trajectory of kidney function decline in Chinese patients with Type 2 diabetes. <i>Kidney International</i> , 2019, 95, 178-187.	5.2	105
14	Clinical considerations when adding a sodium-glucose co-transporter-2 inhibitor to insulin therapy in patients with diabetes mellitus. <i>Diabetes Care</i> , 2019, 25, 312-319.		1
15	Cross-sectional survey of biosimilar insulin utilization in Asia: The Joint Asia Diabetes Evaluation Program. <i>Journal of Diabetes Investigation</i> , 2018, 9, 1312-1322.	2.4	3
16	Association between educational level and cardiovascular disease and all-cause mortality in patients with type 2 diabetes: a prospective study in the Joint Asia Diabetes Evaluation Program. <i>Clinical Epidemiology</i> , 2018, Volume 10, 1561-1571.	3.0	15
17	The impact of maternal gestational weight gain on cardiometabolic risk factors in children. <i>Diabetologia</i> , 2018, 61, 2539-2548.	6.3	47
18	A proof-of-concept study to evaluate the efficacy and safety of BTI320 on post-prandial hyperglycaemia in Chinese subjects with pre-diabetes. <i>BMC Endocrine Disorders</i> , 2018, 18, 59.	2.2	2

#	ARTICLE	IF	CITATIONS
19	Depressive Symptoms, Co-Morbidities, and Glycemic Control in Hong Kong Chinese Elderly Patients With Type 2 Diabetes Mellitus. <i>Frontiers in Endocrinology</i> , 2018, 9, 261.	3.5	21
20	Evolution of Diabetes Care in Hong Kong: From the Hong Kong Diabetes Register to JADE-PEARL Program to RAMP and PEP Program. <i>Endocrinology and Metabolism</i> , 2018, 33, 17.	3.0	14
21	Regular mailing of personalized feedback reports improves glycemic control in diabetes: <sc>A</sc> randomized controlled trial. <i>Journal of Diabetes</i> , 2017, 9, 536-538.	1.8	11
22	In Utero Exposure to Maternal Hyperglycemia Increases Childhood Cardiometabolic Risk in Offspring. <i>Diabetes Care</i> , 2017, 40, 679-686.	8.6	242
23	Low testosterone and clinical outcomes in Chinese men with type 2 diabetes mellitus â€“ Hong Kong Diabetes Registry. <i>Diabetes Research and Clinical Practice</i> , 2017, 123, 97-105.	2.8	17
24	Interactome-transcriptome analysis discovers signatures complementary to GWAS Loci of Type 2 Diabetes. <i>Scientific Reports</i> , 2016, 6, 35228.	3.3	20
25	Retinal Information is Independently Associated with Cardiovascular Disease in Patients with Type 2 diabetes. <i>Scientific Reports</i> , 2016, 6, 19053.	3.3	17
26	Gender, diabetes education, and psychosocial factors are associated with persistent poor glycemic control in patients with type 2 diabetes in the <sc>J</sc>oint <sc>A</sc>sia <sc>D</sc>iabetes <sc>E</sc>valuation (<sc>JADE</sc>) program. <i>Journal of Diabetes</i> , 2016, 8, 109-119.	1.8	23
27	Progression to treatment failure among Chinese patients with type 2 diabetes initiated on metformin versus sulphonylurea monotherapyâ€“The Hong Kong Diabetes Registry. <i>Diabetes Research and Clinical Practice</i> , 2016, 112, 57-64.	2.8	5
28	Modifying Effect of Body Mass Index on Survival in Elderly Type 2 Diabetic Patients: Hong Kong Diabetes Registry. <i>Journal of the American Medical Directors Association</i> , 2016, 17, 276.e15-276.e22.	2.5	5
29	Measuring depression with CES-D in Chinese patients with type 2 diabetes: the validity and its comparison to PHQ-9. <i>BMC Psychiatry</i> , 2015, 15, 198.	2.6	79
30	Borderline ankleâ€“brachial index is associated with increased prevalence of micro- and macrovascular complications in type 2 diabetes: A cross-sectional analysis of 12,772 patients from the Joint Asia Diabetes Evaluation Program. <i>Diabetes and Vascular Disease Research</i> , 2015, 12, 334-341.	2.0	18
31	Effects of Providing Peer Support on Diabetes Management in People With Type 2 Diabetes. <i>Annals of Family Medicine</i> , 2015, 13, S42-S49.	1.9	56
32	The Clinical Utility of SUDOSCAN in Chronic Kidney Disease in Chinese Patients with Type 2 Diabetes. <i>PLoS ONE</i> , 2015, 10, e0134981.	2.5	25
33	Delivery of integrated diabetes care using logistics and information technology â€“ The Joint Asia Diabetes Evaluation (JADE) program. <i>Diabetes Research and Clinical Practice</i> , 2014, 106, S295-S304.	2.8	22
34	Metabolic profiles and treatment gaps in young-onset type 2 diabetes in Asia (the JADE programme): a cross-sectional study of a prospective cohort. <i>Lancet Diabetes and Endocrinology</i> , 2014, 2, 935-943.	11.4	210
35	Premature Mortality and Comorbidities in Young-onset Diabetes: A 7-Year Prospective Analysis. <i>American Journal of Medicine</i> , 2014, 127, 616-624.	1.5	110
36	Effects of Telephone-Based Peer Support in Patients With Type 2 Diabetes Mellitus Receiving Integrated Care. <i>JAMA Internal Medicine</i> , 2014, 174, 972.	5.1	121

#	ARTICLE	IF	CITATIONS
37	A New Tool to Detect Kidney Disease in Chinese Type 2 Diabetes Patients: Comparison of EZSCAN with Standard Screening Methods. <i>Diabetes Technology and Therapeutics</i> , 2011, 13, 937-943.	4.4	27
38	Overweight, family history of diabetes and attending schools of lower academic grading are independent predictors for metabolic syndrome in Hong Kong Chinese adolescents. <i>Archives of Disease in Childhood</i> , 2007, 92, 224-228.	1.9	49