Maree Brennan

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3639168/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Intraspecific variability of quantity and chemical composition of ethanolic knotwood extracts along the stems of three industrially important softwood species: Abies alba, Picea abies and Pseudotsuga menziesii. Holzforschung, 2021, 75, 168-179.	1.9	8
2	A generic information framework for decision-making in a forest-based bio-economy. Annals of Forest Science, 2021, 78, .	2.0	2
3	Post-Anthesis Water-stressed Barley Maintains Grain Specific Weight Through Altered Grain Composition and Plant Architecture. Plants, 2020, 9, 1564.	3.5	1
4	Yield and compositions of bark phenolic extractives from three commercially significant softwoods show intra- and inter-specific variation. Plant Physiology and Biochemistry, 2020, 155, 346-356.	5.8	9
5	Relationship between specific weight of spring barley and malt quality. Journal of Cereal Science, 2020, 95, 103006.	3.7	10
6	Quantitative and qualitative composition of bark polyphenols changes longitudinally with bark maturity in Abies alba Mill Annals of Forest Science, 2020, 77, 1.	2.0	27
7	The Structure of the Barley Husk Influences Its Resistance to Mechanical Stress. Frontiers in Plant Science, 2020, 11, 614334.	3.6	3
8	Increased grain density of spring barley (Hordeum vulgare L.) is associated with an increase in grain nitrogen. Journal of Cereal Science, 2019, 89, 102797.	3.7	9
9	The quality of barley husk-caryopsis adhesion is not correlated with caryopsis cuticle permeability. Journal of Plant Physiology, 2019, 243, 153054.	3.5	0
10	Development and Quality of Barley Husk Adhesion Correlates With Changes in Caryopsis Cuticle Biosynthesis and Composition. Frontiers in Plant Science, 2019, 10, 672.	3.6	5
11	Use of Raman microspectroscopy to predict malting barley husk adhesion quality. Plant Physiology and Biochemistry, 2019, 139, 587-590.	5.8	4
12	Occurrence of fucosylated and non-fucosylated xyloglucans in the cell walls of monocotyledons: An immunofluorescence study. Plant Physiology and Biochemistry, 2019, 139, 428-434.	5.8	11
13	Specific weight of barley grains is determined by traits affecting packing efficiency and by grain density. Journal of the Science of Food and Agriculture, 2019, 99, 2548-2555.	3.5	5
14	Variation in grain skinning among spring barley varieties induced by a controlled environment misting screen. Journal of Agricultural Science, 2017, 155, 317-325.	1.3	7
15	Husk to caryopsis adhesion in barley is influenced by pre- and post-anthesis temperatures through changes in a cuticular cementing layer on the caryopsis. BMC Plant Biology, 2017, 17, 169.	3.6	16
16	Hull to caryopsis adhesion and grain skinning in malting barley: Identification of key growth stages in the adhesion process. Journal of Cereal Science, 2016, 68, 8-15.	3.7	17
17	Wood quality assessment of Pinus radiata (radiata pine) saplings by dynamic mechanical analysis. Wood Science and Technology, 2015, 49, 1239-1250.	3.2	15
18	Using NIR and ATR-FTIR spectroscopy to rapidly detect compression wood in <i>Pinus radiata</i> . Canadian Journal of Forest Research, 2014, 44, 820-830.	1.7	25

#	Article	IF	CITATIONS
19	Pyrolysis gas-chromatography mass-spectrometry (Py-GC/MS) to identify compression wood in <i>Pinus radiata</i> saplings. Holzforschung, 2014, 68, 505-517.	1.9	10
20	Cellulose microfibril angles and cell-wall polymers in different wood types of Pinus radiata. Cellulose, 2012, 19, 1385-1404.	4.9	40
21	Distribution of Fucosylated Xyloglucans among the Walls of Different Cell Types in Monocotyledons Determined by Immunofluorescence Microscopy. Molecular Plant, 2011, 4, 144-156.	8.3	32