

Shin-Ya Kawashiri

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3644011/publications.pdf>

Version: 2024-02-01

91
papers

1,484
citations

361413

20
h-index

377865

34
g-index

94
all docs

94
docs citations

94
times ranked

1904
citing authors

#	ARTICLE	IF	CITATIONS
1	A prediction rule for disease outcome in patients with undifferentiated arthritis using magnetic resonance imaging of the wrists and finger joints and serologic autoantibodies. <i>Arthritis and Rheumatism</i> , 2009, 61, 772-778.	6.7	118
2	Effects of the anti-interleukin-6 receptor antibody, tocilizumab, on serum lipid levels in patients with rheumatoid arthritis. <i>Rheumatology International</i> , 2011, 31, 451-456.	3.0	100
3	A fatal case of acute exacerbation of interstitial lung disease in a patient with rheumatoid arthritis during treatment with tocilizumab. <i>Rheumatology International</i> , 2012, 32, 4023-4026.	3.0	75
4	CD4+CD25 ^{high} CD127 ^{low} Treg Cell Frequency from Peripheral Blood Correlates with Disease Activity in Patients with Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2011, 38, 2517-2521.	2.0	74
5	The power Doppler ultrasonography score from 24 synovial sites or 6 simplified synovial sites, including the metacarpophalangeal joints, reflects the clinical disease activity and level of serum biomarkers in patients with rheumatoid arthritis. <i>Rheumatology</i> , 2011, 50, 962-965.	1.9	73
6	Ultrasonographic examination of rheumatoid arthritis patients who are free of physical synovitis: power Doppler subclinical synovitis is associated with bone erosion. <i>Rheumatology</i> , 2014, 53, 562-569.	1.9	51
7	Multiple Serum Cytokine Profiling to Identify Combinational Diagnostic Biomarkers in Attacks of Familial Mediterranean Fever. <i>Medicine (United States)</i> , 2016, 95, e3449.	1.0	48
8	Proinflammatory Cytokines Synergistically Enhance the Production of Chemokine Ligand 20 (CCL20) from Rheumatoid Fibroblast-like Synovial Cells <i>in vitro</i> and Serum CCL20 Is Reduced <i>in vivo</i> by Biologic Disease-modifying Antirheumatic Drugs. <i>Journal of Rheumatology</i> , 2009, 36, 2397-2402.	2.0	46
9	Musculoskeletal ultrasonography assists the diagnostic performance of the 2010 classification criteria for rheumatoid arthritis. <i>Modern Rheumatology</i> , 2013, 23, 36-43.	1.8	45
10	Platelets and circulating CD34-positive cells as an indicator of the activity of the vicious cycle between hypertension and endothelial dysfunction in elderly Japanese men. <i>Atherosclerosis</i> , 2017, 259, 26-31.	0.8	38
11	Cutaneous vasculitis induced by TNF inhibitors: a report of three cases. <i>Modern Rheumatology</i> , 2010, 20, 86-89.	1.8	37
12	High serum matrix metalloproteinase 3 is characteristic of patients with paraneoplastic remitting seronegative symmetrical synovitis with pitting edema syndrome. <i>Modern Rheumatology</i> , 2012, 22, 584-588.	1.8	35
13	Decrement of serum cartilage oligomeric matrix protein (COMP) in rheumatoid arthritis (RA) patients achieving remission after 6 months of etanercept treatment: Comparison with CRP, IgM-RF, MMP-3 and anti-CCP Ab. <i>Joint Bone Spine</i> , 2010, 77, 418-420.	1.6	29
14	Disease activity score 28 may overestimate the remission induction of rheumatoid arthritis patients treated with tocilizumab: comparison with the remission by the clinical disease activity index. <i>Modern Rheumatology</i> , 2011, 21, 365-369.	1.8	27
15	Clinical outcomes in the first year of remitting seronegative symmetrical synovitis with pitting edema (RS3PE) syndrome. <i>Modern Rheumatology</i> , 2017, 27, 150-154.	1.8	27
16	Handgrip strength and subclinical carotid atherosclerosis in relation to platelet levels among hypertensive elderly Japanese. <i>Oncotarget</i> , 2017, 8, 69362-69369.	1.8	26
17	Ultrasound-detected bone erosion is a relapse risk factor after discontinuation of biologic disease-modifying antirheumatic drugs in patients with rheumatoid arthritis whose ultrasound power Doppler synovitis activity and clinical disease activity are well controlled. <i>Arthritis Research and Therapy</i> , 2017, 19, 108.	3.5	23
18	IgG Anti-“Citruillinated Protein Antibody Variable Domain Glycosylation Increases Before the Onset of Rheumatoid Arthritis and Stabilizes Thereafter: A Cross-sectional Study Encompassing ~1,500 Samples. <i>Arthritis and Rheumatology</i> , 2022, 74, 1147-1158.	5.6	23

#	ARTICLE	IF	CITATIONS
19	Musculoskeletal ultrasonography assists the diagnostic performance of the 2010 classification criteria for rheumatoid arthritis. <i>Modern Rheumatology</i> , 2013, 23, 36-43.	1.8	22
20	Early diagnosis and treatment for remission of clinically amyopathic dermatomyositis complicated by rapid progress interstitial lung disease: a report of two cases. <i>Modern Rheumatology</i> , 2013, 23, 190-194.	1.8	21
21	Synovial inflammation assessed by ultrasonography correlates with MRI-proven osteitis in patients with rheumatoid arthritis. <i>Rheumatology</i> , 2014, 53, 1452-1456.	1.9	21
22	Upregulation of Thrombospondin 1 Expression in Synovial Tissues and Plasma of Rheumatoid Arthritis: Role of Transforming Growth Factor- β 1 toward Fibroblast-like Synovial Cells. <i>Journal of Rheumatology</i> , 2015, 42, 943-947.	2.0	21
23	Ultrasonographic Efficacy of Biologic and Targeted Synthetic Disease-Modifying Antirheumatic Drug Therapy in Rheumatoid Arthritis From a Multicenter Rheumatoid Arthritis Ultrasound Prospective Cohort in Japan. <i>Arthritis Care and Research</i> , 2018, 70, 1719-1726.	3.4	21
24	Prediction of DAS28-ESR remission at 6 months by baseline variables in patients with rheumatoid arthritis treated with etanercept in Japanese population. <i>Modern Rheumatology</i> , 2009, 19, 488-492.	1.8	20
25	Possible mechanism underlying the association between height and vascular remodeling in elderly Japanese men. <i>Oncotarget</i> , 2018, 9, 7749-7757.	1.8	18
26	Type B Insulin Resistance Complicated with Systemic Lupus Erythematosus. <i>Internal Medicine</i> , 2010, 49, 487-490.	0.7	17
27	Reduction in serum levels of substance P in patients with rheumatoid arthritis by etanercept, a tumor necrosis factor inhibitor. <i>Modern Rheumatology</i> , 2011, 21, 244-250.	1.8	17
28	In rheumatoid arthritis patients treated with tocilizumab, the rate of clinical disease activity index (CDAI) remission at 24 weeks is superior in those with higher titers of IgM-rheumatoid factor at baseline. <i>Modern Rheumatology</i> , 2011, 21, 370-374.	1.8	16
29	Combination of MRI-detected bone marrow oedema with 2010 rheumatoid arthritis classification criteria improves the diagnostic probability of early rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 2219-2220.	0.9	16
30	Improvement of plasma endothelin-1 and nitric oxide in patients with systemic sclerosis by bosentan therapy. <i>Rheumatology International</i> , 2014, 34, 221-225.	3.0	16
31	Association between serum bone biomarker levels and therapeutic response to abatacept in patients with rheumatoid arthritis (RA): a multicenter, prospective, and observational RA ultrasound cohort study in Japan. <i>BMC Musculoskeletal Disorders</i> , 2021, 22, 506.	1.9	16
32	Risk factors of adverse events during treatment in elderly patients with rheumatoid arthritis: an observational study. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 346-352.	1.9	15
33	Magnetic Resonance Imaging Bone Edema at Enrollment Predicts Rapid Radiographic Progression in Patients with Early RA: Results from the Nagasaki University Early Arthritis Cohort. <i>Journal of Rheumatology</i> , 2016, 43, 1278-1284.	2.0	14
34	Disease activity score 28 may overestimate the remission induction of rheumatoid arthritis patients treated with tocilizumab: comparison with the remission by the clinical disease activity index. <i>Modern Rheumatology</i> , 2011, 21, 365-369.	1.8	14
35	Monitoring of therapeutic efficacy in a patient with RS3PE syndrome by serologic variables and radiographic methods. <i>Rheumatology International</i> , 2010, 30, 1677-1680.	3.0	13
36	Clopidogrel-associated acute arthritis. <i>Rheumatology International</i> , 2012, 32, 449-450.	3.0	13

#	ARTICLE	IF	CITATIONS
37	Detecting gastrointestinal manifestations in patients with systemic sclerosis using anti-gAChR antibodies. <i>Arthritis Research and Therapy</i> , 2020, 22, 32.	3.5	13
38	Association between thyroid cysts and hypertension by atherosclerosis status: a cross-sectional study. <i>Scientific Reports</i> , 2021, 11, 13922.	3.3	13
39	Switching to the anti-interleukin-6 receptor antibody tocilizumab in rheumatoid arthritis patients refractory to antitumor necrosis factor biologics. <i>Modern Rheumatology</i> , 2010, 20, 40-45.	1.8	13
40	Ultrasound disease activity of bilateral wrist and finger joints at three months reflects the clinical response at six months of patients with rheumatoid arthritis treated with biologic disease-modifying anti-rheumatic drugs. <i>Modern Rheumatology</i> , 2017, 27, 252-256.	1.8	12
41	Impact of single nucleotide polymorphism on short stature and reduced tongue pressure among community-dwelling elderly Japanese participants: a cross-sectional study. <i>Environmental Health and Preventive Medicine</i> , 2017, 22, 62.	3.4	12
42	Association of serum levels of fibrosis-related biomarkers with disease activity in patients with IgG4-related disease. <i>Arthritis Research and Therapy</i> , 2018, 20, 277.	3.5	12
43	Switching to the anti-interleukin-6 receptor antibody tocilizumab in rheumatoid arthritis patients refractory to antitumor necrosis factor biologics. <i>Modern Rheumatology</i> , 2010, 20, 40-45.	1.8	11
44	Association between tongue pressure and subclinical carotid atherosclerosis in relation to platelet levels in hypertensive elderly men: a cross-sectional study. <i>Environmental Health and Preventive Medicine</i> , 2018, 23, 31.	3.4	11
45	Hepatocyte growth factor and carotid intima-media thickness in relation to circulating CD34-positive cell levels. <i>Environmental Health and Preventive Medicine</i> , 2018, 23, 16.	3.4	11
46	In rheumatoid arthritis patients treated with tocilizumab, the rate of clinical disease activity index (CDAI) remission at 24 weeks is superior in those with higher titers of IgM-rheumatoid factor at baseline. <i>Modern Rheumatology</i> , 2011, 21, 370-374.	1.8	10
47	Differences in musculoskeletal ultrasound findings between RS3PE syndrome and elderly-onset rheumatoid arthritis. <i>Clinical Rheumatology</i> , 2020, 39, 1981-1988.	2.2	9
48	Methotrexate Alters the Expression of microRNA in Fibroblast-like Synovial Cells in Rheumatoid Arthritis. <i>International Journal of Molecular Sciences</i> , 2021, 22, 11561.	4.1	9
49	Rituximab-induced Acute Thrombocytopenia in Granulomatosis with Polyangiitis. <i>Internal Medicine</i> , 2018, 57, 2247-2250.	0.7	7
50	Combination of ultrasound power Doppler-verified synovitis and seropositivity accurately identifies patients with early-stage rheumatoid arthritis. <i>International Journal of Rheumatic Diseases</i> , 2019, 22, 842-851.	1.9	7
51	Life prognosis and renal relapse after induction therapy in Japanese patients with proliferative and pure membranous lupus nephritis. <i>Rheumatology</i> , 2021, 60, 2333-2341.	1.9	7
52	Prediction of organ involvement in systemic sclerosis by serum biomarkers and peripheral endothelial function. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 113, 102-108.	0.8	7
53	Prediction of DAS28-CRP remission in patients with rheumatoid arthritis treated with tacrolimus at 6 months by baseline variables. <i>Modern Rheumatology</i> , 2009, 19, 652-656.	1.8	6
54	Takayasu arteritis developing during treatment of ulcerative colitis with infliximab. <i>Modern Rheumatology</i> , 2013, 23, 572-576.	1.8	6

#	ARTICLE	IF	CITATIONS
55	Presence of ultrasound subclinical synovitis and increment of serum vascular endothelial growth factor in a patient with rheumatoid arthritis achieved in sustained clinical remission by treatment with adalimumab and methotrexate. <i>Modern Rheumatology</i> , 2014, 24, 681-684.	1.8	6
56	Synovitis of sternoclavicular and peripheral joints can be detected by ultrasound in patients with SAPHO syndrome. <i>Modern Rheumatology</i> , 2017, 27, 881-885.	1.8	6
57	Diagnostic efficacy of ultrasound detection of enthesitis in peripheral spondyloarthritis. <i>Modern Rheumatology</i> , 2020, 30, 1060-1066.	1.8	6
58	A patient with systemic lupus erythematosus who developed massive small intestinal hemorrhaging during treatment for chronic lupus peritonitis. <i>Modern Rheumatology</i> , 2012, 22, 312-315.	1.8	5
59	Two rare cases of adult-onset phalangeal microgeodic syndrome with magnetic resonance imaging-proven bone edema transiently occurring in winter. <i>Joint Bone Spine</i> , 2013, 80, 523-524.	1.6	5
60	Confirmation of effectiveness of tocilizumab by ultrasonography and magnetic resonance imaging in biologic agent-naïve early-stage rheumatoid arthritis patients. <i>Modern Rheumatology</i> , 2015, 25, 948-953.	1.8	5
61	Association between high psychological distress and poor oral health-related quality of life (OHQoL) in Japanese community-dwelling people: the Nagasaki Islands Study. <i>Environmental Health and Preventive Medicine</i> , 2020, 25, 82.	3.4	5
62	Non-TNF inhibitor switchers versus TNF inhibitor cyclers from multicentre rheumatoid arthritis ultrasonography prospective cohort in Japan. <i>Immunological Medicine</i> , 2020, 43, 115-120.	2.6	5
63	Successful treatment of chronic lupus myocarditis with prednisolone and mizoribine. <i>Modern Rheumatology</i> , 2010, 20, 606-610.	1.8	4
64	An elderly patient with chronic active Epstein-Barr virus infection with mixed cryoglobulinemia and review of the literature. <i>Modern Rheumatology</i> , 2013, 23, 1022-1028.	1.8	4
65	Automated Breast Volume Scanner, a new automated ultrasonic device, is useful to examine joint injuries in patients with rheumatoid arthritis. <i>Modern Rheumatology</i> , 2015, 25, 837-841.	1.8	4
66	Assessment of both articular synovitis and tenosynovitis by ultrasound is useful for evaluations of hand dysfunction in early rheumatoid arthritis patients. <i>Modern Rheumatology</i> , 2017, 27, 605-608.	1.8	4
67	Effect of abatacept treatment on serum osteoclast-related biomarkers in patients with rheumatoid arthritis (RA). <i>Medicine (United States)</i> , 2021, 100, e26592.	1.0	4
68	A case of Sjögren syndrome with pulmonary amyloidosis complicating microscopic polyangiitis. <i>Modern Rheumatology</i> , 2011, 21, 646-650.	1.8	3
69	Phalangeal Microgeodic Syndrome in a Patient with Systemic Lupus Erythematosus. <i>Internal Medicine</i> , 2017, 56, 3385-3387.	0.7	3
70	Certolizumab pegol was effective for treating residual synovitis after total knee arthroplasty in a patient with rheumatoid arthritis: therapeutic monitoring by ultrasound. <i>Journal of Medical Ultrasonics (2001)</i> , 2018, 45, 371-374.	1.3	3
71	Effectiveness and safety of non-tumor necrosis factor inhibitor therapy for anti-human T-cell leukemia virus type 1 antibody-positive rheumatoid arthritis. <i>Modern Rheumatology</i> , 2020, 31, 1-12.	1.8	3
72	Contributing factors of clinical outcomes at 1 year post-diagnosis in early rheumatoid arthritis patients with tightly controlled disease activity in clinical practice: a retrospective study. <i>Modern Rheumatology</i> , 2021, 31, 343-349.	1.8	3

#	ARTICLE	IF	CITATIONS
73	Utility of a simplified ultrasonography scoring system among patients with rheumatoid arthritis. <i>Medicine (United States)</i> , 2021, 100, e23254.	1.0	3
74	The stiff elastographic change of submandibular glands rapidly improves after the introduction of glucocorticoid treatment in patients with IgG4-related dacryoadenitis and sialoadenitis. <i>Modern Rheumatology</i> , 2016, 26, 463-464.	1.8	2
75	Switching from originator infliximab to biosimilar infliximab in Japanese patients with rheumatoid arthritis achieving clinical remission (the IFX-SIRIUS study I). <i>Medicine (United States)</i> , 2020, 99, e21151.	1.0	2
76	Insulin-Like Growth Factor-1 (IGF-1) and Reduced Tongue Pressure in Relation to Atherosclerosis Among Community-Dwelling Elderly Japanese Men: A Cross-Sectional Study. <i>Dysphagia</i> , 2020, 35, 948-954.	1.8	2
77	Usefulness of ultrasonography-proven tenosynovitis to monitor disease activity of a patient with very early rheumatoid arthritis treated by abatacept. <i>Modern Rheumatology</i> , 2013, 23, 582-586.	1.8	2
78	A patient with systemic lupus erythematosus who developed massive small intestinal hemorrhaging during treatment for chronic lupus peritonitis. <i>Modern Rheumatology</i> , 2012, 22, 312-315.	1.8	2
79	Rapid improvement of Clinical Disease Activity Index (CDAI) at 3 months predicts a preferable CDAI outcome at 1 year in active rheumatoid arthritis patients treated with tocilizumab: results from an observational investigation of daily clinical practice. <i>Clinical and Experimental Rheumatology</i> , 2016, 34, 808-812.	0.8	2
80	Comparison of complications during 1-year follow-up between remitting seronegative symmetrical synovitis with pitting edema syndrome and elderly-onset rheumatoid arthritis. <i>Immunological Medicine</i> , 2022, , 1-7.	2.6	2
81	Contribution of an adenine to guanine single nucleotide polymorphism of the matrix metalloproteinase-13 (MMP-13) $\hat{\sim}$ 77 promoter region to the production of anticyclic citrullinated peptide antibodies in patients with HLA-DRB1*shared epitope-negative rheumatoid arthritis. <i>Modern Rheumatology</i> , 2011, 21, 240-243.	1.8	1
82	Usefulness of ultrasonography-proven tenosynovitis to monitor disease activity of a patient with very early rheumatoid arthritis treated by abatacept. <i>Modern Rheumatology</i> , 2013, 23, 582-586.	1.8	1
83	Presence of ultrasound subclinical synovitis and increment of serum vascular endothelial growth factor in a patient with rheumatoid arthritis achieved in sustained clinical remission by treatment with adalimumab and methotrexate. <i>Modern Rheumatology</i> , 0, , 1.	1.8	1
84	Ultrasonography and magnetic resonance imaging findings of rheumatoid arthritis-like arthritis in a patient with adult T-cell leukemia. <i>Modern Rheumatology</i> , 2016, 26, 971-975.	1.8	1
85	Ultrasound-Proven Severe Synovitis Induced by PD-1 Inhibitor Therapy in a Patient Predisposed to Seronegative Inflammatory Arthritis. <i>Case Reports in Rheumatology</i> , 2019, 2019, 1-3.	0.6	1
86	A case of Sjögren syndrome with pulmonary amyloidosis complicating microscopic polyangiitis. <i>Modern Rheumatology</i> , 2011, 21, 646-650.	1.8	1
87	Deux observations rares de syndrome micrographique des phalanges survenant chez lâ€™adulte et comportant un Å“dÅ“me osseux transitoire hivernal. <i>Revue Du Rhumatisme (Edition Francaise)</i> , 2013, 80, 515-516.	0.0	0
88	AB0172â€¦EXPRESSION OF SLAMF6 AND ITS FUNCTIONAL SIGNIFICANCE IN PODOCYTES OF LUPUS NEPHRITIS. , 2019, , .		0
89	Discontinuation of methotrexate in rheumatoid arthritis patients achieving clinical remission by treatment with upadacitinib plus methotrexate (DOPPLER study). <i>Medicine (United States)</i> , 2022, 101, e28463.	1.0	0
90	2. Usefulness and Pitfall of Imaging Modalities in the Management of Patients with Arthritis. <i>The Journal of the Japanese Society of Internal Medicine</i> , 2021, 110, 464-470.	0.0	0

#	ARTICLE	IF	CITATIONS
91	Cross-reactivity of anti-modified protein antibodies is also present in predisease and individuals without rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2022, 81, 1332-1334.	0.9	0