

Kirsi Nuolivirta

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3646889/publications.pdf>

Version: 2024-02-01

56
papers

727
citations

516215

16
h-index

580395

25
g-index

57
all docs

57
docs citations

57
times ranked

767
citing authors

#	ARTICLE	IF	CITATIONS
1	IL-10 gene polymorphism at 1082 A/G is associated with severe rhinovirus bronchiolitis in infants. <i>Pediatric Pulmonology</i> , 2008, 43, 391-395.	1.0	79
2	<i>Bordetella pertussis</i> Infection Is Common in Nonvaccinated Infants Admitted for Bronchiolitis. <i>Pediatric Infectious Disease Journal</i> , 2010, 29, 1013-1015.	1.1	66
3	<i>Bordetella pertussis</i> infection is common in nonvaccinated infants admitted for bronchiolitis. <i>Pediatric Infectious Disease Journal</i> , 2010, 29, 1013-5.	1.1	50
4	GENE POLYMORPHISM OF IFNG +874 T/A AND TLR4 +896 A/G AND RECURRENT INFECTIONS AND WHEEZING IN TODDLERS WITH HISTORY OF BRONCHIOLITIS. <i>Pediatric Infectious Disease Journal</i> , 2009, 28, 1121-1123.	1.1	32
5	The Association of Genetic Variants in Toll-like Receptor 2 Subfamily With Allergy and Asthma After Hospitalization for Bronchiolitis in Infancy. <i>Pediatric Infectious Disease Journal</i> , 2014, 33, 463-466.	1.1	32
6	Lung function by impulse oscillometry at age 5-7 years after bronchiolitis at age 6 months. <i>Pediatric Pulmonology</i> , 2015, 50, 389-395.	1.0	28
7	IL-17A gene polymorphism rs2275913 is associated with the development of asthma after bronchiolitis in infancy. <i>Allergy International</i> , 2018, 67, 109-113.	1.4	28
8	Toll-like Receptor 3 L412F Polymorphisms in Infants With Bronchiolitis and Postbronchiolitis Wheezing. <i>Pediatric Infectious Disease Journal</i> , 2012, 31, 920-923.	1.1	25
9	Prospective study confirms that bronchiolitis in early infancy increases the risk of reduced lung function at 10-13 years of age. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2019, 108, 124-130.	0.7	22
10	Toll-like receptor 2 subfamily gene polymorphisms are associated with <i>Bacillus Calmette-Guérin</i> osteitis following newborn vaccination. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2015, 104, 485-490.	0.7	20
11	Finnish guidelines for the treatment of community-acquired pneumonia and pertussis in children. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2016, 105, 39-43.	0.7	20
12	Polymorphism in the gene encoding toll-like receptor 10 may be associated with asthma after bronchiolitis. <i>Scientific Reports</i> , 2017, 7, 2956.	1.6	20
13	Mannose-Binding Lectin Gene Polymorphisms in Infants with Bronchiolitis and Post-Bronchiolitis Wheezing. <i>Allergy International</i> , 2012, 61, 305-309.	1.4	19
14	Gene Polymorphism of Toll-Like Receptors and Lung Function at Five to Seven Years of Age after Infant Bronchiolitis. <i>PLoS ONE</i> , 2016, 11, e0146526.	1.1	18
15	IL-10 gene polymorphism is associated with preschool atopy and early-life recurrent wheezing after bronchiolitis in infancy. <i>Pediatric Pulmonology</i> , 2017, 52, 14-20.	1.0	18
16	Polymorphism of the rs1800896 IL10 promoter gene protects children from post-bronchiolitis asthma. <i>Pediatric Pulmonology</i> , 2014, 49, 800-806.	1.0	16
17	Post-bronchiolitis wheezing is associated with toll-like receptor 9 rs187084 gene polymorphism. <i>Scientific Reports</i> , 2016, 6, 31165.	1.6	16
18	Using high-flow nasal cannulas for infants with bronchiolitis admitted to paediatric wards is safe and feasible. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2018, 107, 1971-1976.	0.7	16

#	ARTICLE	IF	CITATIONS
19	Obesity and bronchial obstruction in impulse oscillometry at age 5-7 years in a prospective post-bronchiolitis cohort. <i>Pediatric Pulmonology</i> , 2015, 50, 908-914.	1.0	14
20	Rapid detection of functional gene polymorphisms of TLRs and IL-17 using high resolution melting analysis. <i>Scientific Reports</i> , 2017, 7, 41522.	1.6	14
21	Toll-like receptor 1 and IL10 gene polymorphisms are linked to postbronchiolitis asthma in adolescence. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2018, 107, 134-139.	0.7	14
22	Impulse oscillometry at preschool age is a strong predictor of lung function by flow-volume spirometry in adolescence. <i>Pediatric Pulmonology</i> , 2018, 53, 552-558.	1.0	12
23	Haplotype of the Interleukin 17A gene is associated with osteitis after Bacillus Calmette-Guerin vaccination. <i>Scientific Reports</i> , 2017, 7, 11691.	1.6	11
24	Association of MBL2, TLR1, TLR2 and TLR6 Polymorphisms With Production of IFN- γ and IL-12 in BCG Osteitis Survivors R1. <i>Pediatric Infectious Disease Journal</i> , 2017, 36, 135-139.	1.1	11
25	Weight gain in infancy and post-bronchiolitis wheezing. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2012, 101, 38-42.	0.7	10
26	Toll-like receptor 2 subfamily genotypes are not associated with severity of bronchiolitis or postbronchiolitis wheezing in infants. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2013, 102, 1160-1164.	0.7	10
27	Excess weight in preschool children with a history of severe bronchiolitis is associated with asthma. <i>Pediatric Pulmonology</i> , 2015, 50, 424-430.	1.0	9
28	Low eosinophils during bronchiolitis in infancy are associated with lower risk of adulthood asthma. <i>Pediatric Allergy and Immunology</i> , 2015, 26, 668-673.	1.1	9
29	Interleukin 17A gene polymorphism rs2275913 is associated with osteitis after the Bacillus Calmette-Guérin vaccination. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2017, 106, 1837-1841.	0.7	9
30	IL-10 Gene Polymorphisms Are Associated with Post-Bronchiolitis Lung Function Abnormalities at Six Years of Age. <i>PLoS ONE</i> , 2015, 10, e0140799.	1.1	9
31	Toll-like receptor 4 polymorphisms were associated with low serum pro-inflammatory cytokines in BCG osteitis survivors. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2020, 109, 1417-1422.	0.7	8
32	IL33 rs1342326 gene variation is associated with allergic rhinitis at school age after infant bronchiolitis. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2020, 109, 2112-2116.	0.7	8
33	Following up infant bronchiolitis patients provided new evidence for and against the united airway disease hypothesis. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2016, 105, 1355-1360.	0.7	7
34	Interleukin-10 gene polymorphism rs1800896 is associated with post-bronchiolitis asthma at 11-13 years of age. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2019, 108, 2064-2069.	0.7	7
35	TLR5 rs5744174 gene polymorphism is associated with the virus etiology of infant bronchiolitis but not with post-bronchiolitis asthma. <i>Health Science Reports</i> , 2018, 1, e38.	0.6	5
36	IL17F rs763780 single nucleotide polymorphism is associated with asthma after bronchiolitis in infancy. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2021, 110, 222-227.	0.7	5

#	ARTICLE	IF	CITATIONS
37	<i>IL17A</i> gene polymorphisms rs4711998 and rs8193036 are not associated with postbronchiolitis asthma in Finnish children. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2018, 107, 1290-1291.	0.7	4
38	Polymorphisms in the promoter region of <i>IL10</i> gene are associated with virus etiology of infant bronchiolitis. <i>World Journal of Pediatrics</i> , 2018, 14, 594-600.	0.8	4
39	Toll-like receptor 10 rs4129009 gene polymorphism is associated with postbronchiolitis lung function in adolescence. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2020, 109, 1634-1641.	0.7	4
40	Interleukin-1 receptor-associated kinase-4 gene variation may increase postbronchiolitis asthma risk. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2021, 110, 952-958.	0.7	4
41	Interleukin-17 Receptor A gene polymorphism does not increase the risk of <i>Bacillus Calmette-Guérin</i> osteitis. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2020, 109, 1889-1890.	0.7	3
42	Interleukin 1 receptor-like 1 rs13408661/13431828 polymorphism is associated with persistent postbronchiolitis asthma at school age. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2022, 111, 628-635.	0.7	3
43	Preliminary communication suggests overweight was associated with reduced lung function in adolescence after infant bronchiolitis. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2019, 108, 1729-1730.	0.7	2
44	Risk factors for irreversible airway obstruction after infant bronchiolitis. <i>Respiratory Medicine</i> , 2021, 187, 106545.	1.3	2
45	Interferon- γ and interleukin-12 production in relation to gene polymorphisms in <i>Bacillus Calmette-Guérin</i> osteitis. <i>Pediatrics International</i> , 2019, 61, 982-987.	0.2	1
46	<i>IL17RA</i> variations showed no associations with postbronchiolitis asthma or lung function. <i>Pediatrics International</i> , 2021, 63, 196-201.	0.2	1
47	Toll-interacting protein polymorphisms in viral bronchiolitis outcomes. <i>Pediatrics International</i> , 2021, 63, 1103-1107.	0.2	1
48	Toll-like receptor 10 rs10004195 variation may be protective against <i>Bacillus Calmette-Guérin</i> osteitis after newborn vaccination. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2021, 110, 1585-1590.	0.7	1
49	Exploratory and confirmatory studies have different targets and both are needed in clinical research. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2018, 107, 734-735.	0.7	0
50	Interleukin-10 polymorphisms were not associated with lung function at age 11-13 years after infant bronchiolitis. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2020, 109, 198-199.	0.7	0
51	Genetic variations in Toll-like receptors 4 or 7 were not linked to postbronchiolitis lung function in adolescence. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2021, 110, 959-960.	0.7	0
52	Interleukin 17F polymorphisms showed no association with lung function at school age after infant bronchiolitis. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2021, 110, 219-221.	0.7	0
53	Interleukin 17F gene variations showed no association with BCG osteitis risk after newborn vaccination. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2021, 110, 618-623.	0.7	0
54	Variations of interleukin-1 receptor-associated kinase-4 encoding gene were not associated with postbronchiolitis lung function. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2021, 110, 1591-1593.	0.7	0

#	ARTICLE	IF	CITATIONS
55	IL33 rs1342326 polymorphism, though associated with severe post-bronchiolitis asthma, showed no association with lung function. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2021, 110, 2218-2220.	0.7	0
56	Interleukin 17A gene variations and lung function at school age after bronchiolitis in infancy. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2020, , .	0.7	0