Carlos Jaramillo

List of Publications by Citations

Source: https://exaly.com/author-pdf/3648575/carlos-jaramillo-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

170
papers7,848
citations41
h-index84
g-index180
ext. papers9,243
ext. citations5.3
avg, IF5.84
L-index

#	Paper	IF	Citations
170	Amazonia through time: Andean uplift, climate change, landscape evolution, and biodiversity. <i>Science</i> , 2010 , 330, 927-31	33.3	1362
169	Middle Miocene closure of the Central American Seaway. <i>Science</i> , 2015 , 348, 226-9	33.3	373
168	Biological evidence supports an early and complex emergence of the Isthmus of Panama. Proceedings of the National Academy of Sciences of the United States of America, 2015, 112, 6110-5	11.5	340
167	Sensitivity of leaf size and shape to climate: global patterns and paleoclimatic applications. <i>New Phytologist</i> , 2011 , 190, 724-39	9.8	334
166	Cenozoic plant diversity in the neotropics. <i>Science</i> , 2006 , 311, 1893-6	33.3	290
165	The Late Miocene paleogeography of the Amazon Basin and the evolution of the Amazon River system. <i>Earth-Science Reviews</i> , 2010 , 99, 99-124	10.2	240
164	Fracturing of the Panamanian Isthmus during initial collision with South America. <i>Geology</i> , 2011 , 39, 10)0 <i>7</i> -101	0202
163	Middle Eocene rodents from Peruvian Amazonia reveal the pattern and timing of caviomorph origins and biogeography. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2012 , 279, 1319-26	4.4	192
162	Effects of rapid global warming at the Paleocene-Eocene boundary on neotropical vegetation. <i>Science</i> , 2010 , 330, 957-61	33.3	188
161	Late Paleocene fossils from the Cerrejon Formation, Colombia, are the earliest record of Neotropical rainforest. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009 , 106, 18627-32	11.5	182
160	Post-Eocene climate change, niche conservatism, and the latitudinal diversity gradient of New World birds. <i>Journal of Biogeography</i> , 2006 , 33, 770-780	4.1	179
159	Fossil evidence for Cretaceous escalation in angiosperm leaf vein evolution. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 8363-6	11.5	151
158	Giant boid snake from the Palaeocene neotropics reveals hotter past equatorial temperatures. <i>Nature</i> , 2009 , 457, 715-7	50.4	151
157	A palynological zonation for the Cenozoic of the Llanos and Llanos Foothills of Colombia. <i>Palynology</i> , 2011 , 35, 46-84	1.5	108
156	An integrated analysis of an orogen-sedimentary basin pair: Latest Cretaceous-Cenozoic evolution of the linked Eastern Cordillera orogen and the Llanos foreland basin of Colombia. <i>Bulletin of the Geological Society of America</i> , 2008 , 120, 1171-1197	3.9	108
155	Tropical forest responses to increasing atmospheric CO: current knowledge and opportunities for future research. <i>Functional Plant Biology</i> , 2013 , 40, 531-551	2.7	97
154	A 60-million-year Cenozoic history of western Amazonian ecosystems in Contamana, eastern Peru. <i>Gondwana Research</i> , 2016 , 31, 30-59	5.1	90

(2010-2009)

153	Orogenic wedge advance in the northern Andes: Evidence from the Oligocene-Miocene sedimentary record of the Medina Basin, Eastern Cordillera, Colombia. <i>Bulletin of the Geological Society of America</i> , 2009 , 121, 780-800	3.9	88	
152	Miocene flooding events of western Amazonia. <i>Science Advances</i> , 2017 , 3, e1601693	14.3	77	
151	Consequences of shoaling of the Central American Seaway determined from modeling Nd isotopes. <i>Paleoceanography</i> , 2014 , 29, 176-189		71	
150	The Pliocene marine megafauna extinction and its impact on functional diversity. <i>Nature Ecology and Evolution</i> , 2017 , 1, 1100-1106	12.3	70	
149	Clockwise rotation of the Santa Marta massif and simultaneous Paleogene to Neogene deformation of the Plato-San Jorge and Cesar-Rancher basins. <i>Journal of South American Earth Sciences</i> , 2010 , 29, 832-848	2	7°	
148	The palynology of the Cerrejti formation (Upper Paleocene) of northern Colombia. <i>Palynology</i> , 2007 , 31, 153-189	1.5	67	
147	Insect leaf-chewing damage tracks herbivore richness in modern and ancient forests. <i>PLoS ONE</i> , 2014 , 9, e94950	3.7	64	
146	Testing geological models of evolution of the Isthmus of Panama in a phylogenetic framework. <i>Botanical Journal of the Linnean Society</i> , 2013 , 171, 287-300	2.2	59	
145	First North American fossil monkey and early Miocene tropical biotic interchange. <i>Nature</i> , 2016 , 533, 243-6	50.4	59	
144	Comment (1) on "Formation of the Isthmus of Panama" by O'Dea. <i>Science Advances</i> , 2017 , 3, e1602321	14.3	57	
143	Early Paleogene magmatism in the northern Andes: Insights on the effects of Oceanic PlateauBontinent convergence. <i>Earth and Planetary Science Letters</i> , 2012 , 331-332, 97-111	5.3	55	
142	Global Warming and Neotropical Rainforests: A Historical Perspective. <i>Annual Review of Earth and Planetary Sciences</i> , 2013 , 41, 741-766	15.3	54	
141	Responses of legume versus nonlegume tropical tree seedlings to elevated CO2 concentration. <i>Plant Physiology</i> , 2011 , 157, 372-85	6.6	54	
140	Palms (Arecaceae) from a Paleocene rainforest of northern Colombia. <i>American Journal of Botany</i> , 2009 , 96, 1300-12	2.7	53	
139	Systematics and biogeography of crocodylians from the Miocene of Panama. <i>Journal of Vertebrate Paleontology</i> , 2013 , 33, 239-263	1.7	52	
138	Paleocene Malvaceae from northern South America and their biogeographical implications. <i>American Journal of Botany</i> , 2011 , 98, 1337-55	2.7	49	
137	Geographical distribution patterns of Carcharocles megalodon over time reveal clues about extinction mechanisms. <i>Journal of Biogeography</i> , 2016 , 43, 1645-1655	4.1	48	
136	The eastern foothills of the Eastern Cordillera of Colombia: An example of multiple factors controlling structural styles and active tectonics. <i>Bulletin of the Geological Society of America</i> , 2010 , 122, 1946, 1964	3.9	46	

135	Response of tropical vegetation to Paleogene warming. <i>Paleobiology</i> , 2002 , 28, 222-243	2.6	45
134	Rapid regional surface uplift of the northern Altiplano plateau revealed by multiproxy paleoclimate reconstruction. <i>Earth and Planetary Science Letters</i> , 2016 , 447, 33-47	5.3	45
133	Neotropical mammal diversity and the Great American Biotic Interchange: spatial and temporal variation in South America's fossil record. <i>Frontiers in Genetics</i> , 2014 , 5, 451	4.5	44
132	Early-subduction-related orogeny in the northern Andes: Turonian to Eocene magmatic and provenance record in the Santa Marta Massif and Rancheria Basin, northern Colombia. <i>Terra Nova</i> , 2011 , 23, 26-34	3	43
131	Paratropical floral extinction in the Late Palaeocene Early Eocene. <i>Journal of the Geological Society</i> , 2007 , 164, 323-332	2.7	43
130	Sequence stratigraphic interpretations from palynofacies, dinocyst and lithological data of Upper Eocenellower Oligocene strata in southern Mississippi and Alabama, U.S. Gulf Coast. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 1999 , 145, 259-302	2.9	41
129	A new small short-snouted dyrosaurid (Crocodylomorpha, Mesoeucrocodylia) from the Paleocene of northeastern Colombia. <i>Journal of Vertebrate Paleontology</i> , 2010 , 30, 139-162	1.7	40
128	Intraplate subsidence and basin filling adjacent to an oceanic arctontinent collision: a case from the southern Caribbean-South America plate margin. <i>Basin Research</i> , 2011 , 23, 403-422	3.2	39
127	Revised stratigraphy of Neogene strata in the Cocinetas Basin, La Guajira, Colombia. <i>Swiss Journal of Palaeontology</i> , 2015 , 134, 5-43	1.2	37
126	Fossil Araceae from a Paleocene neotropical rainforest in Colombia. <i>American Journal of Botany</i> , 2008 , 95, 1569-83	2.7	37
125	Is the trace fossil Macaronichnus an indicator of temperate to cold waters? Exploring the paradox of its occurrence in tropical coasts. <i>Geology</i> , 2010 , 38, 651-654	5	36
124	Extinction at the end-Cretaceous and the origin of modern Neotropical rainforests. <i>Science</i> , 2021 , 372, 63-68	33.3	36
123	Onset of fault reactivation in the Eastern Cordillera of Colombia and proximal Llanos Basin; response to CaribbeanBouth American convergence in early Palaeogene time. <i>Geological Society Special Publication</i> , 2013 , 377, 285-314	1.7	35
122	An integrative geologic, geochronologic and geochemical study of Gorgona Island, Colombia: Implications for the formation of the Caribbean Large Igneous Province. <i>Earth and Planetary Science Letters</i> , 2011 , 309, 324-336	5.3	35
121	New podocnemidid turtle (Testudines: Pleurodira) from the middlelipper Paleocene of South America. <i>Journal of Vertebrate Paleontology</i> , 2010 , 30, 367-382	1.7	34
120	Menispermaceae from the Cerrejon Formation, middle to late Paleocene, Colombia. <i>American Journal of Botany</i> , 2008 , 95, 954-73	2.7	34
119	Tectonic controls on Cenozoic foreland basin development in the north-eastern Andes, Colombia. <i>Basin Research</i> , 2010 , 22, 874	3.2	33
118	A new longirostrine dyrosaurid (Crocodylomorpha, Mesoeucrocodylia) from the Paleocene of north-eastern Colombia: biogeographic and behavioural implications for New-World Dyrosauridae. <i>Palaeontology</i> , 2011 , 54, 1095-1116	2.9	32

(2010-2012)

117	The paleogene synorogenic succession in the northwestern Maracaibo block: Tracking intraplate uplifts and changes in sediment delivery systems. <i>Journal of South American Earth Sciences</i> , 2012 , 39, 93-111	2	31
116	The final phase of tropical lowland conditions in the axial zone of the Eastern Cordillera of Colombia: Evidence from three palynological records. <i>Journal of South American Earth Sciences</i> , 2012 , 39, 157-169	2	31
115	Phytogeographic History and Phylogeny of the Humiriaceae. <i>International Journal of Plant Sciences</i> , 2010 , 171, 392-408	2.6	31
114	Permineralized fruits from the late Eocene of Panama give clues of the composition of forests established early in the uplift of Central America. <i>Review of Palaeobotany and Palynology</i> , 2012 , 175, 10-24	1.7	30
113	Phytogeographic implications of fossil endocarps of Menispermaceae from the Paleocene of Colombia. <i>American Journal of Botany</i> , 2011 , 98, 2004-17	2.7	30
112	Neogene molluscs, shallow marine paleoenvironments, and chronostratigraphy of the Guajira Peninsula, Colombia. <i>Swiss Journal of Palaeontology</i> , 2015 , 134, 45-75	1.2	29
111	Reply to Lessios and Marko et al.: Early and progressive migration across the Isthmus of Panama is robust to missing data and biases. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, E5767-8	11.5	29
110	New turtles (Chelonia) from the late Eocene through late Miocene of the Panama Canal Basin. <i>Journal of Paleontology</i> , 2012 , 86, 539-557	1.1	28
109	A new Late Miocene chondrichthyan assemblage from the Chagres Formation, Panama. <i>Journal of South American Earth Sciences</i> , 2015 , 60, 56-70	2	27
108	Carnivorans at the Great American Biotic Interchange: new discoveries from the northern neotropics. <i>Die Naturwissenschaften</i> , 2014 , 101, 965-74	2	27
107	New pelomedusoid turtles from the late Palaeocene Cerrej Formation of Colombia and their implications for phylogeny and body size evolution. <i>Journal of Systematic Palaeontology</i> , 2012 , 10, 313-	3347	27
106	Late Eocene marine incursion in north-western South America. <i>Palaeogeography, Palaeoclimatology, Palaeoecology,</i> 2008 , 264, 140-146	2.9	27
105	Sharks and rays (Chondrichthyes, Elasmobranchii) from the late Miocene Gatun Formation of Panama. <i>Journal of Paleontology</i> , 2013 , 87, 755-774	1.1	26
104	Early Miocene chondrichthyans from the Culebra Formation, Panama: A window into marine vertebrate faunas before closure the Central American Seaway. <i>Journal of South American Earth Sciences</i> , 2013 , 42, 159-170	2	26
103	Ariid sea catfishes from the coeval Pirabas (Northeastern Brazil), Cantaure, Castillo (Northwestern Venezuela), and Castilletes (North Colombia) formations (early Miocene), with description of three new species. <i>Swiss Journal of Palaeontology</i> , 2013 , 132, 45-68	1.2	25
102	Comment on "The Response of Vegetation on the Andean Flank in Western Amazonia to Pleistocene Climate Change". <i>Science</i> , 2011 , 333, 1825-1825	33.3	25
101	Palaeontological evidence for the last temporal occurrence of the ancient western Amazonian river outflow into the Caribbean. <i>PLoS ONE</i> , 2013 , 8, e76202	3.7	24
100	Extinct peccary Lynorcaloccidentale (Tayassuidae, Tayassuinae) from the Miocene of Panama and correlations to North America. <i>Journal of Paleontology</i> , 2010 , 84, 288-298	1.1	24

99	Fossil Crocodilians from the High Guajira Peninsula of Colombia: Neogene faunal change in northernmost South America. <i>Journal of Vertebrate Paleontology</i> , 2016 , 36, e1110586	1.7	24
98	Fruits of an Dld WorldItribe (Phytocreneae; Icacinaceae) from the Paleogene of North and South America. <i>Systematic Botany</i> , 2012 , 37, 784-794	0.7	23
97	A fruit and leaves of Rhamnaceous affinities from the late Cretaceous (Maastrichtian) of Colombia. <i>American Journal of Botany</i> , 2010 , 97, 71-9	2.7	23
96	The Origin and Diversification of the Hyperdiverse Flora in the ChocBiogeographic Region. <i>Frontiers in Plant Science</i> , 2019 , 10, 1328	6.2	23
95	New bothremydid turtle (Testudines, Pleurodira) from the Paleocene of northeastern Colombia. <i>Journal of Paleontology</i> , 2012 , 86, 688-698	1.1	22
94	A New Early Miocene (Aquitanian) Elasmobranchii Assemblage from the la Guajira Peninsula, Colombia. <i>Ameghiniana</i> , 2016 , 53, 77	0.9	22
93	Exceptional preservation of mid-Cretaceous marine arthropods and the evolution of novel forms via heterochrony. <i>Science Advances</i> , 2019 , 5, eaav3875	14.3	21
92	A new blunt-snouted dyrosaurid, Anthracosuchus balrogus gen. et sp. nov. (Crocodylomorpha, Mesoeucrocodylia), from the Palaeocene of Colombia. <i>Historical Biology</i> , 2015 , 27, 998-1020	1.1	21
91	Origins of BiodiversityResponse. <i>Science</i> , 2011 , 331, 399-400	33.3	21
90	The oldest frog crabs (Decapoda: Brachyura: Raninoida) from the Aptian of northern South America. <i>Journal of Crustacean Biology</i> , 2012 , 32, 405-420	0.8	20
89	Fossil evidence for earliest Neogene American faunal interchange: Boa (Serpentes, Boinae) from the early Miocene of Panama. <i>Journal of Vertebrate Paleontology</i> , 2012 , 32, 1328-1334	1.7	20
88	MARINE PALEOENVIRONMENTS OF MIOCENE-PLIOCENE FORMATIONS OF NORTH-CENTRAL FALCON STATE, VENEZUELA. <i>Journal of Foraminiferal Research</i> , 2010 , 40, 266-282	1.1	20
87	Mining threatens Colombian ecosystems. <i>Science</i> , 2018 , 359, 1475	33.3	18
86	Insights into the Neotropics prior to the Great American Biotic Interchange: new evidence of mammalian predators from the Miocene of Northern Colombia. <i>Journal of Vertebrate Paleontology</i> , 2016 , 36, e1029581	1.7	18
85	Fossil woods (Malvaceae) from the lower Miocene (early to mid-Burdigalian) part of the Cucaracha Formation of Panama (Central America) and their biogeographic implications. <i>Review of Palaeobotany and Palynology</i> , 2014 , 209, 11-34	1.7	18
84	Early to Middle Miocene Turtles from the Northernmost Tip of South America: Giant Testudinids, Chelids, and Podocnemidids from the Castilletes Formation, Colombia. <i>Ameghiniana</i> , 2015 , 52, 188-203	0.9	17
83	First Central American record of Anthracotheriidae (Mammalia, Bothriodontinae) from the early Miocene of Panama. <i>Journal of Vertebrate Paleontology</i> , 2013 , 33, 421-433	1.7	17
82	U-Pb LA-ICP-MS GEOCHRONOLOGY AND GEOCHEMISTRY OF JURASSIC VOLCANIC AND PLUTONIC ROCKS FROM THE PUTUMAYO REGION (SOUTHERN COLOMBIA): TECTONIC SETTING AND REGIONAL CORRELATIONS. <i>Boletin De Geologia</i> 2016 , 38, 21-38	0.4	17

81	Preface: La Guajira, Colombia: a new window into the Cenozoic neotropical biodiversity and the Great American Biotic Interchange. <i>Swiss Journal of Palaeontology</i> , 2015 , 134, 1-4	1.2	16
80	Climate change during the Early Paleogene in the BogotBasin (Colombia) inferred from paleosol carbon isotope stratigraphy, major oxides, and environmental magnetism. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2013 , 388, 115-127	2.9	16
79	Tracing the fossil pollen record of Hedyosmum (Chloranthaceae), an old lineage with recent Neotropical diversification. <i>Grana</i> , 2013 , 52, 161-180	0.8	16
78	Early Miocene CO estimates from a Neotropical fossil leaf assemblage exceed 400 ppm. <i>American Journal of Botany</i> , 2018 , 105, 1929-1937	2.7	16
77	Magmatic evolution of Panama Canal volcanic rocks: A record of arc processes and tectonic change. <i>PLoS ONE</i> , 2017 , 12, e0176010	3.7	15
76	Variations in angiosperm leaf vein density have implications for interpreting life form in the fossil record. <i>Geology</i> , 2014 , 42, 919-922	5	15
75	Marine mammals from the Miocene of Panama. Journal of South American Earth Sciences, 2010, 30, 167-	1275	15
74	New floridatragulines (Mammalia, Camelidae) from the early Miocene Las Cascadas Formation, Panama. <i>Journal of Vertebrate Paleontology</i> , 2012 , 32, 456-475	1.7	15
73	New material of Chelus colombiana (Testudines; Pleurodira) from the lower Miocene of Colombia. Journal of Vertebrate Paleontology, 2008 , 28, 1206-1212	1.7	15
72	Endemic palm species shed light on habitat shifts and the assembly of the Cerrado and Restinga floras. <i>Molecular Phylogenetics and Evolution</i> , 2017 , 110, 127-133	4.1	14
71	A New Pliocene Capybara (Rodentia, Caviidae) from Northern South America (Guajira, Colombia), and its Implications for the Great American Biotic Interchange. <i>Journal of Mammalian Evolution</i> , 2017 , 24, 111-125	2.2	14
70	HUESSER HORIZON: A LAKE AND A MARINE INCURSION IN NORTHWESTERN SOUTH AMERICA DURING THE EARLY MIOCENE. <i>Palaios</i> , 2009 , 24, 199-210	1.6	14
69	U/Pb LA-MC-ICP-MS Zircon Geochronology and Geochemistry from a Postcollisional Biotite Granite of the Baja Guajira Basin, Colombia: Implications for Late Cretaceous and Neogene CaribbeanBouth American Tectonics. <i>Journal of Geology</i> , 2009 , 117, 685-692	2	14
68	A molecular evaluation of bulk organic carbon-isotope chemostratigraphy for terrestrial correlations: An example from two Paleocene Eocene tropical sequences. <i>Palaeogeography, Palaeoclimatology, Palaeoecology,</i> 2009 , 277, 173-183	2.9	14
67	Disproportionate extinction of South American mammals drove the asymmetry of the Great American Biotic Interchange. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 26281-26287	11.5	14
66	A Late Cretaceous Piper (Piperaceae) from Colombia and diversification patterns for the genus. <i>American Journal of Botany</i> , 2015 , 102, 273-89	2.7	13
65	Two tropical conifers show strong growth and water-use efficiency responses to altered CO2 concentration. <i>Annals of Botany</i> , 2016 , 118, 1113-1125	4.1	13
64	New Miocene Caribbean gavialoids and patterns of longirostry in crocodylians. <i>Journal of Systematic Palaeontology</i> , 2019 , 17, 1049-1075	2.3	13

63	Early Eocene Spore and Pollen Assemblages from the Laguna del Hunco Fossil Lake Beds, Patagonia, Argentina. <i>International Journal of Plant Sciences</i> , 2020 , 181, 594-615	2.6	12
62	Paleoenvironmental reconstruction for the lower Pliocene Arroyo Piedras section (Tubar Delenation of South American Earth Sciences, 2012 , 39, 170-183	2	11
61	Late Eocene-Early Oligocene paleofloristic patterns in southern Mississippi and Alabama, US Gulf Coast. <i>Review of Palaeobotany and Palynology</i> , 1996 , 91, 23-34	1.7	11
60	Neogene Proto-Caribbean porcupinefishes (Diodontidae). <i>PLoS ONE</i> , 2017 , 12, e0181670	3.7	11
59	Palaeontological framework from Pirabas Formation (North Brazil) used as potential model for equatorial carbonate platform. <i>Marine Micropaleontology</i> , 2020 , 154, 101813	1.7	11
58	Neogene sloth assemblages (Mammalia, Pilosa) of the Cocinetas Basin (La Guajira, Colombia): implications for the Great American Biotic Interchange. <i>Palaeontology</i> , 2016 , 59, 563-582	2.9	11
57	Paleogene Salvinia (Salviniaceae) from Colombia and their paleobiogeographic implications. <i>Review of Palaeobotany and Palynology</i> , 2017 , 246, 85-108	1.7	10
56	Phytogeographic History of the Humiriaceae (Part 2). <i>International Journal of Plant Sciences</i> , 2014 , 175, 828-840	2.6	10
55	Palynological composition of a Lower Cretaceous South American tropical sequence: climatic implications and diversity comparisons with other latitudes. <i>American Journal of Botany</i> , 2012 , 99, 1819	- 2 7	10
54	Improving the taxonomy of fossil pollen using convolutional neural networks and superresolution microscopy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 28496-28505	11.5	10
53	Neogene precipitation, vegetation, and elevation history of the Central Andean Plateau. <i>Science Advances</i> , 2020 , 6, eaaz4724	14.3	10
52	Canopy structure in Late Cretaceous and Paleocene forests as reconstructed from carbon isotope analyses of fossil leaves. <i>Geology</i> , 2019 , 47, 977-981	5	9
51	Biodiversity only makes sense in the light of evolution. <i>Journal of Biosciences</i> , 2014 , 39, 333-7	2.3	9
50	Quantitative analysis of Cenozoic palynofloras from Patagonia, southern South America. <i>Palynology</i> , 2013 , 37, 246-258	1.5	9
49	Biotic community and landscape changes around the Eocene®ligocene transition at Shapaja, Peruvian Amazonia: Regional or global drivers?. <i>Global and Planetary Change</i> , 2021 , 202, 103512	4.2	9
48	New early Miocene protoceratids (Mammalia, Artiodactyla) from Panama. <i>Journal of Vertebrate Paleontology</i> , 2015 , 35, e970688	1.7	8
47	Mangrove Distribution during the Holocene in TribuglGulf, Colombia1. <i>Biotropica</i> , 2000 , 32, 14-22	2.3	8
46	Middle to Late Paleocene Leguminosae fruits and leaves from Colombia. <i>Australian Systematic Botany</i> , 2019 ,	1	7

(2011-2014)

45	Paleocene wind-dispersed fruits and seeds from Colombia and their implications for early Neotropical rainforests. <i>Acta Palaeobotanica</i> , 2014 , 54, 197-229	0.9	7	
44	New observations and reinterpretation on the enigmatic taxon Colombitherium (?Pyrotheria, Mammalia) from Colombia. <i>Palaeontology</i> , 2010 , 53, 319-325	2.9	7	
43	Comment on "The response of vegetation on the Andean flank in western Amazonia to Pleistocene climate change". <i>Science</i> , 2011 , 333, 1825; author reply 1825	33.3	7	
42	Trans-Amazon Drilling Project (TADP): origins and evolution of the forests, climate, and hydrology of the South American tropics. <i>Scientific Drilling</i> ,20, 41-49		7	
41	The Neogene Record of Northern South American Native Ungulates. <i>Smithsonian Contributions To Paleobiology</i> , 2018 , iv-67		7	
40	Drastic Vegetation Change in the Guajira Peninsula (Colombia) During the Neogene. <i>Paleoceanography and Paleoclimatology</i> , 2020 , 35, e2020PA003933	3.3	7	
39	Selective extinction against redundant species buffers functional diversity. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2020 , 287, 20201162	4.4	7	
38	The Pliocene B leistocene palynology of the Negro River, Brazil. <i>Palynology</i> , 2019 , 43, 223-243	1.5	7	
37	Unravelling the widening of the earliest Andean northern orogen: Maastrichtian to early Eocene intra-basinal deformation in the northern Eastern Cordillera of Colombia. <i>Basin Research</i> , 2021 , 33, 809-	-845	7	
36	Provenance analysis of the Pliocene Ware Formation in the Guajira Peninsula, northern Colombia: Paleodrainage implications. <i>Journal of South American Earth Sciences</i> , 2018 , 81, 66-77	2	7	
35	Sedimentology and Palynostratigraphy of a Pliocene-Pleistocene (Piacenzian to Gelasian) deposit in the lower Negro River: Implications for the establishment of large rivers in Central Amazonia. <i>Journal of South American Earth Sciences</i> , 2017 , 79, 215-229	2	6	
34	Karatophyllum bromelioides L.D. Gomez revisited: a probable fossil CAM bromeliad. <i>American Journal of Botany</i> , 2011 , 98, 1905-8	2.7	6	
33	Probabilistic correlation of single stratigraphic samples: A generalized approach for biostratigraphic data. <i>AAPG Bulletin</i> , 2012 , 96, 235-244	2.5	6	
32	Early Miocene marine palynology of the Colombian Caribbean Margin: biostratigraphic and paleoceanographic implications. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2020 , 558, 109955	5 ^{2.9}	6	
31	19-Million-Year-Old Spondioid Fruits from Panama Reveal a Dynamic Dispersal History for Anacardiaceae. <i>International Journal of Plant Sciences</i> , 2019 , 180, 479-492	2.6	5	
30	The Biogeography of the Araucarian Dispersed PollenCyclusphaera. <i>International Journal of Plant Sciences</i> , 2013 , 174, 489-498	2.6	5	
29	A neotropical Miocene pollen database employing image-based search and semantic modeling. <i>Applications in Plant Sciences</i> , 2014 , 2, 1400030	2.3	5	
28	The Origin of the Modern Amazon Rainforest: Implications of the Palynological and Palaeobotanical Record 2011 , 317-334		5	

27	New records of Humiriaceae fossil fruits from the Oligocene and Early Miocene of the western Azuero Peninsula, Panam\(\textit{Boletin De La Sociedad Geologica Mexicana, 2018}\), 70, 223-239	1.7	5
26	Early Records of Melastomataceae from the Middlellate Paleocene Rain Forests of South America Conflict with Laurasian Origins. <i>International Journal of Plant Sciences</i> , 2021 , 182, 401-412	2.6	5
25	The search for an elusive worm in the tropics, the past as a key to the present, and reverse uniformitarianism. <i>Scientific Reports</i> , 2019 , 9, 18402	4.9	5
24	Anllsis de secuencias y procedencia EN sucesiones sedimentarias de grano fino: un ejemplo de la Formacili Umir y base de la Formacili Lisama, en el sector de Simacota (Santander, Colombia). <i>Boletin De Geologia</i> , 2016 , 38, 51-72	0.4	4
23	Miocene heterozoan carbonate systems from the western Atlantic equatorial margin in South America: The Pirabas formation. <i>Sedimentary Geology</i> , 2020 , 407, 105739	2.8	4
22	Isotope sclerochronology indicates enhanced seasonal precipitation in northern South America (Colombia) during the Mid-Miocene Climatic Optimum. <i>Geology</i> , 2020 , 48, 668-672	5	4
21	Compositional and diversity comparisons between the palynological records of the Neogene (Solim Formation) and Holocene sediments of Western Amazonia. <i>Palynology</i> , 2021 , 45, 3-14	1.5	4
20	Floristic and climatic reconstructions of two Lower Cretaceous successions from Peru. <i>Palynology</i> , 2018 , 42, 420-433	1.5	3
19	First report of Hadrosia Cooper, 1983 in South America and its biostratigraphical and palaeobiogeographical implications. <i>Cretaceous Research</i> , 2012 , 34, 257-267	1.8	3
18	The Tertiary Fusagasugsuccession; A Record Of The Complex Latest Cretaceous-Pre-Miocene Deformation In An Area Between The Magdalena Valley And Sabana De Bogots 2003,		3
17	Miocene Freshwater Dolphins from La Venta, Huila, Colombia Suggest Independent Invasions of Riverine Environments in Tropical South America. <i>Journal of Vertebrate Paleontology</i> , 2020 , 40, e18120	78 ^{.7}	3
16	Phylogenetic and ecological correlates of pollen morphological diversity in a Neotropical rainforest. <i>Biotropica</i> , 2021 , 53, 74-85	2.3	3
15	Aquatic ecosystems in a newly formed ecospace: Early Pliocene lakes in the Central Andean Altiplano. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2018 , 490, 218-226	2.9	3
14	Long-term topographic growth and decay constrained by 3D thermo-kinematic modeling: Tectonic evolution of the Antioquia Altiplano, Northern Andes. <i>Global and Planetary Change</i> , 2021 , 203, 103553	4.2	3
13	Palynostratigraphy of the Ramon and Solim\(\text{B} \) formations in the Acre Basin, Brazil. <i>Journal of South American Earth Sciences</i> , 2020 , 103, 102720	2	2
12	A Pliocene-Pleistocene continental biota from Venezuela. <i>Swiss Journal of Palaeontology</i> , 2021 , 140, 9	1.2	2
11	Pollen morphology of the Amacayacu Forest dynamics plot, Western Amazon, Colombia. <i>Palynology</i> , 2020 , 44, 32-79	1.5	2
10	Paleoclimatic and paleoecological reconstruction of a middle to late Eocene South American tropical dry forest. <i>Global and Planetary Change</i> , 2021 , 205, 103617	4.2	2

LIST OF PUBLICATIONS

9	3D models related to the publication: Neogene sloth assemblages (Mammalia, Pilosa) of the Cocinetas Basin (La Guajira, Colombia): implications for the Great American Biotic Interchange. MorphoMuseuM, 2016 , 2, e3	0.1	1
8	Increased megathrust shear force drives topographic uplift in the Colombian coastal forearc. <i>Tectonophysics</i> , 2021 , 820, 229132	3.1	1
7	A new Miocene turtle from Colombia sheds light on the evolutionary history of the extant genus Mesoclemmys Gray, 1873. <i>Journal of Vertebrate Paleontology</i> , 2019 , 39, e1716777	1.7	1
6	New acritarch and peridinioid dinoflagellate cyst species from the Oligocene Miocene of Colombia. <i>Review of Palaeobotany and Palynology</i> , 2021 , 290, 104427	1.7	О
5	Mid to late Holocene sea-level rise and precipitation variability recorded in the fringe mangroves of the Caribbean coast of Panama. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2022 , 592, 110918	8 ^{2.9}	О
4	Tryptophan usage by Helicobacter pylori differs among strains. <i>Scientific Reports</i> , 2019 , 9, 873	4.9	
3	(087 D 90) Proposal to treat the use of a hyphen in the name of a fossil-genus as an orthographical error. <i>Taxon</i> , 2015 , 64, 863-863	0.8	
2	Unraveling the record of a tropical continental Cretaceous-Paleogene boundary in northern Colombia, South America. <i>Journal of South American Earth Sciences</i> , 2022 , 114, 103717	2	
1	Effects of consecutive dry and wet days on the forestBavanna boundary in north-west South America. <i>Global Ecology and Biogeography</i> , 2022 , 31, 347	6.1	