
Nicola Lamberti

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/365280/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Exercise in Patients on Dialysis: A Multicenter, Randomized Clinical Trial. Journal of the American Society of Nephrology: JASN, 2017, 28, 1259-1268.	3.0	272
2	Effects of low-intensity endurance and resistance training on mobility in chronic stroke survivors: a pilot randomized controlled study. European Journal of Physical and Rehabilitation Medicine, 2017, 53, 228-239.	1.1	49
3	Robot-assisted gait training is not superior to intensive overground walking in multiple sclerosis with severe disability (the RAGTIME study): A randomized controlled trial. Multiple Sclerosis Journal, 2020, 26, 716-724.	1.4	43
4	The Role of Deconditioning in the End-Stage Renal Disease Myopathy: Physical Exercise Improves Altered Resting Muscle Oxygen Consumption. American Journal of Nephrology, 2015, 41, 329-336.	1.4	41
5	Impact of renal dysfunction on in-hospital mortality of patients with severe chronic obstructive pulmonary disease: a single-center Italian study. International Urology and Nephrology, 2016, 48, 1121-1127.	0.6	37
6	Effects of a Robot-Assisted Arm Training Plus Hand Functional Electrical Stimulation on Recovery After Stroke: A Randomized Clinical Trial. Archives of Physical Medicine and Rehabilitation, 2020, 101, 309-316.	0.5	29
7	Effectiveness of blood flowâ€restricted slow walking on mobility in severe multiple sclerosis: A pilot randomized trial. Scandinavian Journal of Medicine and Science in Sports, 2020, 30, 1999-2009.	1.3	28
8	A Toe Flexion NIRS assisted Test for Rapid Assessment of Foot Perfusion in Peripheral Arterial Disease: Feasibility, Validity, and Diagnostic Accuracy. European Journal of Vascular and Endovascular Surgery, 2017, 54, 187-194.	0.8	25
9	Reliability of the Vascular Claudication Reporting in Diabetic Patients With Peripheral Arterial Disease. Angiology, 2015, 66, 365-374.	0.8	24
10	Changes in exercise capacity and risk of all-cause mortality in patients with peripheral artery disease: a 10-year retrospective cohort study. Internal and Emergency Medicine, 2020, 15, 289-298.	1.0	22
11	Muscle oxygen consumption by NIRS and mobility in multiple sclerosis patients. BMC Neurology, 2013, 13, 52.	0.8	21
12	Gender Differences in Outcomes Following a Pain-Free, Home-Based Exercise Program for Claudication. Journal of Women's Health, 2019, 28, 1313-1321.	1.5	21
13	Respiratory muscle impairment in dialysis patients: can minimal dose of exercise limit the damage? A Preliminary study in a sample of patients enrolled in the EXCITE trial. Journal of Nephrology, 2016, 29, 863-869.	0.9	20
14	Effects of a "test in-train out" walking program versus supervised standard rehabilitation in chronic stroke patients: a feasibility and pilot randomized study. European Journal of Physical and Rehabilitation Medicine, 2016, 52, 279-87.	1.1	19
15	Structured Home-Based Exercise Versus Invasive Treatment. Angiology, 2016, 67, 772-780.	0.8	18
16	The effectiveness of Robot-Assisted Gait Training versus conventional therapy on mobility in severely disabled progressive MultiplE sclerosis patients (RAGTIME): study protocol for a randomized controlled trial. Trials, 2017, 18, 88.	0.7	18
17	Biomarkers of Muscle Metabolism in Peripheral Artery Disease: A Dynamic NIRS-Assisted Study to Detect Adaptations Following Revascularization and Exercise Training. Diagnostics, 2020, 10, 312.	1.3	16
18	Rehabilitative Exercise Reduced the Impact of Peripheral Artery Disease on Vascular Outcomes in Elderly Patients with Claudication: A Three-Year Single Center Retrospective Study. Journal of Clinical Medicine, 2019, 8, 210.	1.0	14

Nicola Lamberti

#	Article	lF	CITATIONS
19	Infrared Thermography as a Diagnostic Tool for the Assessment of Patients with Symptomatic Peripheral Arterial Disease Undergoing Infrafemoral Endovascular Revascularisations. Diagnostics, 2021, 11, 1701.	1.3	13
20	Acute oxygenation changes on ischemic foot of a novel intermittent pneumatic compression device and of an existing sequential device in severe peripheral arterial disease. BMC Cardiovascular Disorders, 2014, 14, 40.	0.7	12
21	Rehabilitation Improves Mitochondrial Energetics in Progressive Multiple Sclerosis: The Significant Role of Robot-Assisted Gait Training and of the Personalized Intensity. Diagnostics, 2020, 10, 834.	1.3	12
22	Altered erythroidâ€related miRNA levels as a possible novel biomarker for detection of autologous blood transfusion misuse in sport. Transfusion, 2019, 59, 2709-2721.	0.8	11
23	Don't stop walking: the in-home rehabilitation program for peripheral artery disease patients during the COVID-19 pandemic. Internal and Emergency Medicine, 2021, 16, 1307-1315.	1.0	11
24	Construct Validity and Responsiveness of the COVID-19 Yorkshire Rehabilitation Scale (C19-YRS) in a Cohort of Italian Hospitalized COVID-19 Patients. International Journal of Environmental Research and Public Health, 2022, 19, 6696.	1.2	11
25	Changes in hemoglobin profile reflect autologous blood transfusion misuse in sports. Internal and Emergency Medicine, 2018, 13, 517-526.	1.0	10
26	Structured pain-free exercise progressively improves ankle-brachial index and walking ability in patients with claudication and compressible arteries: an observational study. Internal and Emergency Medicine, 2022, 17, 439-449.	1.0	8
27	Home-based exercise for elderly patients with intermittent claudication limited by osteoarticular disorders – feasibility and effectiveness of a low-intensity programme. Vasa - European Journal of Vascular Medicine, 2018, 47, 227-234.	0.6	8
28	Impact of comorbidity on outcome in kidney transplant recipients: a retrospective study in Italy. Internal and Emergency Medicine, 2016, 11, 825-832.	1.0	7
29	Functional recovery in multiple sclerosis patients undergoing rehabilitation programs is associated with plasma levels of hemostasis inhibitors. Multiple Sclerosis and Related Disorders, 2020, 44, 102319.	0.9	7
30	Lack of a "Weekend Effect―for Renal Transplant Recipients. Angiology, 2017, 68, 366-373.	0.8	6
31	Effectiveness of Home-Based Pain-Free Exercise versus Walking Advice in Patients with Peripheral Artery Disease: A Randomized Controlled Trial. Methods and Protocols, 2021, 4, 29.	0.9	6
32	Beneficial Effects of Robot-Assisted Gait Training on Functional Recovery in Women after Stroke: A Cohort Study. Medicina (Lithuania), 2021, 57, 1200.	0.8	6
33	Antidoping attitudes among elite athletes: a cross sectional study in biathlon using a suitably developed questionnaire. Journal of Sports Medicine and Physical Fitness, 2017, 57, 610-623.	0.4	5
34	A Personalized Patient-Centered Intervention to Empower through Physical Activity the Patient in the Dialysis Center: Study Protocol for a Pragmatic Nonrandomized Clinical Trial. Methods and Protocols, 2020, 3, 83.	0.9	5
35	Combining a supervised and home-based task-oriented circuit training improves walking endurance in patients with multiple sclerosis. The MS_TOCT randomized-controlled trial. Multiple Sclerosis and Related Disorders, 2022, 60, 103721.	0.9	5
36	The legacy effect of a home walking exercise program in kidney failure patients on dialysis. Nephrology Dialysis Transplantation, 0, , .	0.4	5

NICOLA LAMBERTI

#	Article	IF	CITATIONS
37	Performance Assessment of Patient on Dialysis. Kidney and Blood Pressure Research, 2014, 39, 176-179.	0.9	4
38	Weekend Effect and in-Hospital Mortality in Elderly Patients with Acute Kidney Injury: A Retrospective Analysis of a National Hospital Database in Italy. Journal of Clinical Medicine, 2020, 9, 1815.	1.0	4
39	Renal Transplantation and Outcome. Progress in Transplantation, 2016, 26, 397-398.	0.4	3
40	Baseline and overtime variations of soluble adhesion molecule plasma concentrations are associated with mobility recovery after rehabilitation in multiple sclerosis patients. Journal of Neuroimmunology, 2021, 352, 577473.	1.1	3
41	Motor Cortical Activation Assessment in Progressive Multiple Sclerosis Patients Enrolled in Gait Rehabilitation: A Secondary Analysis of the RAGTIME Trial Assisted by Functional Near-Infrared Spectroscopy. Diagnostics, 2021, 11, 1068.	1.3	3
42	Planning the International Competition Schedules for the Health of Elite Athletes: A 21-Year Retrospective Study Evaluating the Effectiveness and Economic Impact in an Olympic Sport. PLoS ONE, 2015, 10, e0130338.	1.1	2
43	Restless Leg Syndrome in Peripheral Artery Disease: Prevalence among Patients with Claudication and Benefits from Low-Intensity Exercise. Journal of Clinical Medicine, 2019, 8, 1403.	1.0	2
44	Cortical Oxygenation during a Motor Task to Evaluate Recovery in Subacute Stroke Patients: A Study with Near-Infrared Spectroscopy. Neurology International, 2022, 14, 322-335.	1.3	2
45	A Fitness-Fatigue Model of Performance in Peripheral Artery Disease: Predicted and Measured Effects of a Pain-Free Exercise Program. Journal of Personalized Medicine, 2022, 12, 397.	1.1	2
46	A near-infrared spectroscopy-assisted test discriminates patients with peripheral arterial disease and venous insufficiency with changes of foot oxygenation following light elastic compression therapy. Vasa - European Journal of Vascular Medicine, 2019, 48, 361-367.	0.6	1
47	Beyond the Patient's Report: Self-Reported, Subjective, Objective and Estimated Walking Disability in Patients with Peripheral Artery Disease. Diagnostics, 2021, 11, 1991.	1.3	1
48	Seasonal pattern in elderly hospitalized with acute kidney injury: a retrospective nationwide study in Italy. International Urology and Nephrology, 2022, 54, 3243-3253.	0.6	1
49	Response to Letter to the Editor. Archives of Physical Medicine and Rehabilitation, 2020, 101, 925-926.	0.5	0
50	Epigenetics and doping in sports—The role of microRNAs. , 2021, , 269-284.		0
51	MO870: The Legacy Effect of a Home Walking Exercise Program in Kidney Failure Patients on Dialysis. Nephrology Dialysis Transplantation, 2022, 37, .	0.4	0