
Judy Geissler

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3662184/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Generation and characterization of a control and patient-derived human iPSC line containing the Hermansky Pudlak type 2 (HPS2) associated heterozygous compound mutation in AP3B1. Stem Cell Research, 2021, 54, 102444.	0.7	3
2	Generation and characterization of a human iPSC line SANi007-A from a patient with a heterozygous dominant mutation in ELANE. Stem Cell Research, 2021, 55, 102440.	0.7	1
3	Generation and characterization of a human iPSC line SANi008-A from a Chédiak-Higashi Syndrome patient. Stem Cell Research, 2021, 55, 102442.	0.7	1
4	Generation and characterization of a human iPSC line SANi006-A from a Gray Platelet Syndrome patient. Stem Cell Research, 2021, 55, 102443.	0.7	1
5	MKL1 deficiency results in a severe neutrophil motility defect due to impaired actin polymerization. Blood, 2020, 135, 2171-2181.	1.4	29
6	Genetic variation of human neutrophil Fcl̂³ receptors and SIRPα in antibodyâ€dependent cellular cytotoxicity towards cancer cells. European Journal of Immunology, 2018, 48, 344-354.	2.9	28
7	Hermansky-Pudlak syndrome type 2: Aberrant pre-mRNA splicing and mislocalization of granule proteins in neutrophils. Human Mutation, 2017, 38, 1402-1411.	2.5	21
8	Mutation in an exonic splicing enhancer site causing chronic granulomatous disease. Blood Cells, Molecules, and Diseases, 2017, 66, 50-57.	1.4	13
9	Factor H-Related (FHR)-1 and FHR-2 Form Homo- and Heterodimers, while FHR-5 Circulates Only As Homodimer in Human Plasma. Frontiers in Immunology, 2017, 8, 1328.	4.8	38
10	Complement Factor H-Related Protein 3 Serum Levels Are Low Compared to Factor H and Mainly Determined by Gene Copy Number Variation in CFHR3. PLoS ONE, 2016, 11, e0152164.	2.5	30
11	Primary Immunodeficiency Caused by an Exonized Retroposed Gene Copy Inserted in the <i>CYBB</i> Gene. Human Mutation, 2014, 35, 486-496.	2.5	38
12	Neutrophil responsiveness to IgG, as determined by fixed ratios of mRNA levels for activating and inhibitory Fcl ³ RII (CD32), is stable over time and unaffected by cytokines. Blood, 2006, 108, 584-590.	1.4	35