

# Katsuyuki Miura

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3663077/publications.pdf>

Version: 2024-02-01

272  
papers

7,541  
citations

81743

39  
h-index

74018

75  
g-index

278  
all docs

278  
docs citations

278  
times ranked

10209  
citing authors

#	ARTICLE	IF	CITATIONS
1	The Japanese Society of Hypertension Guidelines for the Management of Hypertension (JSH 2019). Hypertension Research, 2019, 42, 1235-1481.	1.5	1,047
2	Cardiovascular Disease and Risk Factors in Asia. Circulation, 2008, 118, 2702-2709.	1.6	604
3	Dietary Sources of Sodium in China, Japan, the United Kingdom, and the United States, Women and Men Aged 40 to 59 Years: The INTERMAP Study. Journal of the American Dietetic Association, 2010, 110, 736-745.	1.3	440
4	Epidemiology of Hypertension in Japan. Circulation Journal, 2013, 77, 2226-2231.	0.7	155
5	Conversion of Urine Proteinâ€“Creatinine Ratio or Urine Dipstick Protein to Urine Albuminâ€“Creatinine Ratio for Use in Chronic Kidney Disease Screening and Prognosis. Annals of Internal Medicine, 2020, 173, 426-435.	2.0	144
6	Adiposity and risk of decline in glomerular filtration rate: meta-analysis of individual participant data in a global consortium. BMJ: British Medical Journal, 2019, 364, k5301.	2.4	139
7	Blood pressure categories and long-term risk of cardiovascular disease according to age group in Japanese men and women. Hypertension Research, 2012, 35, 947-953.	1.5	134
8	Four Blood Pressure Indexes and the Risk of Stroke and Myocardial Infarction in Japanese Men and Women. Circulation, 2009, 119, 1892-1898.	1.6	118
9	Incidence, Management and Short-Term Outcome of Stroke in a General Population of 1.4 Million Japaneseâ€“Shiga Stroke Registry â€“. Circulation Journal, 2017, 81, 1636-1646.	0.7	118
10	Dietary sodium-to-potassium ratio as a risk factor for stroke, cardiovascular disease and all-cause mortality in Japan: the NIPPON DATA80 cohort study. BMJ Open, 2016, 6, e011632.	0.8	104
11	Association of extremely high levels of high-density lipoprotein cholesterol with cardiovascular mortality in a pooled analysis of 9 cohort studies including 43,407 individuals: The EPOCHâ€“JAPAN study. Journal of Clinical Lipidology, 2018, 12, 674-684.e5.	0.6	101
12	Rationale and Descriptive Analysis of Specific Health Guidance: the Nationwide Lifestyle Intervention Program Targeting Metabolic Syndrome in Japan. Journal of Atherosclerosis and Thrombosis, 2018, 25, 308-322.	0.9	88
13	Relation Between Serum Total Cholesterol Level and Cardiovascular Disease Stratified by Sex and Age Group: A Pooled Analysis of 65,594 Individuals From 10 Cohort Studies in Japan. Journal of the American Heart Association, 2012, 1, e001974.	1.6	84
14	Time to Consider Use of the Sodium-to-Potassium Ratio for Practical Sodium Reduction and Potassium Increase. Nutrients, 2017, 9, 700.	1.7	84
15	Having few remaining teeth is associated with a low nutrient intake and low serum albumin levels in middle-aged and older Japanese individuals: findings from the NIPPON DATA2010. Environmental Health and Preventive Medicine, 2019, 24, 1.	1.4	84
16	Relationship of Dietary Linoleic Acid to Blood Pressure. Hypertension, 2008, 52, 408-414.	1.3	76
17	Relation of Dietary Sodium (Salt) to Blood Pressure and Its Possible Modulation by Other Dietary Factors. Hypertension, 2018, 71, 631-637.	1.3	76
18	Associations of socioeconomic status with prevalence, awareness, treatment, and control of hypertension in a general Japanese population. Journal of Hypertension, 2017, 35, 401-408.	0.3	74

#	ARTICLE	IF	CITATIONS
19	Impact of Metabolic Syndrome on the Risk of Cardiovascular Disease Mortality in the United States and in Japan. <i>American Journal of Cardiology</i> , 2014, 113, 84-89.	0.7	69
20	Salt intake and prevalence of overweight/obesity in Japan, China, the United Kingdom, and the United States: the INTERMAP Study. <i>American Journal of Clinical Nutrition</i> , 2019, 110, 34-40.	2.2	69
21	Population attributable numbers and fractions of deaths due to smoking: A pooled analysis of 180,000 Japanese. <i>Preventive Medicine</i> , 2011, 52, 60-65.	1.6	63
22	Volumetric changes in the aging rat brain and its impact on cognitive and locomotor functions. <i>Experimental Gerontology</i> , 2017, 99, 69-79.	1.2	63
23	Influence of Smoking Combined with Another Risk Factor on the Risk of Mortality from Coronary Heart Disease and Stroke: Pooled Analysis of 10 Japanese Cohort Studies. <i>Cerebrovascular Diseases</i> , 2012, 33, 480-491.	0.8	62
24	HbA1c and the Risks for All-Cause and Cardiovascular Mortality in the General Japanese Population. <i>Diabetes Care</i> , 2013, 36, 3759-3765.	4.3	61
25	Plasma 25-hydroxyvitamin D concentration and subsequent risk of total and site specific cancers in Japanese population: large case-cohort study within Japan Public Health Center-based Prospective Study cohort. <i>BMJ: British Medical Journal</i> , 2018, 360, k671.	2.4	61
26	Low-carbohydrate diets and cardiovascular and total mortality in Japanese: a 29-year follow-up of NIPPON DATA80. <i>British Journal of Nutrition</i> , 2014, 112, 916-924.	1.2	59
27	Statin Use and Risk of Cerebral Aneurysm Rupture: A Hospital-based Case-control Study in Japan. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2014, 23, 343-348.	0.7	58
28	Epidemiology of Cardiovascular Risk Factors in Asian Countries. <i>Circulation Journal</i> , 2013, 77, 2851-2859.	0.7	56
29	Six random specimens of daytime casual urine on different days are sufficient to estimate daily sodium/potassium ratio in comparison to 7-day 24-h urine collections. <i>Hypertension Research</i> , 2014, 37, 765-771.	1.5	56
30	Fiber-rich diet with brown rice improves endothelial function in type 2 diabetes mellitus: A randomized controlled trial. <i>PLoS ONE</i> , 2017, 12, e0179869.	1.1	52
31	Long-chain n-3 polyunsaturated fatty acids intake and cardiovascular disease mortality risk in Japanese: A 24-year follow-up of NIPPON DATA80. <i>Atherosclerosis</i> , 2014, 232, 384-389.	0.4	51
32	Increased Aortic Calcification Is Associated With Arterial Stiffness Progression in Multiethnic Middle-Aged Men. <i>Hypertension</i> , 2017, 69, 102-108.	1.3	51
33	Mutant <i>KCNJ3</i> and <i>KCNJ5</i> Potassium Channels as Novel Molecular Targets in Bradyarrhythmias and Atrial Fibrillation. <i>Circulation</i> , 2019, 139, 2157-2169.	1.6	51
34	Epidemiology of hypertension in Japan: beyond the new 2019 Japanese guidelines. <i>Hypertension Research</i> , 2020, 43, 1344-1351.	1.5	49
35	The Relationship between Very High Levels of Serum High-Density Lipoprotein Cholesterol and Cause-Specific Mortality in a 20-Year Follow-Up Study of Japanese General Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2016, 23, 800-809.	0.9	48
36	Lipoprotein-associated phospholipase A2 is related to risk of subclinical atherosclerosis but is not supported by Mendelian randomization analysis in a general Japanese population. <i>Atherosclerosis</i> , 2016, 246, 141-147.	0.4	48

#	ARTICLE	IF	CITATIONS
37	Dietary Salt Intake and Blood Pressure in a Representative Japanese Population: Baseline Analyses of NIPPON DATA80. <i>Journal of Epidemiology</i> , 2010, 20, S524-S530.	1.1	45
38	Relationship of Insulin Resistance to Prevalence and Progression of Coronary Artery Calcification Beyond Metabolic Syndrome Components. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2016, 36, 1703-1708.	1.1	44
39	Relationship of dietary cholesterol to blood pressure: the INTERMAP study. <i>Journal of Hypertension</i> , 2011, 29, 222-228.	0.3	42
40	Serum n-6 fatty acids and lipoprotein subclasses in middle-aged men: the population-based cross-sectional ERA-JUMP Study. <i>American Journal of Clinical Nutrition</i> , 2010, 91, 1195-1203.	2.2	41
41	Ipragliflozin, a sodium-glucose cotransporter 2 inhibitor, reduces bodyweight and fat mass, but not muscle mass, in Japanese type 2 diabetes patients treated with insulin: A randomized clinical trial. <i>Journal of Diabetes Investigation</i> , 2019, 10, 1012-1021.	1.1	41
42	Integration of Data from NIPPON DATA80/90 and National Nutrition Survey in Japan: For Cohort Studies of Representative Japanese on Nutrition. <i>Journal of Epidemiology</i> , 2010, 20, S506-S514.	1.1	40
43	Secular trends of the impact of overweight and obesity on hypertension in Japan, 1980-2010. <i>Hypertension Research</i> , 2015, 38, 790-795.	1.5	39
44	Smoking, Smoking Cessation, and Measures of Subclinical Atherosclerosis in Multiple Vascular Beds in Japanese Men. <i>Journal of the American Heart Association</i> , 2016, 5, .	1.6	39
45	Seasonal variation in home blood pressure: findings from nationwide web-based monitoring in Japan. <i>BMJ Open</i> , 2018, 8, e017351.	0.8	39
46	Relationship of dietary monounsaturated fatty acids to blood pressure. <i>Journal of Hypertension</i> , 2013, 31, 1144-1150.	0.3	38
47	The association of fish consumption and its urinary metabolites with cardiovascular risk factors: the International Study of Macro-/Micronutrients and Blood Pressure (INTERMAP). <i>American Journal of Clinical Nutrition</i> , 2020, 111, 280-290.	2.2	37
48	Cross-Sectional Comparison of Coronary Artery Calcium Scores Between Caucasian Men in the United States and Japanese Men in Japan: The Multi-Ethnic Study of Atherosclerosis and the Shiga Epidemiological Study of Subclinical Atherosclerosis. <i>American Journal of Epidemiology</i> , 2014, 180, 590-598.	1.6	36
49	Continuous decline in mortality from coronary heart disease in Japan despite a continuous and marked rise in total cholesterol: Japanese experience after the Seven Countries Study. <i>International Journal of Epidemiology</i> , 2015, 44, 1614-1624.	0.9	36
50	Estimating 24-h urinary sodium/potassium ratio from casual (spot) urinary sodium/potassium ratio: the INTERSALT Study. <i>International Journal of Epidemiology</i> , 2017, 46, dyw287.	0.9	34
51	Epidemiology and prevention of hypertension in Japanese: how could Japan get longevity?. <i>EPMA Journal</i> , 2011, 2, 59-64.	3.3	33
52	HOMA-IR and the risk of hyperuricemia: A prospective study in non-diabetic Japanese men. <i>Diabetes Research and Clinical Practice</i> , 2014, 106, 154-160.	1.1	33
53	Smoking increases the risk of all-cause and cardiovascular mortality in patients with chronic kidney disease. <i>Kidney International</i> , 2015, 88, 1144-1152.	2.6	32
54	Significant inverse association of equol-producer status with coronary artery calcification but not dietary isoflavones in healthy Japanese men. <i>British Journal of Nutrition</i> , 2017, 117, 260-266.	1.2	31

#	ARTICLE	IF	CITATIONS
55	Diurnal variation of urinary sodium-to-potassium ratio in free-living Japanese individuals. <i>Hypertension Research</i> , 2017, 40, 658-664.	1.5	31
56	Lifetime cigarette smoking is associated with abdominal obesity in a community-based sample of Japanese men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Preventive Medicine Reports</i> , 2016, 4, 225-232.	0.8	30
57	Effectiveness of a Self-monitoring Device for Urinary Sodium-to-Potassium Ratio on Dietary Improvement in Free-Living Adults: a Randomized Controlled Trial. <i>Journal of Epidemiology</i> , 2018, 28, 41-47.	1.1	30
58	Relationship of serum irisin levels to prevalence and progression of coronary artery calcification: A prospective, population-based study. <i>International Journal of Cardiology</i> , 2018, 267, 177-182.	0.8	30
59	Lifetime Risk of Stroke and Coronary Heart Disease Deaths According to Blood Pressure Level. <i>Hypertension</i> , 2019, 73, 52-59.	1.3	30
60	Relationship between non-high-density lipoprotein cholesterol and the long-term mortality of cardiovascular diseases: NIPPON DATA 90. <i>International Journal of Cardiology</i> , 2016, 220, 262-267.	0.8	29
61	The National Integrated Project for Prospective Observation of Non-communicable Disease and its Trends in the Aged 2010 (NIPPON DATA2010): Objectives, Design, and Population Characteristics. <i>Journal of Epidemiology</i> , 2018, 28, S2-S9.	1.1	29
62	Associations between Rice, Noodle, and Bread Intake and Sleep Quality in Japanese Men and Women. <i>PLoS ONE</i> , 2014, 9, e105198.	1.1	29
63	Age-specific impact of diabetes mellitus on the risk of cardiovascular mortality: An overview from the evidence for Cardiovascular Prevention from Observational Cohorts in the Japan Research Group (EPOCH-JAPAN). <i>Journal of Epidemiology</i> , 2017, 27, 123-129.	1.1	28
64	Mendelian randomization analysis in three Japanese populations supports a causal role of alcohol consumption in lowering low-density lipid cholesterol levels and particle numbers. <i>Atherosclerosis</i> , 2016, 254, 242-248.	0.4	27
65	Serum magnesium, phosphorus, and calcium levels and subclinical calcific aortic valve disease: A population-based study. <i>Atherosclerosis</i> , 2018, 273, 145-152.	0.4	27
66	Urinary sodium-to-potassium ratio and intake of sodium and potassium among men and women from multiethnic general populations: the INTERSALT Study. <i>Hypertension Research</i> , 2019, 42, 1590-1598.	1.5	27
67	Coronary Artery Calcification by Computed Tomography in Epidemiologic Research and Cardiovascular Disease Prevention. <i>Journal of Epidemiology</i> , 2012, 22, 188-198.	1.1	26
68	Proton magnetic resonance spectroscopy assessment of metabolite status of the anterior cingulate cortex in chronic pain patients and healthy controls. <i>Journal of Pain Research</i> , 2017, Volume 10, 287-293.	0.8	26
69	Relationship Between Socioeconomic Status and the Prevalence of Underweight, Overweight or Obesity in a General Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S10-S16.	1.1	26
70	Brachial-ankle pulse wave velocity is associated with coronary calcification among 1131 healthy middle-aged men. <i>International Journal of Cardiology</i> , 2015, 189, 67-72.	0.8	24
71	Absolute risk score for stroke, myocardial infarction, and all cardiovascular disease: Japan Arteriosclerosis Longitudinal Study. <i>Hypertension Research</i> , 2019, 42, 567-579.	1.5	24
72	Comparison of blood pressure values—self-measured at home, measured at an unattended office, and measured at a conventional attended office. <i>Hypertension Research</i> , 2019, 42, 1726-1737.	1.5	23

#	ARTICLE	IF	CITATIONS
73	Lipoprotein particle profiles compared with standard lipids in association with coronary artery calcification in the general Japanese population. <i>Atherosclerosis</i> , 2014, 236, 237-243.	0.4	22
74	Trends in antipsychotic prescriptions for Japanese outpatients during 2006-2012: a descriptive epidemiological study. <i>Pharmacoepidemiology and Drug Safety</i> , 2017, 26, 642-656.	0.9	22
75	Association of blood levels of marine omega-3 fatty acids with coronary calcification and calcium density in Japanese men. <i>European Journal of Clinical Nutrition</i> , 2019, 73, 783-792.	1.3	22
76	A cross-sectional association of obesity with coronary calcium among Japanese, Koreans, Japanese Americans, and US Whites. <i>European Heart Journal Cardiovascular Imaging</i> , 2013, 14, 921-927.	0.5	21
77	[Scientific Statement]. <i>Hypertension Research</i> , 2013, 36, 1020-1025.	1.5	21
78	Inclusion of a Genetic Risk Score into a Validated Risk Prediction Model for Colorectal Cancer in Japanese Men Improves Performance. <i>Cancer Prevention Research</i> , 2017, 10, 535-541.	0.7	21
79	Re-evaluation of the associations of egg intake with serum total cholesterol and cause-specific and total mortality in Japanese women. <i>European Journal of Clinical Nutrition</i> , 2018, 72, 841-847.	1.3	21
80	Twelve-year trends of increasing overweight and obesity in patients with diabetes: the Shiga Diabetes Clinical Survey. <i>Endocrine Journal</i> , 2018, 65, 527-536.	0.7	21
81	Association of blood pressure with estimates of 24-h urinary sodium and potassium excretion from repeated single-spot urine samples. <i>Hypertension Research</i> , 2019, 42, 411-418.	1.5	21
82	Relationship between 5-Year Decline in Instrumental Activity of Daily Living and Accumulation of Cardiovascular Risk Factors: NIPPON DATA90. <i>Journal of Atherosclerosis and Thrombosis</i> , 2010, 17, 64-72.	0.9	20
83	Food sources of dietary sodium in the Japanese adult population: the international study of macro-/micronutrients and blood pressure (INTERMAP). <i>European Journal of Nutrition</i> , 2017, 56, 1269-1280.	1.8	20
84	A pooled analysis of the association of isolated low levels of high-density lipoprotein cholesterol with cardiovascular mortality in Japan. <i>European Journal of Epidemiology</i> , 2017, 32, 547-557.	2.5	20
85	Urinary Sodium-to-Potassium Ratio Tracks the Changes in Salt Intake during an Experimental Feeding Study Using Standardized Low-Salt and High-Salt Meals among Healthy Japanese Volunteers. <i>Nutrients</i> , 2017, 9, 951.	1.7	20
86	Brain Metabolite Changes in the Anterior Cingulate Cortex of Chronic Low Back Pain Patients and Correlations Between Metabolites and Psychological State. <i>Clinical Journal of Pain</i> , 2018, 34, 657-663.	0.8	20
87	The Japanese Society of Hypertensionâ€™Digest of plan for the future. <i>Hypertension Research</i> , 2018, 41, 989-990.	1.5	20
88	The impact of equol-producing status in modifying the effect of soya isoflavones on risk factors for CHD: a systematic review of randomised controlled trials. <i>Journal of Nutritional Science</i> , 2016, 5, e30.	0.7	19
89	Relation of unprocessed, processed red meat and poultry consumption to blood pressure in East Asian and Western adults. <i>Journal of Hypertension</i> , 2016, 34, 1721-1729.	0.3	19
90	Macronutrient Intake and Socioeconomic Status: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S17-S22.	1.1	19

#	ARTICLE	IF	CITATIONS
91	Relationship Between Step Counts and Cerebral Small Vessel Disease in Japanese Men. <i>Stroke</i> , 2020, 51, 3584-3591.	1.0	19
92	High-density lipoprotein particle concentration and subclinical atherosclerosis of the carotid arteries in Japanese men. <i>Atherosclerosis</i> , 2015, 239, 444-450.	0.4	18
93	Associations of High-Density Lipoprotein Particle and High-Density Lipoprotein Cholesterol With Alcohol Intake, Smoking, and Body Mass Index—The INTERLIPID Study. <i>Circulation Journal</i> , 2018, 82, 2557-2565.	0.7	18
94	Intracranial Artery Stenosis and Its Association With Conventional Risk Factors in a General Population of Japanese Men. <i>Stroke</i> , 2019, 50, 2967-2969.	1.0	18
95	Dietary Inflammatory Index Positively Associated With High-Sensitivity C-Reactive Protein Level in Japanese From NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2020, 30, 98-107.	1.1	18
96	Cumulative impact of axial, structural, and repolarization ECG findings on long-term cardiovascular mortality among healthy individuals in Japan: National Integrated Project for Prospective Observation of Non-Communicable Disease and its Trends in the Aged, 1980 and 1990. <i>European Journal of Preventive Cardiology</i> , 2014, 21, 1501-1508.	0.8	17
97	The Association between Glomerular Filtration Rate Estimated on Admission and Acute Stroke Outcome: The Shiga Stroke Registry. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 570-579.	0.9	17
98	Associations between Socioeconomic Status and the Prevalence and Treatment of Hypercholesterolemia in a General Japanese Population: NIPPON DATA2010. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 606-620.	0.9	17
99	Socioeconomic Status and Knowledge of Cardiovascular Risk Factors: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S46-S52.	1.1	17
100	Agreement between 24-h dietary recalls and 24-h urine collections for estimating sodium intake in China, Japan, UK, USA. <i>Journal of Hypertension</i> , 2019, 37, 814-819.	0.3	17
101	Vegetable Protein Intake was Inversely Associated with Cardiovascular Mortality in a 15-Year Follow-Up Study of the General Japanese Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 198-206.	0.9	17
102	Vitamin B6 intake and incidence of diabetic retinopathy in Japanese patients with type 2 diabetes: analysis of data from the Japan Diabetes Complications Study (JDCS). <i>European Journal of Nutrition</i> , 2020, 59, 1585-1594.	1.8	17
103	Relationship between carbohydrate and dietary fibre intake and the risk of cardiovascular disease mortality in Japanese: 24-year follow-up of NIPPON DATA80. <i>European Journal of Clinical Nutrition</i> , 2020, 74, 67-76.	1.3	17
104	Cross-sectional association between exposure to particulate matter and inflammatory markers in the Japanese general population: NIPPON DATA2010. <i>Environmental Pollution</i> , 2016, 213, 460-467.	3.7	16
105	Two-Year Survival After First-Ever Stroke in a General Population of 1.4 Million Japanese—Shiga Stroke Registry. <i>Circulation Journal</i> , 2018, 82, 2549-2556.	0.7	16
106	Association Between Body Mass Index and All-Cause Death in Japanese Population: Pooled Individual Participant Data Analysis of 13 Cohort Studies. <i>Journal of Epidemiology</i> , 2019, 29, 457-463.	1.1	16
107	Long-Term Survival after Stroke in 1.4 Million Japanese Population: Shiga Stroke and Heart Attack Registry. <i>Journal of Stroke</i> , 2020, 22, 336-344.	1.4	16
108	Fasting but not casual blood glucose is associated with pancreatic cancer mortality in Japanese: EPOCH-JAPAN. <i>Cancer Causes and Control</i> , 2017, 28, 625-633.	0.8	15

#	ARTICLE	IF	CITATIONS
109	Variation of Risk Factors for Cause-Specific Reintubation: A Preliminary Study. <i>Canadian Respiratory Journal</i> , 2018, 2018, 1-6.	0.8	15
110	Socioeconomic Status Associated With Urinary Sodium and Potassium Excretion in Japan: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S29-S34.	1.1	15
111	Meat intake and incidence of cardiovascular disease in Japanese patients with type 2 diabetes: analysis of the Japan Diabetes Complications Study (JDCS). <i>European Journal of Nutrition</i> , 2019, 58, 281-290.	1.8	15
112	Metabolic changes induced by dapagliflozin, an SGLT2 inhibitor, in Japanese patients with type 2 diabetes treated by oral anti-diabetic agents: A randomized, clinical trial. <i>Diabetes Research and Clinical Practice</i> , 2022, 186, 109781.	1.1	15
113	Association between antidepressant use during pregnancy and autism spectrum disorder in children: a retrospective cohort study based on Japanese claims data. <i>Maternal Health, Neonatology and Perinatology</i> , 2019, 5, 1.	1.0	14
114	Physical activity levels in American and Japanese men from the ERA-JUMP Study and associations with metabolic syndrome. <i>Journal of Sport and Health Science</i> , 2020, 9, 170-178.	3.3	14
115	Isolated systolic hypertension and 29-year cardiovascular mortality risk in Japanese adults aged 30–49 years. <i>Journal of Hypertension</i> , 2020, 38, 2230-2236.	0.3	14
116	Is the Proportion of Carbohydrate Intake Associated with the Incidence of Diabetes Complications? An Analysis of the Japan Diabetes Complications Study. <i>Nutrients</i> , 2017, 9, 113.	1.7	13
117	Development of a High-Sensitivity Method for the Measurement of Human Nasal A $\beta$ 42, Tau, and Phosphorylated Tau. <i>Journal of Alzheimer's Disease</i> , 2018, 62, 737-744.	1.2	13
118	Differences Between Coronary Artery Calcification and Aortic Artery Calcification in Relation to Cardiovascular Disease Risk Factors in Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 452-464.	0.9	13
119	Association between plant-based diets and blood pressure in the INTERMAP study. <i>BMJ Nutrition, Prevention and Health</i> , 2020, 3, 133-142.	1.9	13
120	Food Sources of Dietary Potassium in the Adult Japanese Population: The International Study of Macro-/Micronutrients and Blood Pressure (INTERMAP). <i>Nutrients</i> , 2020, 12, 787.	1.7	13
121	A Combination of Blood Pressure and Total Cholesterol Increases the Lifetime Risk of Coronary Heart Disease Mortality: EPOCH-JAPAN. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021, 28, 6-24.	0.9	13
122	Long-term outcomes associated with prolonged PR interval in the general Japanese population. <i>International Journal of Cardiology</i> , 2015, 184, 291-293.	0.8	12
123	Associations of serum LDL particle concentration with carotid intima-media thickness and coronary artery calcification. <i>Journal of Clinical Lipidology</i> , 2016, 10, 1195-1202.e1.	0.6	12
124	Relationship of three different types of low-carbohydrate diet to cardiometabolic risk factors in a Japanese population: the INTERMAP/INTERLIPID Study. <i>European Journal of Nutrition</i> , 2016, 55, 1515-1524.	1.8	12
125	Effects of $\beta$ -estradiol on cold-sensitive receptor channel TRPM8 in ovariectomized rats. <i>Experimental Animals</i> , 2017, 66, 337-343.	0.7	12
126	Relationships among Socioeconomic Factors and Self-rated Health in Japanese Adults: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S66-S72.	1.1	12



#	ARTICLE	IF	CITATIONS
127	The relationship between serum levels of LOX-1 ligand containing ApoAI as a novel marker of dysfunctional HDL and coronary artery calcification in middle-aged Japanese men. <i>Atherosclerosis</i> , 2020, 313, 20-25.	0.4	12
128	Association between antidepressant use during pregnancy and congenital anomalies in children: A retrospective cohort study based on Japanese claims data. <i>Congenital Anomalies (discontinued)</i> , 2020, 60, 180-188.	0.3	12
129	The Association Between Coronary Artery Calcification and Subclinical Cerebrovascular Diseases in Men: An Observational Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020, 27, 995-1009.	0.9	12
130	High long-chain n-3 fatty acid intake attenuates the effect of high resting heart rate on cardiovascular mortality risk: A 24-year follow-up of Japanese general population. <i>Journal of Cardiology</i> , 2014, 64, 218-224.	0.8	11
131	Serum level of LOX-1 ligand containing ApoB is associated with increased carotid intima-media thickness in Japanese community-dwelling men, especially those with hypercholesterolemia. <i>Journal of Clinical Lipidology</i> , 2016, 10, 172-180.e1.	0.6	11
132	Socioeconomic Inequalities in Oral Health among Middle-Aged and Elderly Japanese: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S59-S65.	1.1	11
133	Change in Pericardial Fat Volume and Cardiovascular Risk Factors in a General Population of Japanese Men. <i>Circulation Journal</i> , 2018, 82, 2542-2548.	0.7	11
134	Cardiovascular Risk Assessment Chart by Dietary Factors in Japan. <i>NIPPON DATA80</i> . <i>Circulation Journal</i> , 2019, 83, 1254-1260.	0.7	11
135	Socioeconomic and lifestyle factors associated with depressive tendencies in general Japanese men and women: NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , 2019, 24, 37.	1.4	11
136	Quantitative CT analysis of honeycombing area predicts mortality in idiopathic pulmonary fibrosis with definite usual interstitial pneumonia pattern: A retrospective cohort study. <i>PLoS ONE</i> , 2019, 14, e0214278.	1.1	11
137	Relationship of household salt intake level with long-term all-cause and cardiovascular disease mortality in Japan: NIPPON DATA80. <i>Hypertension Research</i> , 2020, 43, 132-139.	1.5	11
138	Association of Total Marine Fatty Acids, Eicosapentaenoic and Docosahexaenoic Acids, With Aortic Stiffness in Koreans, Whites, and Japanese Americans. <i>American Journal of Hypertension</i> , 2013, 26, 1321-1327.	1.0	10
139	Dietary tofu intake and long-term risk of death from stroke in a general population. <i>Clinical Nutrition</i> , 2018, 37, 182-188.	2.3	10
140	Relationships among Food Group Intakes, Household Expenditure, and Education Attainment in a General Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S23-S28.	1.1	10
141	Reduced Lung Function and Cerebral Small Vessel Disease in Japanese Men: the Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 1009-1021.	0.9	10
142	Coronary Artery Calcium Progression Among the US and Japanese Men. <i>Circulation: Cardiovascular Imaging</i> , 2019, 12, e008104.	1.3	10
143	Proteinuria and Reduced Estimated Glomerular Filtration Rate are Independently Associated With Lower Cognitive Abilities in Apparently Healthy Community-Dwelling Elderly Men in Japan: A Cross-sectional Study. <i>Journal of Epidemiology</i> , 2020, 30, 244-252.	1.1	10
144	JSH Statement: Tokyo declaration promoting salt reduction by the Japanese Society of Hypertension. <i>Hypertension Research</i> , 2020, 43, 1133-1134.	1.5	10

#	ARTICLE	IF	CITATIONS
145	Sorafenib exposure and its correlation with response and safety in advanced hepatocellular carcinoma: results from an observational retrospective study. <i>Cancer Chemotherapy and Pharmacology</i> , 2020, 86, 129-139.	1.1	10
146	Estimation of 10-Year Risk of Death from Coronary Heart Disease, Stroke, and Cardiovascular Disease in a Pooled Analysis of Japanese Cohorts: EPOCH-JAPAN. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021, 28, 816-825.	0.9	10
147	Dietary Intake of Potassium and Associated Dietary Factors among Representative Samples of Japanese General Population: NIPPON DATA 80/90. <i>Journal of Epidemiology</i> , 2010, 20, S567-S575.	1.1	9
148	Relation of Serum Leptin and Adiponectin Level to Serum C-Reactive Protein: The INTERLIPID Study. <i>International Journal of Vascular Medicine</i> , 2013, 2013, 1-7.	0.4	9
149	Holter monitoring for the screening of cardiac disease in diabetes mellitus: The non-invasive Holter monitoring observation of new cardiac events in diabetics study. <i>Diabetes and Vascular Disease Research</i> , 2015, 12, 396-404.	0.9	9
150	Relationship of type of work with health-related quality of life. <i>Quality of Life Research</i> , 2015, 24, 2927-2932.	1.5	9
151	Maximum BMI and microvascular complications in a cohort of Japanese patients with type 2 diabetes: the Japan Diabetes Complications Study. <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 790-797.	1.2	9
152	Validation of the european SCORE risk chart in the healthy middle-aged Japanese. <i>Atherosclerosis</i> , 2016, 252, 116-121.	0.4	9
153	International Comparison of Abdominal Fat Distribution Among Four Populations: The ERA-JUMP Study. <i>Metabolic Syndrome and Related Disorders</i> , 2018, 16, 166-173.	0.5	9
154	Comparison of carotid plaque burden among healthy middle-aged men living in the US, Japan, and South Korea. <i>International Journal of Cardiology</i> , 2018, 266, 245-249.	0.8	9
155	Associations of Overweight, Obesity, and Underweight With High Serum Total Cholesterol Level Over 30 Years Among the Japanese Elderly: NIPPON DATA 80, 90, and 2010. <i>Journal of Epidemiology</i> , 2019, 29, 133-138.	1.1	9
156	Elevated Fasting Blood Glucose Levels Are Associated With Lower Cognitive Function, With a Threshold in Non-Diabetic Individuals: A Population-Based Study. <i>Journal of Epidemiology</i> , 2020, 30, 121-127.	1.1	9
157	The association between subjective health perception and lifestyle factors in Shiga prefecture, Japan: a cross-sectional study. <i>BMC Public Health</i> , 2020, 20, 1786.	1.2	9
158	Relationship Between Non-fasting Triglycerides and Cardiovascular Disease Mortality in a 20-year Follow-up Study of a Japanese General Population: NIPPON DATA90. <i>Journal of Epidemiology</i> , 2022, 32, 303-313.	1.1	9
159	Association of Alcohol Consumption With Fat Deposition in a Community-Based Sample of Japanese Men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Epidemiology</i> , 2019, 29, 205-212.	1.1	9
160	Relationship between Kidney Function and Subclinical Atherosclerosis Progression Evaluated by Coronary Artery Calcification. <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 1359-1371.	0.9	9
161	Does the flushing response modify the relationship between alcohol intake and hypertension in the Japanese population? NIPPON DATA2010. <i>Hypertension Research</i> , 2016, 39, 670-679.	1.5	8
162	Impacts of chronic kidney disease and diabetes on cardiovascular mortality in a general Japanese population: A 20-year follow-up of the NIPPON DATA90 study. <i>European Journal of Preventive Cardiology</i> , 2017, 24, 505-513.	0.8	8

#	ARTICLE	IF	CITATIONS
163	Overweight or underweight and the risk of decline in activities of daily living in a 22-year cohort study of a Japanese sample. <i>Geriatrics and Gerontology International</i> , 2018, 18, 799-805.	0.7	8
164	Factors Related to Participation in Health Examinations for Japanese National Health Insurance: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S53-S58.	1.1	8
165	The impact of sex on risk of cardiovascular disease and all-cause mortality in adults with or without diabetes mellitus: A comparison between the U.S. and Japan. <i>Journal of Diabetes and Its Complications</i> , 2019, 33, 417-423.	1.2	8
166	Waist Circumference and Domain-Specific Cognitive Function Among Non-Demented Japanese Older Adults Stratified by Sex: Results from the Takashima Cognition Study. <i>Journal of Alzheimer's Disease</i> , 2020, 73, 887-896.	1.2	8
167	Inclusion of a gene-environment interaction between alcohol consumption and the aldehyde dehydrogenase 2 genotype in a risk prediction model for upper aerodigestive tract cancer in Japanese men. <i>Cancer Science</i> , 2020, 111, 3835-3844.	1.7	8
168	A genome-wide association study in Japanese identified one variant associated with a preference for a Japanese dietary pattern. <i>European Journal of Clinical Nutrition</i> , 2021, 75, 937-945.	1.3	8
169	Seasonal Variation in Incidence of Stroke in a General Population of 1.4 Million Japanese: The Shiga Stroke Registry. <i>Cerebrovascular Diseases</i> , 2022, 51, 75-81.	0.8	8
170	Comparability in coronary artery calcium scores on CT scan between two community-based cohort studies. <i>International Journal of Cardiology</i> , 2011, 149, 244-245.	0.8	7
171	Interaction between dietary marine-derived n-3 fatty acids intake and J-point elevation on the risk of cardiac death: a 24-year follow-up of Japanese men. <i>Heart</i> , 2013, 99, 1024-1029.	1.2	7
172	Exclusion of emphysematous lung from dose-volume estimates of risk improves prediction of radiation pneumonitis. <i>Radiation Oncology</i> , 2017, 12, 160.	1.2	7
173	Association of Work Situation With Cardiovascular Disease Mortality Risk Among Working-Age Japanese Men—A 20-Year Follow-up of NIPPON DATA90. <i>Circulation Journal</i> , 2019, 83, 1506-1513.	0.7	7
174	One-year weight loss maintenance outcomes following a worksite-based weight reduction program among Japanese men with cardiovascular risk factors. <i>Journal of Occupational Health</i> , 2019, 61, 189-196.	1.0	7
175	The association of home and accurately measured office blood pressure with coronary artery calcification among general Japanese men. <i>Journal of Hypertension</i> , 2019, 37, 1676-1681.	0.3	7
176	Factors associated with intra-individual visit-to-visit variability of blood pressure in four countries: the INTERMAP study. <i>Journal of Human Hypertension</i> , 2019, 33, 229-236.	1.0	7
177	The relationship between repeated measurement of casual and 24-h urinary sodium-to-potassium ratio in patients with chronic kidney disease. <i>Journal of Human Hypertension</i> , 2019, 33, 286-297.	1.0	7
178	Health status of workers approximately 60 years of age and the risk of early death after compulsory retirement: A cohort study. <i>Journal of Occupational Health</i> , 2020, 62, e12088.	1.0	7
179	Alcohol consumption and cognitive function in elderly Japanese men. <i>Alcohol</i> , 2020, 85, 145-152.	0.8	7
180	Smoking habits and progression of coronary and aortic artery calcification: A 5-year follow-up of community-dwelling Japanese men. <i>International Journal of Cardiology</i> , 2020, 314, 89-94.	0.8	7

#	ARTICLE	IF	CITATIONS
181	Two-Year Recurrence After First-Ever Stroke in a General Population of 1.4 Million Japanese Patientsâ€”The Shiga Stroke and Heart Attack Registry Study â€”. <i>Circulation Journal</i> , 2020, 84, 943-948.	0.7	7
182	Impact of hypertension stratified by diabetes on the lifetime risk of cardiovascular disease mortality in Japan: a pooled analysis of data from the Evidence for Cardiovascular Prevention from Observational Cohorts in Japan study. <i>Hypertension Research</i> , 2020, 43, 1437-1444.	1.5	7
183	Relationship of Higher-level Functional Capacity With Long-term Mortality in Japanese Older People: NIPPON DATA90. <i>Journal of Epidemiology</i> , 2023, 33, 136-141.	1.1	7
184	Sodium Intake and Incidence of Diabetes Complications in Elderly Patients with Type 2 Diabetesâ€”Analysis of Data from the Japanese Elderly Diabetes Intervention Study (J-EDIT). <i>Nutrients</i> , 2021, 13, 689.	1.7	7
185	A Comparison of Segment-Specific and Composite Measures of Carotid Intima-Media Thickness and their Relationships with Coronary Calcium. <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 282-295.	0.9	7
186	Trends in medical performance in diabetic patients in primary care clinics compared with those in hospitals: Shiga Diabetes Clinical Survey, Japan, 2000â€”2012. <i>Diabetology International</i> , 2017, 8, 59-68.	0.7	6
187	Cardiovascular risk and blood pressure lowering treatment among elderly individuals. <i>Journal of Hypertension</i> , 2018, 36, 410-418.	0.3	6
188	Association of alcohol consumption and aortic calcification in healthy men aged 40â€”49 years for the ERA JUMP Study. <i>Atherosclerosis</i> , 2018, 268, 84-91.	0.4	6
189	JSH Statement: Kyoto declaration on hypertension research in Asia. <i>Hypertension Research</i> , 2019, 42, 759-760.	1.5	6
190	Estimating the costâ€”effectiveness of screening for hepatitis C virus infection in Japan. <i>Hepatology Research</i> , 2020, 50, 542-556.	1.8	6
191	Differences between home blood pressure and strictly measured office blood pressure and their determinants in Japanese men. <i>Hypertension Research</i> , 2021, 44, 80-87.	1.5	6
192	A genome-wide association study on confection consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort Study. <i>British Journal of Nutrition</i> , 2021, 126, 1843-1851.	1.2	6
193	Epidemiology and control of hypertension in Japan: a comparison with Western countries. <i>Journal of Human Hypertension</i> , 2021, , .	1.0	6
194	The 21-Year Trend of Stroke Incidence in a General Japanese Population: Results from the Takashima Stroke Registry, 1990â€”2010. <i>Cerebrovascular Diseases</i> , 2022, 51, 570-576.	0.8	6
195	Serum Ferritin, Insulin Resistance, and Î²-cell Dysfunction: A Prospective Study in Normoglycemic Japanese Men. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2017, 125, 12-20.	0.6	5
196	Self-reported Sleep Duration and Subclinical Atherosclerosis in a General Population of Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 186-198.	0.9	5
197	Plasma C-peptide and glycated albumin and subsequent risk of cancer: From a large prospective caseâ€”cohort study in Japan. <i>International Journal of Cancer</i> , 2019, 144, 718-729.	2.3	5
198	Cluster-randomized controlled trial for the early promotion of clinic visits for untreated hypertension. <i>Hypertension Research</i> , 2021, 44, 355-362.	1.5	5

#	ARTICLE	IF	CITATIONS
199	A genome-wide association study on fish consumption in a Japanese population—the Japan Multi-Institutional Collaborative Cohort study. <i>European Journal of Clinical Nutrition</i> , 2021, 75, 480-488.	1.3	5
200	Risk Factors That Most Accurately Predict Coronary Artery Disease Based on the Duration of Follow-up—NIPPON DATA80. <i>Circulation Journal</i> , 2021, 85, 908-913.	0.7	5
201	Association between socioeconomic status and physical inactivity in a general Japanese population: NIPPON DATA2010. <i>PLoS ONE</i> , 2021, 16, e0254706.	1.1	5
202	Exercise Habits Are Associated with Improved Long-Term Mortality Risks in the Nationwide General Japanese Population: A 20-Year Follow-Up of the NIPPON DATA90 Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020, 252, 253-262.	0.5	5
203	Perioperative Adiponectin Measurement is Useful for Prediction of Postoperative Infection in Patients with Colorectal Cancer. <i>Annals of Surgical Oncology</i> , 2016, 23, 540-545.	0.7	4
204	Association of Coronary Artery Calcification with Estimated Coronary Heart Disease Risk from Prediction Models in a Community-Based Sample of Japanese Men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 477-489.	0.9	4
205	Passive Smoking at Home by Socioeconomic Factors in a Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S40-S45.	1.1	4
206	Time-Related Changes in Relationships Between the Keys Score, Dietary Lipids, and Serum Total Cholesterol in Japan—NIPPON DATA80/90/2010. <i>Circulation Journal</i> , 2018, 83, 147-155.	0.7	4
207	The Relationship of Dietary Cholesterol with Serum Low-Density Lipoprotein Cholesterol and Confounding by Reverse Causality: The INTERLIPID Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 170-182.	0.9	4
208	Liver fat accumulation assessed by computed tomography is an independent risk factor for diabetes mellitus in a population-based study: SESSA (Shiga Epidemiological Study of Subclinical) Tj ETQq0 0 0 rgBT /Overload 110 Tf 50 377 Td (A	1.1	4
209	Recent status of self-measured home blood pressure in the Japanese general population: a modern database on self-measured home blood pressure (MDAS). <i>Hypertension Research</i> , 2020, 43, 1403-1412.	1.5	4
210	Association of Red Meat Intake with the Risk of Cardiovascular Mortality in General Japanese Stratified by Kidney Function: NIPPON DATA80. <i>Nutrients</i> , 2020, 12, 3707.	1.7	4
211	Coronary Artery Calcium Assessed Years Before Was Positively Associated With Subtle White Matter Injury of the Brain in Asymptomatic Middle-Aged Men: The Framingham Heart Study. <i>Circulation: Cardiovascular Imaging</i> , 2021, 14, e011753.	1.3	4
212	Developing a health economic model for Asians with type 2 diabetes based on the Japan Diabetes Complications Study and the Japanese Elderly Diabetes Intervention Trial. <i>BMJ Open Diabetes Research and Care</i> , 2021, 9, e002177.	1.2	4
213	Independent Prognostic Value of Single and Multiple Non-Specific 12-Lead Electrocardiographic Findings for Long-Term Cardiovascular Outcomes: A Prospective Cohort Study. <i>PLoS ONE</i> , 2016, 11, e0157563.	1.1	4
214	School-based routine screenings of electrocardiograms for the diagnosis of long QT syndrome. <i>Europace</i> , 2022, 24, 1496-1503.	0.7	4
215	Eating Slowly Is Associated with Undernutrition among Community-Dwelling Adult Men and Older Adult Women. <i>Nutrients</i> , 2022, 14, 54.	1.7	4
216	Overall sleep status and high sensitivity C-reactive protein: a prospective study in Japanese factory workers. <i>Journal of Sleep Research</i> , 2014, 23, 717-727.	1.7	3

#	ARTICLE	IF	CITATIONS
217	Stroke registries in the world: a systematic review. <i>Nosotchu</i> , 2018, 40, 331-342.	0.0	3
218	Serum long-chain n-3 polyunsaturated fatty acids and aortic calcification in middle-aged men: The population-based cross-sectional ERA-JUMP study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2019, 29, 837-846.	1.1	3
219	Anthropometric Obesity Indices were Stronger than CT-Based Indices in Associations with Carotid Intima-Media Thickness in Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 1102-1114.	0.9	3
220	Cross-sectional association of bone mineral density with coronary artery calcification in an international multi-ethnic population-based cohort of men aged 40â€“49: ERA JUMP study. <i>IJC Heart and Vasculature</i> , 2020, 30, 100618.	0.6	3
221	Relationship Between Calcium Intake and Impaired Activities of Daily Living in a Japanese Population: NIPPON DATA90. <i>Journal of Epidemiology</i> , 2021, 31, 119-124.	1.1	3
222	Association between socioeconomic status and prolonged television viewing time in a general Japanese population: NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , 2021, 26, 57.	1.4	3
223	Association between Stress-Coping Strategy and Functional Disability in the General Older Adult Population: The Takashima Study. <i>Gerontology</i> , 2022, 68, 699-706.	1.4	3
224	A genome-wide association study on meat consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort study. <i>Journal of Nutritional Science</i> , 2021, 10, e61.	0.7	3
225	Overall nutrient and total fat intake among Japanese people: The INTERLIPID Study Japan. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2017, 26, 837-848.	0.3	3
226	Ventricular Premature Complexes and Their Associated Factors in a General Population of Japanese Men. <i>American Journal of Cardiology</i> , 2022, 169, 51-56.	0.7	3
227	Association of equol producing status with aortic calcification in middle-aged Japanese men: The ERA JUMP study. <i>International Journal of Cardiology</i> , 2022, 352, 158-164.	0.8	3
228	Trends in Prevalence, Treatment, and Control of Hypertension According to 40-Year-Old Life Expectancy at Prefectures in Japan from the National Health and Nutrition Surveys. <i>Nutrients</i> , 2022, 14, 1219.	1.7	3
229	NIPPON DATA80/90 Nutrition Study: Appendix Tables. <i>Journal of Epidemiology</i> , 2010, 20, S587-S596.	1.1	2
230	Analysis of factors associated with maintenance discontinuation in implant patients. SpringerPlus, 2015, 4, 767.	1.2	2
231	Progression of coronary artery calcium in Japanese American men and white men in the ERA JUMP study. <i>International Journal of Cardiology</i> , 2017, 228, 672-676.	0.8	2
232	Data on alcohol consumption and coronary artery calcification among asymptomatic middle-aged men for the ERA-JUMP study. <i>Data in Brief</i> , 2018, 17, 1091-1098.	0.5	2
233	The Influence of the Japanese Nationwide Cardiovascular Prevention System Health Guidance on Smoking Cessation Among Smokers: A Propensity Score Matching Analysis. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 323-334.	0.9	2
234	Differences in Lifestyle Improvements With the Intention to Prevent Cardiovascular Diseases by Socioeconomic Status in a Representative Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S35-S39.	1.1	2

#	ARTICLE	IF	CITATIONS
235	The JAGUAR Score Predicts 1-Month Disability/Death in Ischemic Stroke Patient Ineligible for Recanalization Therapy. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2018, 27, 2579-2586.	0.7	2
236	JSH Statement: Asahikawa declaration in promotion of diversity by the Japanese society of hypertension—the JSH Asahikawa declaration. <i>Hypertension Research</i> , 2019, 42, 1483-1484.	1.5	2
237	Electrocardiographic Left Atrial Abnormality and B-Type Natriuretic Peptide in a General Japanese Population: NIPPON DATA2010. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021, 28, 34-43.	0.9	2
238	Relationship of Four Blood Pressure Indexes to Subclinical Cerebrovascular Diseases Assessed by Brain MRI in General Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 174-187.	0.9	2
239	Lipoprotein Particle Profiles Compared With Standard Lipids in the Association With Subclinical Aortic Valve Calcification in Apparently Healthy Japanese Men. <i>Circulation Journal</i> , 2021, 85, 1076-1082.	0.7	2
240	Relationship of Ambient Temperature Parameters to Stroke Incidence in a Japanese Population—the Takashima Stroke Registry, Japan, 1988–2010. <i>Circulation Journal</i> , 2021, 85, 2215-2221.	0.7	2
241	Relationships of Alcohol Consumption with Coronary Risk Factors and Macro- and Micro-Nutrient Intake in Japanese People: The INTERLIPID Study. <i>Journal of Nutritional Science and Vitaminology</i> , 2021, 67, 28-38.	0.2	2
242	The association of reproductive history with hypertension and obesity according to menopausal status: the J-MICC Study. <i>Hypertension Research</i> , 2022, 45, 708-714.	1.5	2
243	Predictors of lower limb fractures in general Japanese: NIPPON DATA90. <i>PLoS ONE</i> , 2022, 17, e0261716.	1.1	2
244	Premature Atrial Contractions and Their Determinants in a General Population of Japanese Men. <i>Circulation Journal</i> , 2022, 86, 1298-1306.	0.7	2
245	Diet, Nutrients, and the Prevention of Hypertension. <i>Current Nutrition Reports</i> , 2012, 1, 87-92.	2.1	1
246	Alcohol Consumption, Hospitalization and Medical Expenditure: A Large Epidemiological Study on the Medical Insurance System in Japan. <i>Alcohol and Alcoholism</i> , 2015, 50, 236-243.	0.9	1
247	Dexmedetomidine attenuates the positive chronotropic effects of intravenous atropine in patients with bradycardia during spinal anaesthesia: a retrospective study. <i>JA Clinical Reports</i> , 2018, 4, 70.	0.2	1
248	Carotid Intima-Media Thickness and Plaque in Apparently Healthy Japanese Individuals with an Estimated 10-Year Absolute Risk of CAD Death According to the Japan Atherosclerosis Society (JAS) Guidelines 2012: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 746-746.	0.9	1
249	Lipoprotein particles and coronary artery calcium in middle-aged US-White and Japanese men. <i>Open Heart</i> , 2019, 6, e001119.	0.9	1
250	Factors Associated with Lower Cognitive Performance Scores Among Older Japanese Men in Hawaii and Japan. <i>Journal of Alzheimer's Disease</i> , 2021, 81, 403-412.	1.2	1
251	Prologue: Special Spotlight Issue on Japan. <i>Journal of Human Hypertension</i> , 2021, , .	1.0	1
252	A genome-wide association study on adherence to low-carbohydrate diets in Japanese. <i>European Journal of Clinical Nutrition</i> , 2022, , .	1.3	1

#	ARTICLE	IF	CITATIONS
253	Association between C-Reactive Protein Levels and Functional Disability in the General Older-Population: The Takashima Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2023, 30, 56-65.	0.9	1
254	Eighteen-year trends in the management of patients with diabetes in the Shiga Diabetes Clinical Survey: overall trends and differences by age group. <i>Diabetology International</i> , 2022, 13, 566-574.	0.7	1
255	Scientific evidences for the cut-off points of blood pressure. <i>Health Evaluation and Promotion</i> , 2015, 42, 280-286.	0.0	0
256	ICâ€œPaâ€œ12: OBJECTIVELY MEASURED PHYSICAL ACTIVITY AND BRAIN VOLUME IN JAPANESE ADULT MEN: FINDINGS FROM THE SHIGA EPIDEMIOLOGICAL STUDY OF SUBCLINICAL ATHEROSCLEROSIS. <i>Alzheimer's and Dementia</i> , 2019, 15, P96.	0.4	0
257	Dynamic changes of mitral annulus in patients with degenerative mitral regurgitation and chronic atrial fibrillation undergoing mitral valve reconstruction. <i>General Thoracic and Cardiovascular Surgery</i> , 2020, 68, 1405-1411.	0.4	0
258	Abstract P096: Association Of Accurately Measured Office, Self-measured Home, And Ambulatory Blood Pressure And Their Variability With Intracranial Arterial Stenosis. <i>Circulation</i> , 2021, 143, .	1.6	0
259	Association between adherence to warfarin and thrombotic events in patients with antiphospholipid syndrome in Japan: A claimsâ€œbased retrospective cohort study. <i>Pharmacoepidemiology and Drug Safety</i> , 2021, , .	0.9	0
260	913Factors of premature atrial contractions among general Japanese men. <i>International Journal of Epidemiology</i> , 2021, 50, .	0.9	0
261	Impact of resting heart rate on cardiovascular mortality according to serum albumin levels in a 24-year follow-up study on a general Japanese population: NIPPON DATA80. <i>Journal of Epidemiology</i> , 2021, , .	1.1	0
262	Effects of Changes in Lifestyle on Weight Loss before and after Intervention with Active Support Including Specific Health Guidance. <i>Health Evaluation and Promotion</i> , 2018, 45, 374-381.	0.0	0
263	Preface for the Special Issue â€œCardiovascular Risk Factors and Socioeconomic Status in Japan: NIPPON DATA2010â€œ. <i>Journal of Epidemiology</i> , 2018, 28, S1-S1.	1.1	0
264	Prediction of radiation pneumonitis using dose-volume histogram parameters with high attenuation in two types of cancer: A retrospective study. <i>PLoS ONE</i> , 2020, 15, e0244143.	1.1	0
265	Abstract P172: Relationship of Four Blood Pressure Indexes to Subclinical Cerebrovascular Diseases Assessed by Brain MRI in General Japanese Men. <i>Circulation</i> , 2020, 141, .	1.6	0
266	Title is missing!. , 2020, 15, e0244143.		0
267	Title is missing!. , 2020, 15, e0244143.		0
268	Title is missing!. , 2020, 15, e0244143.		0
269	Title is missing!. , 2020, 15, e0244143.		0
270	Title is missing!. , 2020, 15, e0244143.		0



#	ARTICLE	IF	CITATIONS
271	Title is missing!. , 2020, 15, e0244143.		0
272	Differential Association of Serum n-3 Polyunsaturated Fatty Acids with Various Cerebrovascular Lesions in Japanese Men. Cerebrovascular Diseases, 2022, 51, 774-780.	0.8	0