

Katsuyuki Miura

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3663077/katsuyuki-miura-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

248
papers

4,756
citations

35
h-index

60
g-index

278
ext. papers

6,320
ext. citations

4.1
avg. IF

5.26
L-index

#	Paper	IF	Citations
248	The 21-Year Trend of Stroke Incidence in a General Japanese Population: Results from the Takashima Stroke Registry, 1990-2010.. <i>Cerebrovascular Diseases</i> , 2022 , 1-7	3.2	0
247	Predictors of lower limb fractures in general Japanese: NIPPON DATA90.. <i>PLoS ONE</i> , 2022 , 17, e0261716	3.7	0
246	Trends in Prevalence, Treatment, and Control of Hypertension According to 40-Year-Old Life Expectancy at Prefectures in Japan from the National Health and Nutrition Surveys.. <i>Nutrients</i> , 2022 , 14,	6.7	1
245	Metabolic changes induced by dapagliflozin, an SGLT2 inhibitor, in Japanese patients with type 2 diabetes treated by oral anti-diabetic agents: A randomized, clinical trial.. <i>Diabetes Research and Clinical Practice</i> , 2022 , 186, 109781	7.4	0
244	Differential Association of Serum n-3 Polyunsaturated Fatty Acids with Various Cerebrovascular Lesions in Japanese Men.. <i>Cerebrovascular Diseases</i> , 2022 , 1-7	3.2	
243	A genome-wide association study on meat consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort study. <i>Journal of Nutritional Science</i> , 2021 , 10, e61	2.7	0
242	Risk Factors That Most Accurately Predict Coronary Artery Disease Based on the Duration of Follow-up - NIPPON DATA80. <i>Circulation Journal</i> , 2021 , 85, 908-913	2.9	1
241	Factors Associated with Lower Cognitive Performance Scores Among Older Japanese Men in Hawaii and Japan. <i>Journal of Alzheimer's Disease</i> , 2021 , 81, 403-412	4.3	
240	Association between socioeconomic status and prolonged television viewing time in a general Japanese population: NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , 2021 , 26, 57	4.2	3
239	Lipoprotein Particle Profiles Compared With Standard Lipids in the Association With Subclinical Aortic Valve Calcification in Apparently Healthy Japanese Men. <i>Circulation Journal</i> , 2021 , 85, 1076-1082	2.9	1
238	A Combination of Blood Pressure and Total Cholesterol Increases the Lifetime Risk of Coronary Heart Disease Mortality: EPOCH-JAPAN. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021 , 28, 6-24	4	4
237	Electrocardiographic Left Atrial Abnormality and B-Type Natriuretic Peptide in a General Japanese Population: NIPPON DATA2010. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021 , 28, 34-43	4	1
236	A genome-wide association study in Japanese identified one variant associated with a preference for a Japanese dietary pattern. <i>European Journal of Clinical Nutrition</i> , 2021 , 75, 937-945	5.2	2
235	Cluster-randomized controlled trial for the early promotion of clinic visits for untreated hypertension. <i>Hypertension Research</i> , 2021 , 44, 355-362	4.7	0
234	Differences between home blood pressure and strictly measured office blood pressure and their determinants in Japanese men. <i>Hypertension Research</i> , 2021 , 44, 80-87	4.7	2
233	A genome-wide association study on fish consumption in a Japanese population-the Japan Multi-Institutional Collaborative Cohort study. <i>European Journal of Clinical Nutrition</i> , 2021 , 75, 480-488	5.2	0
232	Relationship of higher-level functional capacity with long-term mortality in Japanese older people: NIPPON DATA90. <i>Journal of Epidemiology</i> , 2021 ,	3.4	2

231	A genome-wide association study on confection consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort Study. <i>British Journal of Nutrition</i> , 2021 , 126, 1843-1851	3.6	1
230	Relationship Between Calcium Intake and Impaired Activities of Daily Living in a Japanese Population: NIPPON DATA90. <i>Journal of Epidemiology</i> , 2021 , 31, 119-124	3.4	1
229	Sodium Intake and Incidence of Diabetes Complications in Elderly Patients with Type 2 Diabetes-Analysis of Data from the Japanese Elderly Diabetes Intervention Study (J-EDIT). <i>Nutrients</i> , 2021 , 13,	6.7	2
228	Association between socioeconomic status and physical inactivity in a general Japanese population: NIPPON DATA2010. <i>PLoS ONE</i> , 2021 , 16, e0254706	3.7	2
227	Coronary Artery Calcium Assessed Years Before Was Positively Associated With Subtle White Matter Injury of the Brain in Asymptomatic Middle-Aged Men: The Framingham Heart Study. <i>Circulation: Cardiovascular Imaging</i> , 2021 , 14, e011753	3.9	0
226	Relationship of Ambient Temperature Parameters to Stroke Incidence in a Japanese Population - Takashima Stroke Registry, Japan, 1988-2010. <i>Circulation Journal</i> , 2021 , 85, 2215-2221	2.9	
225	Seasonal Variation in Incidence of Stroke in a General Population of 1.4 Million Japanese: The Shiga Stroke Registry. <i>Cerebrovascular Diseases</i> , 2021 , 1-7	3.2	2
224	Developing a health economic model for Asians with type 2 diabetes based on the Japan Diabetes Complications Study and the Japanese Elderly Diabetes Intervention Trial. <i>BMJ Open Diabetes Research and Care</i> , 2021 , 9,	4.5	2
223	Estimation of 10-Year Risk of Death from Coronary Heart Disease, Stroke, and Cardiovascular Disease in a Pooled Analysis of Japanese Cohorts: EPOCH-JAPAN. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021 , 28, 816-825	4	4
222	Association between Stress-Coping Strategy and Functional Disability in the General Older Adult Population: The Takashima Study. <i>Gerontology</i> , 2021 , 1-8	5.5	0
221	Relationships of Alcohol Consumption with Coronary Risk Factors and Macro- and Micro-Nutrient Intake in Japanese People: The INTERLIPID Study. <i>Journal of Nutritional Science and Vitaminology</i> , 2021 , 67, 28-38	1.1	1
220	Relationship Between Step Counts and Cerebral Small Vessel Disease in Japanese Men. <i>Stroke</i> , 2020 , 51, 3584-3591	6.7	5
219	Smoking habits and progression of coronary and aortic artery calcification: A 5-year follow-up of community-dwelling Japanese men. <i>International Journal of Cardiology</i> , 2020 , 314, 89-94	3.2	3
218	Two-Year Recurrence After First-Ever Stroke in a General Population of 1.4 Million Japanese Patients - The Shiga Stroke and Heart Attack Registry Study. <i>Circulation Journal</i> , 2020 , 84, 943-948	2.9	2
217	Dynamic changes of mitral annulus in patients with degenerative mitral regurgitation and chronic atrial fibrillation undergoing mitral valve reconstruction. <i>General Thoracic and Cardiovascular Surgery</i> , 2020 , 68, 1405-1411	1.6	
216	Food Sources of Dietary Potassium in the Adult Japanese Population: The International Study of Macro-/Micronutrients and Blood Pressure (INTERMAP). <i>Nutrients</i> , 2020 , 12,	6.7	6
215	The Association Between Coronary Artery Calcification and Subclinical Cerebrovascular Diseases in Men: An Observational Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020 , 27, 995-1009	4	5
214	Epidemiology of hypertension in Japan: beyond the new 2019 Japanese guidelines. <i>Hypertension Research</i> , 2020 , 43, 1344-1351	4.7	13

213	Sorafenib exposure and its correlation with response and safety in advanced hepatocellular carcinoma: results from an observational retrospective study. <i>Cancer Chemotherapy and Pharmacology</i> , 2020 , 86, 129-139	3.5	4
212	Impact of hypertension stratified by diabetes on the lifetime risk of cardiovascular disease mortality in Japan: a pooled analysis of data from the Evidence for Cardiovascular Prevention from Observational Cohorts in Japan study. <i>Hypertension Research</i> , 2020 , 43, 1437-1444	4.7	3
211	Long-Term Survival after Stroke in 1.4 Million Japanese Population: Shiga Stroke and Heart Attack Registry. <i>Journal of Stroke</i> , 2020 , 22, 336-344	5.6	7
210	Exercise Habits Are Associated with Improved Long-Term Mortality Risks in the Nationwide General Japanese Population: A 20-Year Follow-Up of the NIPPON DATA90 Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020 , 252, 253-262	2.4	0
209	Prediction of radiation pneumonitis using dose-volume histogram parameters with high attenuation in two types of cancer: A retrospective study. <i>PLoS ONE</i> , 2020 , 15, e0244143	3.7	
208	Health status of workers approximately 60 years of age and the risk of early death after compulsory retirement: A cohort study. <i>Journal of Occupational Health</i> , 2020 , 62, e12088	2.3	3
207	Alcohol consumption and cognitive function in elderly Japanese men. <i>Alcohol</i> , 2020 , 85, 145-152	2.7	3
206	Liver fat accumulation assessed by computed tomography is an independent risk factor for diabetes mellitus in a population-based study: SESSA (Shiga Epidemiological Study of Subclinical Atherosclerosis). <i>Diabetes Research and Clinical Practice</i> , 2020 , 160, 108002	7.4	2
205	Waist Circumference and Domain-Specific Cognitive Function Among Non-Demented Japanese Older Adults Stratified by Sex: Results from the Takashima Cognition Study. <i>Journal of Alzheimer's Disease</i> , 2020 , 73, 887-896	4.3	6
204	Relationship of household salt intake level with long-term all-cause and cardiovascular disease mortality in Japan: NIPPON DATA80. <i>Hypertension Research</i> , 2020 , 43, 132-139	4.7	6
203	The association of fish consumption and its urinary metabolites with cardiovascular risk factors: the International Study of Macro-/Micronutrients and Blood Pressure (INTERMAP). <i>American Journal of Clinical Nutrition</i> , 2020 , 111, 280-290	7	18
202	Estimating the cost-effectiveness of screening for hepatitis C virus infection in Japan. <i>Hepatology Research</i> , 2020 , 50, 542-556	5.1	2
201	Isolated systolic hypertension and 29-year cardiovascular mortality risk in Japanese adults aged 30-49 years. <i>Journal of Hypertension</i> , 2020 , 38, 2230-2236	1.9	7
200	The relationship between serum levels of LOX-1 ligand containing ApoAI as a novel marker of dysfunctional HDL and coronary artery calcification in middle-aged Japanese men. <i>Atherosclerosis</i> , 2020 , 313, 20-25	3.1	1
199	Cross-sectional association of bone mineral density with coronary artery calcification in an international multi-ethnic population-based cohort of men aged 40-49: ERA JUMP study. <i>IJC Heart and Vasculature</i> , 2020 , 30, 100618	2.4	0
198	Conversion of Urine Protein-Creatinine Ratio or Urine Dipstick Protein to Urine Albumin-Creatinine Ratio for Use in Chronic Kidney Disease Screening and Prognosis : An Individual Participant-Based Meta-analysis. <i>Annals of Internal Medicine</i> , 2020 , 173, 426-435	8	39
197	Association between plant-based diets and blood pressure in the INTERMAP study. <i>BMJ Nutrition, Prevention and Health</i> , 2020 , 3, 133-142	6.7	0
196	Association between antidepressant use during pregnancy and congenital anomalies in children: A retrospective cohort study based on Japanese claims data. <i>Congenital Anomalies (discontinued)</i> , 2020 , 60, 180-188	1.1	3

195	Recent status of self-measured home blood pressure in the Japanese general population: a modern database on self-measured home blood pressure (MDAS). <i>Hypertension Research</i> , 2020 , 43, 1403-1412	4.7	3
194	Inclusion of a gene-environment interaction between alcohol consumption and the aldehyde dehydrogenase 2 genotype in a risk prediction model for upper aerodigestive tract cancer in Japanese men. <i>Cancer Science</i> , 2020 , 111, 3835-3844	6.9	3
193	The association between subjective health perception and lifestyle factors in Shiga prefecture, Japan: a cross-sectional study. <i>BMC Public Health</i> , 2020 , 20, 1786	4.1	4
192	Dietary Inflammatory Index Positively Associated With High-Sensitivity C-Reactive Protein Level in Japanese From NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2020 , 30, 98-107	3.4	10
191	Proteinuria and Reduced Estimated Glomerular Filtration Rate are Independently Associated With Lower Cognitive Abilities in Apparently Healthy Community-Dwelling Elderly Men in Japan: A Cross-sectional Study. <i>Journal of Epidemiology</i> , 2020 , 30, 244-252	3.4	3
190	Elevated Fasting Blood Glucose Levels Are Associated With Lower Cognitive Function, With a Threshold in Non-Diabetic Individuals: A Population-Based Study. <i>Journal of Epidemiology</i> , 2020 , 30, 121-127	3.4	1
189	Vitamin B6 intake and incidence of diabetic retinopathy in Japanese patients with type 2 diabetes: analysis of data from the Japan Diabetes Complications Study (JDCS). <i>European Journal of Nutrition</i> , 2020 , 59, 1585-1594	5.2	5
188	Relationship between carbohydrate and dietary fibre intake and the risk of cardiovascular disease mortality in Japanese: 24-year follow-up of NIPPON DATA80. <i>European Journal of Clinical Nutrition</i> , 2020 , 74, 67-76	5.2	4
187	Physical activity levels in American and Japanese men from the ERA-JUMP Study and associations with metabolic syndrome. <i>Journal of Sport and Health Science</i> , 2020 , 9, 170-178	8.2	2
186	Prediction of radiation pneumonitis using dose-volume histogram parameters with high attenuation in two types of cancer: A retrospective study 2020 , 15, e0244143		
185	Prediction of radiation pneumonitis using dose-volume histogram parameters with high attenuation in two types of cancer: A retrospective study 2020 , 15, e0244143		
184	Prediction of radiation pneumonitis using dose-volume histogram parameters with high attenuation in two types of cancer: A retrospective study 2020 , 15, e0244143		
183	Prediction of radiation pneumonitis using dose-volume histogram parameters with high attenuation in two types of cancer: A retrospective study 2020 , 15, e0244143		
182	Prediction of radiation pneumonitis using dose-volume histogram parameters with high attenuation in two types of cancer: A retrospective study 2020 , 15, e0244143		
181	Prediction of radiation pneumonitis using dose-volume histogram parameters with high attenuation in two types of cancer: A retrospective study 2020 , 15, e0244143		
180	Cardiovascular Risk Assessment Chart by Dietary Factors in Japan - NIPPON DATA80. <i>Circulation Journal</i> , 2019 , 83, 1254-1260	2.9	5
179	Comparison of blood pressure values-self-measured at home, measured at an unattended office, and measured at a conventional attended office. <i>Hypertension Research</i> , 2019 , 42, 1726-1737	4.7	12
178	Socioeconomic and lifestyle factors associated with depressive tendencies in general Japanese men and women: NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , 2019 , 24, 37	4.2	9

177	Salt intake and prevalence of overweight/obesity in Japan, China, the United Kingdom, and the United States: the INTERMAP Study. <i>American Journal of Clinical Nutrition</i> , 2019 , 110, 34-40	7	36
176	Association of Work Situation With Cardiovascular Disease Mortality Risk Among Working-Age Japanese Men - A 20-Year Follow-up of NIPPON DATA90. <i>Circulation Journal</i> , 2019 , 83, 1506-1513	2.9	4
175	Anthropometric Obesity Indices were Stronger than CT-Based Indices in Associations with Carotid Intima-Media Thickness in Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019 , 26, 1102-1114 ⁴		2
174	Urinary sodium-to-potassium ratio and intake of sodium and potassium among men and women from multiethnic general populations: the INTERSALT Study. <i>Hypertension Research</i> , 2019 , 42, 1590-1598 ^{4.7}	4.7	15
173	The impact of sex on risk of cardiovascular disease and all-cause mortality in adults with or without diabetes mellitus: A comparison between the U.S. and Japan. <i>Journal of Diabetes and Its Complications</i> , 2019 , 33, 417-423	3.2	6
172	Quantitative CT analysis of honeycombing area predicts mortality in idiopathic pulmonary fibrosis with definite usual interstitial pneumonia pattern: A retrospective cohort study. <i>PLoS ONE</i> , 2019 , 14, e0214278	3.7	8
171	Differences Between Coronary Artery Calcification and Aortic Artery Calcification in Relation to Cardiovascular Disease Risk Factors in Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019 , 26, 452-464	4	3
170	One-year weight loss maintenance outcomes following a worksite-based weight reduction program among Japanese men with cardiovascular risk factors. <i>Journal of Occupational Health</i> , 2019 , 61, 189-196 ^{2.3}	2.3	3
169	Coronary Artery Calcium Progression Among the US and Japanese Men. <i>Circulation: Cardiovascular Imaging</i> , 2019 , 12, e008104	3.9	3
168	Mutant KCNJ3 and KCNJ5 Potassium Channels as Novel Molecular Targets in Bradyarrhythmias and Atrial Fibrillation. <i>Circulation</i> , 2019 , 139, 2157-2169	16.7	23
167	Association of blood pressure with estimates of 24-h urinary sodium and potassium excretion from repeated single-spot urine samples. <i>Hypertension Research</i> , 2019 , 42, 411-418	4.7	15
166	The Relationship of Dietary Cholesterol with Serum Low-Density Lipoprotein Cholesterol and Confounding by Reverse Causality: The INTERLIPID Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019 , 26, 170-182	4	4
165	Associations of Overweight, Obesity, and Underweight With High Serum Total Cholesterol Level Over 30 Years Among the Japanese Elderly: NIPPON DATA 80, 90, and 2010. <i>Journal of Epidemiology</i> , 2019 , 29, 133-138	3.4	3
164	Association of blood levels of marine omega-3 fatty acids with coronary calcification and calcium density in Japanese men. <i>European Journal of Clinical Nutrition</i> , 2019 , 73, 783-792	5.2	13
163	The Japanese Society of Hypertension Guidelines for the Management of Hypertension (JSH 2019). <i>Hypertension Research</i> , 2019 , 42, 1235-1481	4.7	468
162	Intracranial Artery Stenosis and Its Association With Conventional Risk Factors in a General Population of Japanese Men. <i>Stroke</i> , 2019 , 50, 2967-2969	6.7	13
161	Serum long-chain n-3 polyunsaturated fatty acids and aortic calcification in middle-aged men: The population-based cross-sectional ERA-JUMP study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2019 , 29, 837-846	4.5	3
160	Carotid Intima-Media Thickness and Plaque in Apparently Healthy Japanese Individuals with an Estimated 10-Year Absolute Risk of CAD Death According to the Japan Atherosclerosis Society (JAS) Guidelines 2012: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2019 , 26, 716	4	

159	Association of Alcohol Consumption With Fat Deposition in a Community-Based Sample of Japanese Men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Epidemiology</i> , 2019 , 29, 205-212	3.4	6
158	Absolute risk score for stroke, myocardial infarction, and all cardiovascular disease: Japan Arteriosclerosis Longitudinal Study. <i>Hypertension Research</i> , 2019 , 42, 567-579	4.7	9
157	Lipoprotein particles and coronary artery calcium in middle-aged US-White and Japanese men. <i>Open Heart</i> , 2019 , 6, e001119	3	
156	IC-P-112: OBJECTIVELY MEASURED PHYSICAL ACTIVITY AND BRAIN VOLUME IN JAPANESE ADULT MEN: FINDINGS FROM THE SHIGA EPIDEMIOLOGICAL STUDY OF SUBCLINICAL ATHEROSCLEROSIS 2019 , 15, P96-P96		
155	Agreement between 24-h dietary recalls and 24-h urine collections for estimating sodium intake in China, Japan, UK, USA: the International Study of Macro- and Micro-nutrients and Blood Pressure. <i>Journal of Hypertension</i> , 2019 , 37, 814-819	1.9	9
154	The association of home and accurately measured office blood pressure with coronary artery calcification among general Japanese men. <i>Journal of Hypertension</i> , 2019 , 37, 1676-1681	1.9	4
153	Plasma C-peptide and glycated albumin and subsequent risk of cancer: From a large prospective case-cohort study in Japan. <i>International Journal of Cancer</i> , 2019 , 144, 718-729	7.5	2
152	Ipragliflozin, a sodium-glucose cotransporter 2 inhibitor, reduces bodyweight and fat mass, but not muscle mass, in Japanese type 2 diabetes patients treated with insulin: A randomized clinical trial. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 1012-1021	3.9	25
151	Factors associated with intra-individual visit-to-visit variability of blood pressure in four countries: the INTERMAP study. <i>Journal of Human Hypertension</i> , 2019 , 33, 229-236	2.6	3
150	The relationship between repeated measurement of casual and 24-h urinary sodium-to-potassium ratio in patients with chronic kidney disease. <i>Journal of Human Hypertension</i> , 2019 , 33, 286-297	2.6	4
149	Lifetime Risk of Stroke and Coronary Heart Disease Deaths According to Blood Pressure Level: EPOCH-JAPAN (Evidence for Cardiovascular Prevention From Observational Cohorts in Japan). <i>Hypertension</i> , 2019 , 73, 52-59	8.5	19
148	Adiposity and risk of decline in glomerular filtration rate: meta-analysis of individual participant data in a global consortium. <i>BMJ, The</i> , 2019 , 364, k5301	5.9	85
147	Having few remaining teeth is associated with a low nutrient intake and low serum albumin levels in middle-aged and older Japanese individuals: findings from the NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , 2019 , 24, 1	4.2	36
146	Association between antidepressant use during pregnancy and autism spectrum disorder in children: a retrospective cohort study based on Japanese claims data. <i>Maternal Health, Neonatology and Perinatology</i> , 2019 , 5, 1	3.4	9
145	Vegetable Protein Intake was Inversely Associated with Cardiovascular Mortality in a 15-Year Follow-Up Study of the General Japanese Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019 , 26, 198-206	4	12
144	Association Between Body Mass Index and All-Cause Death in Japanese Population: Pooled Individual Participant Data Analysis of 13 Cohort Studies. <i>Journal of Epidemiology</i> , 2019 , 29, 457-463	3.4	8
143	Meat intake and incidence of cardiovascular disease in Japanese patients with type 2 diabetes: analysis of the Japan Diabetes Complications Study (JDACS). <i>European Journal of Nutrition</i> , 2019 , 58, 281-290	5.3	9
142	Development of a High-Sensitivity Method for the Measurement of Human Nasal Aβ ₂ , Tau, and Phosphorylated Tau. <i>Journal of Alzheimer's Disease</i> , 2018 , 62, 737-744	4.3	7

141	Data on alcohol consumption and coronary artery calcification among asymptomatic middle-aged men for the ERA-JUMP study. <i>Data in Brief</i> , 2018 , 17, 1091-1098	1.2	2
140	Relation of Dietary Sodium (Salt) to Blood Pressure and Its Possible Modulation by Other Dietary Factors: The INTERMAP Study. <i>Hypertension</i> , 2018 , 71, 631-637	8.5	52
139	Seasonal variation in home blood pressure: findings from nationwide web-based monitoring in Japan. <i>BMJ Open</i> , 2018 , 8, e017351	3	24
138	Association of extremely high levels of high-density lipoprotein cholesterol with cardiovascular mortality in a pooled analysis of 9 cohort studies including 43,407 individuals: The EPOCH-JAPAN study. <i>Journal of Clinical Lipidology</i> , 2018 , 12, 674-684.e5	4.9	61
137	Brain Metabolite Changes in the Anterior Cingulate Cortex of Chronic Low Back Pain Patients and Correlations Between Metabolites and Psychological State. <i>Clinical Journal of Pain</i> , 2018 , 34, 657-663	3.5	10
136	Re-evaluation of the associations of egg intake with serum total cholesterol and cause-specific and total mortality in Japanese women. <i>European Journal of Clinical Nutrition</i> , 2018 , 72, 841-847	5.2	12
135	Association of Coronary Artery Calcification with Estimated Coronary Heart Disease Risk from Prediction Models in a Community-Based Sample of Japanese Men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2018 , 25, 477-489	4	3
134	International Comparison of Abdominal Fat Distribution Among Four Populations: The ERA-JUMP Study. <i>Metabolic Syndrome and Related Disorders</i> , 2018 , 16, 166-173	2.6	5
133	Overweight or underweight and the risk of decline in activities of daily living in a 22-year cohort study of a Japanese sample. <i>Geriatrics and Gerontology International</i> , 2018 , 18, 799-805	2.9	5
132	Plasma 25-hydroxyvitamin D concentration and subsequent risk of total and site specific cancers in Japanese population: large case-cohort study within Japan Public Health Center-based Prospective Study cohort. <i>BMJ, The</i> , 2018 , 360, k671	5.9	41
131	The Association between Glomerular Filtration Rate Estimated on Admission and Acute Stroke Outcome: The Shiga Stroke Registry. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018 , 25, 570-579	4	11
130	Serum magnesium, phosphorus, and calcium levels and subclinical calcific aortic valve disease: A population-based study. <i>Atherosclerosis</i> , 2018 , 273, 145-152	3.1	12
129	Dietary tofu intake and long-term risk of death from stroke in a general population. <i>Clinical Nutrition</i> , 2018 , 37, 182-188	5.9	7
128	Cardiovascular risk and blood pressure lowering treatment among elderly individuals: Evidence for Cardiovascular Prevention from Observational Cohorts in Japan. <i>Journal of Hypertension</i> , 2018 , 36, 410-418	1.9	5
127	Two-Year Survival After First-Ever Stroke in a General Population of 1.4 Million Japanese - Shiga Stroke Registry. <i>Circulation Journal</i> , 2018 , 82, 2549-2556	2.9	11
126	Change in Pericardial Fat Volume and Cardiovascular Risk Factors in a General Population of Japanese Men. <i>Circulation Journal</i> , 2018 , 82, 2542-2548	2.9	4
125	Comparison of carotid plaque burden among healthy middle-aged men living in the US, Japan, and South Korea. <i>International Journal of Cardiology</i> , 2018 , 266, 245-249	3.2	5
124	Rationale and Descriptive Analysis of Specific Health Guidance: the Nationwide Lifestyle Intervention Program Targeting Metabolic Syndrome in Japan. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018 , 25, 308-322	4	51

123	Effectiveness of a Self-monitoring Device for Urinary Sodium-to-Potassium Ratio on Dietary Improvement in Free-Living Adults: a Randomized Controlled Trial. <i>Journal of Epidemiology</i> , 2018 , 28, 41-47	3.4	21
122	Twelve-year trends of increasing overweight and obesity in patients with diabetes: the Shiga Diabetes Clinical Survey. <i>Endocrine Journal</i> , 2018 , 65, 527-536	2.9	11
121	Relationship of serum irisin levels to prevalence and progression of coronary artery calcification: A prospective, population-based study. <i>International Journal of Cardiology</i> , 2018 , 267, 177-182	3.2	15
120	Reduced Lung Function and Cerebral Small Vessel Disease in Japanese Men: the Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2018 , 25, 1009-1021	4	6
119	Effects of Changes in Lifestyle on Weight Loss before and after Intervention with Active Support Including Specific Health Guidance. <i>Health Evaluation and Promotion</i> , 2018 , 45, 374-381	0.1	
118	Preface for the Special Issue "Cardiovascular Risk Factors and Socioeconomic Status in Japan: NIPPON DATA2010". <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S1	3.4	
117	Association of alcohol consumption and aortic calcification in healthy men aged 40-49 years for the ERA JUMP Study. <i>Atherosclerosis</i> , 2018 , 268, 84-91	3.1	4
116	Stroke registries in the world: a systematic review. <i>Nosotchu</i> , 2018 , 40, 331-342	0.1	1
115	Relationships among Socioeconomic Factors and Self-rated Health in Japanese Adults: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S66-S72	3.4	9
114	Socioeconomic Inequalities in Oral Health among Middle-Aged and Elderly Japanese: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S59-S65	3.4	10
113	The Influence of the Japanese Nationwide Cardiovascular Prevention System Health Guidance on Smoking Cessation Among Smokers: A Propensity Score Matching Analysis. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018 , 25, 323-334	4	1
112	Dexmedetomidine attenuates the positive chronotropic effects of intravenous atropine in patients with bradycardia during spinal anaesthesia: a retrospective study. <i>JA Clinical Reports</i> , 2018 , 4, 70	0.6	0
111	Self-reported Sleep Duration and Subclinical Atherosclerosis in a General Population of Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018 , 25, 186-198	4	5
110	Variation of Risk Factors for Cause-Specific Reintubation: A Preliminary Study. <i>Canadian Respiratory Journal</i> , 2018 , 2018, 3654251	2.1	10
109	The National Integrated Project for Prospective Observation of Non-communicable Disease and its Trends in the Aged 2010 (NIPPON DATA2010): Objectives, Design, and Population Characteristics. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S2-S9	3.4	20
108	Relationships among Food Group Intakes, Household Expenditure, and Education Attainment in a General Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S23-S28	3.4	5
107	Relationship Between Socioeconomic Status and the Prevalence of Underweight, Overweight or Obesity in a General Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S10-S16	3.4	19
106	Macronutrient Intake and Socioeconomic Status: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S17-S22	3.4	11

105	Socioeconomic Status Associated With Urinary Sodium and Potassium Excretion in Japan: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S29-S34	3.4	11
104	Differences in Lifestyle Improvements With the Intention to Prevent Cardiovascular Diseases by Socioeconomic Status in a Representative Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S35-S39	3.4	2
103	Associations between Socioeconomic Status and the Prevalence and Treatment of Hypercholesterolemia in a General Japanese Population: NIPPON DATA2010. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018 , 25, 606-620	4	10
102	Passive Smoking at Home by Socioeconomic Factors in a Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S40-S45	3.4	3
101	Time-Related Changes in Relationships Between the Keys Score, Dietary Lipids, and Serum Total Cholesterol in Japan - NIPPON DATA80/90/2010. <i>Circulation Journal</i> , 2018 , 83, 147-155	2.9	2
100	Associations of High-Density Lipoprotein Particle and High-Density Lipoprotein Cholesterol With Alcohol Intake, Smoking, and Body Mass Index - The INTERLIPID Study. <i>Circulation Journal</i> , 2018 , 82, 2557-2565 ⁹	2.9	9
99	Socioeconomic Status and Knowledge of Cardiovascular Risk Factors: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S46-S52	3.4	9
98	The JAGUAR Score Predicts 1-Month Disability/Death in Ischemic Stroke Patient Ineligible for Recanalization Therapy. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2018 , 27, 2579-2586	2.8	2
97	Factors Related to Participation in Health Examinations for Japanese National Health Insurance: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S53-S58	3.4	4
96	Food sources of dietary sodium in the Japanese adult population: the international study of macro-/micronutrients and blood pressure (INTERMAP). <i>European Journal of Nutrition</i> , 2017 , 56, 1269-1280	5.3	17
95	Age-specific impact of diabetes mellitus on the risk of cardiovascular mortality: An overview from the evidence for Cardiovascular Prevention from Observational Cohorts in the Japan Research Group (EPOCH-JAPAN). <i>Journal of Epidemiology</i> , 2017 , 27, 123-129	3.4	22
94	Significant inverse association of equol-producer status with coronary artery calcification but not dietary isoflavones in healthy Japanese men. <i>British Journal of Nutrition</i> , 2017 , 117, 260-266	3.6	25
93	Diurnal variation of urinary sodium-to-potassium ratio in free-living Japanese individuals. <i>Hypertension Research</i> , 2017 , 40, 658-664	4.7	23
92	Associations of socioeconomic status with prevalence, awareness, treatment, and control of hypertension in a general Japanese population: NIPPON DATA2010. <i>Journal of Hypertension</i> , 2017 , 35, 401-408	1.9	59
91	Fasting but not casual blood glucose is associated with pancreatic cancer mortality in Japanese: EPOCH-JAPAN. <i>Cancer Causes and Control</i> , 2017 , 28, 625-633	2.8	13
90	Trends in antipsychotic prescriptions for Japanese outpatients during 2006-2012: a descriptive epidemiological study. <i>Pharmacoepidemiology and Drug Safety</i> , 2017 , 26, 642-656	2.6	17
89	Impacts of chronic kidney disease and diabetes on cardiovascular mortality in a general Japanese population: A 20-year follow-up of the NIPPON DATA90 study. <i>European Journal of Preventive Cardiology</i> , 2017 , 24, 505-513	3.9	5
88	Fiber-rich diet with brown rice improves endothelial function in type 2 diabetes mellitus: A randomized controlled trial. <i>PLoS ONE</i> , 2017 , 12, e0179869	3.7	38

87	Incidence, Management and Short-Term Outcome of Stroke in a General Population of 1.4 Million Japanese - Shiga Stroke Registry. <i>Circulation Journal</i> , 2017 , 81, 1636-1646	2.9	75
86	Exclusion of emphysematous lung from dose-volume estimates of risk improves prediction of radiation pneumonitis. <i>Radiation Oncology</i> , 2017 , 12, 160	4.2	4
85	Proton magnetic resonance spectroscopy assessment of metabolite status of the anterior cingulate cortex in chronic pain patients and healthy controls. <i>Journal of Pain Research</i> , 2017 , 10, 287-293	3.9	21
84	Volumetric changes in the aging rat brain and its impact on cognitive and locomotor functions. <i>Experimental Gerontology</i> , 2017 , 99, 69-79	4.5	34
83	Inclusion of a Genetic Risk Score into a Validated Risk Prediction Model for Colorectal Cancer in Japanese Men Improves Performance. <i>Cancer Prevention Research</i> , 2017 , 10, 535-541	3.2	17
82	Trends in medical performance in diabetic patients in primary care clinics compared with those in hospitals: Shiga Diabetes Clinical Survey, Japan, 2000-2012. <i>Diabetology International</i> , 2017 , 8, 59-68	2.3	6
81	Serum Ferritin, Insulin Resistance, and Ecell Dysfunction: A Prospective Study in Normoglycemic Japanese Men. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2017 , 125, 12-20	2.3	3
80	A pooled analysis of the association of isolated low levels of high-density lipoprotein cholesterol with cardiovascular mortality in Japan. <i>European Journal of Epidemiology</i> , 2017 , 32, 547-557	12.1	14
79	Increased Aortic Calcification Is Associated With Arterial Stiffness Progression in Multiethnic Middle-Aged Men. <i>Hypertension</i> , 2017 , 69, 102-108	8.5	37
78	Progression of coronary artery calcium in Japanese American men and white men in the ERA JUMP study. <i>International Journal of Cardiology</i> , 2017 , 228, 672-676	3.2	1
77	Effects of Estradiol on cold-sensitive receptor channel TRPM8 in ovariectomized rats. <i>Experimental Animals</i> , 2017 , 66, 337-343	1.8	9
76	Urinary Sodium-to-Potassium Ratio Tracks the Changes in Salt Intake during an Experimental Feeding Study Using Standardized Low-Salt and High-Salt Meals among Healthy Japanese Volunteers. <i>Nutrients</i> , 2017 , 9,	6.7	15
75	Is the Proportion of Carbohydrate Intake Associated with the Incidence of Diabetes Complications?-An Analysis of the Japan Diabetes Complications Study. <i>Nutrients</i> , 2017 , 9,	6.7	8
74	Time to Consider Use of the Sodium-to-Potassium Ratio for Practical Sodium Reduction and Potassium Increase. <i>Nutrients</i> , 2017 , 9,	6.7	55
73	Overall nutrient and total fat intake among Japanese people: The INTERLIPID Study Japan. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2017 , 26, 837-848	1	3
72	Estimating 24-h urinary sodium/potassium ratio from casual (RpotR urinary sodium/potassium ratio: the INTERSALT Study. <i>International Journal of Epidemiology</i> , 2017 , 46, 1564-1572	7.8	24
71	Dietary sodium-to-potassium ratio as a risk factor for stroke, cardiovascular disease and all-cause mortality in Japan: the NIPPON DATA80 cohort study. <i>BMJ Open</i> , 2016 , 6, e011632	3	53
70	Relation of unprocessed, processed red meat and poultry consumption to blood pressure in East Asian and Western adults. <i>Journal of Hypertension</i> , 2016 , 34, 1721-9	1.9	11

69	Relationship between non-high-density lipoprotein cholesterol and the long-term mortality of cardiovascular diseases: NIPPON DATA 90. <i>International Journal of Cardiology</i> , 2016 , 220, 262-7	3.2	16
68	Relationship of Insulin Resistance to Prevalence and Progression of Coronary Artery Calcification Beyond Metabolic Syndrome Components: Shiga Epidemiological Study of Subclinical Atherosclerosis. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2016 , 36, 1703-8	9.4	26
67	Lipoprotein-associated phospholipase A2 is related to risk of subclinical atherosclerosis but is not supported by Mendelian randomization analysis in a general Japanese population. <i>Atherosclerosis</i> , 2016 , 246, 141-7	3.1	36
66	Serum level of LOX-1 ligand containing ApoB is associated with increased carotid intima-media thickness in Japanese community-dwelling men, especially those with hypercholesterolemia LOX-1 ligand and IMT in Japanese. <i>Journal of Clinical Lipidology</i> , 2016 , 10, 172-80.e1	4.9	10
65	Associations of serum LDL particle concentration with carotid intima-media thickness and coronary artery calcification. <i>Journal of Clinical Lipidology</i> , 2016 , 10, 1195-1202.e1	4.9	9
64	Relationship of three different types of low-carbohydrate diet to cardiometabolic risk factors in a Japanese population: the INTERMAP/INTERLIPID Study. <i>European Journal of Nutrition</i> , 2016 , 55, 1515-24	5.2	6
63	Independent Prognostic Value of Single and Multiple Non-Specific 12-Lead Electrocardiographic Findings for Long-Term Cardiovascular Outcomes: A Prospective Cohort Study. <i>PLoS ONE</i> , 2016 , 11, e0157563	3.7	3
62	The Relationship between Very High Levels of Serum High-Density Lipoprotein Cholesterol and Cause-Specific Mortality in a 20-Year Follow-Up Study of Japanese General Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2016 , 23, 800-9	4	33
61	Perioperative Adiponectin Measurement is Useful for Prediction of Postoperative Infection in Patients with Colorectal Cancer. <i>Annals of Surgical Oncology</i> , 2016 , 23, 540-545	3.1	4
60	The impact of equol-producing status in modifying the effect of soya isoflavones on risk factors for CHD: a systematic review of randomised controlled trials. <i>Journal of Nutritional Science</i> , 2016 , 5, e30	2.7	16
59	Cross-sectional association between exposure to particulate matter and inflammatory markers in the Japanese general population: NIPPON DATA2010. <i>Environmental Pollution</i> , 2016 , 213, 460-467	9.3	13
58	Does the flushing response modify the relationship between alcohol intake and hypertension in the Japanese population? NIPPON DATA2010. <i>Hypertension Research</i> , 2016 , 39, 670-9	4.7	8
57	Maximum BMI and microvascular complications in a cohort of Japanese patients with type 2 diabetes: the Japan Diabetes Complications Study. <i>Journal of Diabetes and Its Complications</i> , 2016 , 30, 790-7	3.2	5
56	Smoking, Smoking Cessation, and Measures of Subclinical Atherosclerosis in Multiple Vascular Beds in Japanese Men. <i>Journal of the American Heart Association</i> , 2016 , 5,	6	26
55	Mendelian randomization analysis in three Japanese populations supports a causal role of alcohol consumption in lowering low-density lipid cholesterol levels and particle numbers. <i>Atherosclerosis</i> , 2016 , 254, 242-248	3.1	18
54	Validation of the european SCORE risk chart in the healthy middle-aged Japanese. <i>Atherosclerosis</i> , 2016 , 252, 116-121	3.1	6
53	Lifetime cigarette smoking is associated with abdominal obesity in a community-based sample of Japanese men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Preventive Medicine Reports</i> , 2016 , 4, 225-32	2.6	26
52	Long-term outcomes associated with prolonged PR interval in the general Japanese population. <i>International Journal of Cardiology</i> , 2015 , 184, 291-293	3.2	9

51	Continuous decline in mortality from coronary heart disease in Japan despite a continuous and marked rise in total cholesterol: Japanese experience after the Seven Countries Study. <i>International Journal of Epidemiology</i> , 2015 , 44, 1614-24	7.8	20
50	Brachial-ankle pulse wave velocity is associated with coronary calcification among 1131 healthy middle-aged men. <i>International Journal of Cardiology</i> , 2015 , 189, 67-72	3.2	20
49	Relationship of type of work with health-related quality of life. <i>Quality of Life Research</i> , 2015 , 24, 2927-33	7	9
48	Secular trends of the impact of overweight and obesity on hypertension in Japan, 1980-2010. <i>Hypertension Research</i> , 2015 , 38, 790-5	4.7	32
47	Smoking increases the risk of all-cause and cardiovascular mortality in patients with chronic kidney disease. <i>Kidney International</i> , 2015 , 88, 1144-52	9.9	24
46	Analysis of factors associated with maintenance discontinuation in implant patients. <i>SpringerPlus</i> , 2015 , 4, 767		1
45	Scientific evidences for the cut-off points of blood pressure. <i>Health Evaluation and Promotion</i> , 2015 , 42, 280-286	0.1	
44	Alcohol consumption, hospitalization and medical expenditure: a large epidemiological study on the medical insurance system in Japan. <i>Alcohol and Alcoholism</i> , 2015 , 50, 236-43	3.5	1
43	Holter monitoring for the screening of cardiac disease in diabetes mellitus: The non-invasive Holter monitoring observation of new cardiac events in diabetics study. <i>Diabetes and Vascular Disease Research</i> , 2015 , 12, 396-404	3.3	7
42	High-density lipoprotein particle concentration and subclinical atherosclerosis of the carotid arteries in Japanese men. <i>Atherosclerosis</i> , 2015 , 239, 444-50	3.1	16
41	Cumulative impact of axial, structural, and repolarization ECG findings on long-term cardiovascular mortality among healthy individuals in Japan: National Integrated Project for Prospective Observation of Non-Communicable Disease and its Trends in the Aged, 1980 and 1990. <i>European Journal of Preventive Cardiology</i> , 2014 , 21, 1501-8	3.9	12
40	Lipoprotein particle profiles compared with standard lipids in association with coronary artery calcification in the general Japanese population. <i>Atherosclerosis</i> , 2014 , 236, 237-43	3.1	17
39	HOMA-IR and the risk of hyperuricemia: a prospective study in non-diabetic Japanese men. <i>Diabetes Research and Clinical Practice</i> , 2014 , 106, 154-60	7.4	18
38	Long-chain n-3 polyunsaturated fatty acids intake and cardiovascular disease mortality risk in Japanese: a 24-year follow-up of NIPPON DATA80. <i>Atherosclerosis</i> , 2014 , 232, 384-9	3.1	42
37	Statin use and risk of cerebral aneurysm rupture: a hospital-based case-control study in Japan. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2014 , 23, 343-8	2.8	50
36	Cross-sectional comparison of coronary artery calcium scores between Caucasian men in the United States and Japanese men in Japan: the multi-ethnic study of atherosclerosis and the Shiga epidemiological study of subclinical atherosclerosis. <i>American Journal of Epidemiology</i> , 2014 , 180, 590-8	3.8	29
35	Six random specimens of daytime casual urine on different days are sufficient to estimate daily sodium/potassium ratio in comparison to 7-day 24-h urine collections. <i>Hypertension Research</i> , 2014 , 37, 765-71	4.7	43
34	Low-carbohydrate diets and cardiovascular and total mortality in Japanese: a 29-year follow-up of NIPPON DATA80. <i>British Journal of Nutrition</i> , 2014 , 112, 916-24	3.6	35

33	Overall sleep status and high sensitivity C-reactive protein: a prospective study in Japanese factory workers. <i>Journal of Sleep Research</i> , 2014 , 23, 717-727	5.8	2
32	High long-chain n-3 fatty acid intake attenuates the effect of high resting heart rate on cardiovascular mortality risk: a 24-year follow-up of Japanese general population. <i>Journal of Cardiology</i> , 2014 , 64, 218-24	3	9
31	Impact of metabolic syndrome on the risk of cardiovascular disease mortality in the United States and in Japan. <i>American Journal of Cardiology</i> , 2014 , 113, 84-9	3	60
30	Associations between rice, noodle, and bread intake and sleep quality in Japanese men and women. <i>PLoS ONE</i> , 2014 , 9, e105198	3.7	18
29	A cross-sectional association of obesity with coronary calcium among Japanese, Koreans, Japanese Americans, and U.S. whites. <i>European Heart Journal Cardiovascular Imaging</i> , 2013 , 14, 921-7	4.1	16
28	[Scientific statement] Report of the Salt Reduction Committee of the Japanese Society of Hypertension(2) Goal and strategies of dietary salt reduction in the management of hypertension. <i>Hypertension Research</i> , 2013 , 36, 1020-5	4.7	15
27	Interaction between dietary marine-derived n-3 fatty acids intake and J-point elevation on the risk of cardiac death: a 24-year follow-up of Japanese men. <i>Heart</i> , 2013 , 99, 1024-9	5.1	6
26	Association of total marine fatty acids, eicosapentaenoic and docosahexaenoic acids, with aortic stiffness in Koreans, whites, and Japanese Americans. <i>American Journal of Hypertension</i> , 2013 , 26, 1321-7	7.3	8
25	Relation of Serum Leptin and Adiponectin Level to Serum C-Reactive Protein: The INTERLIPID Study. <i>International Journal of Vascular Medicine</i> , 2013 , 2013, 601364	1.2	9
24	HbA1c and the risks for all-cause and cardiovascular mortality in the general Japanese population: NIPPON DATA90. <i>Diabetes Care</i> , 2013 , 36, 3759-65	14.6	42
23	Epidemiology of hypertension in Japan: where are we now?. <i>Circulation Journal</i> , 2013 , 77, 2226-31	2.9	130
22	Epidemiology of cardiovascular risk factors in Asian countries. <i>Circulation Journal</i> , 2013 , 77, 2851-9	2.9	44
21	Relationship of dietary monounsaturated fatty acids to blood pressure: the International Study of Macro/Micronutrients and Blood Pressure. <i>Journal of Hypertension</i> , 2013 , 31, 1144-50	1.9	25
20	Coronary artery calcification by computed tomography in epidemiologic research and cardiovascular disease prevention. <i>Journal of Epidemiology</i> , 2012 , 22, 188-98	3.4	23
19	Diet, Nutrients, and the Prevention of Hypertension. <i>Current Nutrition Reports</i> , 2012 , 1, 87-92	6	0
18	Relation between serum total cholesterol level and cardiovascular disease stratified by sex and age group: a pooled analysis of 65 594 individuals from 10 cohort studies in Japan. <i>Journal of the American Heart Association</i> , 2012 , 1, e001974	6	64
17	Influence of smoking combined with another risk factor on the risk of mortality from coronary heart disease and stroke: pooled analysis of 10 Japanese cohort studies. <i>Cerebrovascular Diseases</i> , 2012 , 33, 480-91	3.2	50
16	Blood pressure categories and long-term risk of cardiovascular disease according to age group in Japanese men and women. <i>Hypertension Research</i> , 2012 , 35, 947-53	4.7	87

15	Comparability in coronary artery calcium scores on CT scan between two community-based cohort studies. <i>International Journal of Cardiology</i> , 2011 , 149, 244-245	3.2	7
14	Population attributable numbers and fractions of deaths due to smoking: a pooled analysis of 180,000 Japanese. <i>Preventive Medicine</i> , 2011 , 52, 60-5	4.3	45
13	Relationship of dietary cholesterol to blood pressure: the INTERMAP study. <i>Journal of Hypertension</i> , 2011 , 29, 222-8	1.9	37
12	Epidemiology and prevention of hypertension in Japanese: how could Japan get longevity?. <i>EPMA Journal</i> , 2011 , 2, 59-64	8.8	28
11	Relationship between 5-year decline in instrumental activity of daily living and accumulation of cardiovascular risk factors: NIPPON DATA90. <i>Journal of Atherosclerosis and Thrombosis</i> , 2010 , 17, 64-72	4	17
10	Serum n-6 fatty acids and lipoprotein subclasses in middle-aged men: the population-based cross-sectional ERA-JUMP study. <i>American Journal of Clinical Nutrition</i> , 2010 , 91, 1195-203	7	35
9	Integration of data from NIPPON DATA80/90 and National Nutrition Survey in Japan: for cohort studies of representative Japanese on nutrition. <i>Journal of Epidemiology</i> , 2010 , 20 Suppl 3, S506-14	3.4	35
8	Dietary salt intake and blood pressure in a representative Japanese population: baseline analyses of NIPPON DATA80. <i>Journal of Epidemiology</i> , 2010 , 20 Suppl 3, S524-30	3.4	38
7	Dietary intake of potassium and associated dietary factors among representative samples of Japanese general population: NIPPON DATA 80/90. <i>Journal of Epidemiology</i> , 2010 , 20 Suppl 3, S567-75	3.4	7
6	NIPPON DATA80/90 Nutrition Study: Appendix Tables. <i>Journal of Epidemiology</i> , 2010 , 20, S587-S596	3.4	0
5	Dietary sources of sodium in China, Japan, the United Kingdom, and the United States, women and men aged 40 to 59 years: the INTERMAP study. <i>Journal of the American Dietetic Association</i> , 2010 , 110, 736-45		359
4	Four blood pressure indexes and the risk of stroke and myocardial infarction in Japanese men and women: a meta-analysis of 16 cohort studies. <i>Circulation</i> , 2009 , 119, 1892-8	16.7	88
3	Relationship of dietary linoleic acid to blood pressure. The International Study of Macro-Micronutrients and Blood Pressure Study [corrected]. <i>Hypertension</i> , 2008 , 52, 408-14	8.5	60
2	Cardiovascular disease and risk factors in Asia: a selected review. <i>Circulation</i> , 2008 , 118, 2702-9	16.7	500
1	Eighteen-year trends in the management of patients with diabetes in the Shiga Diabetes Clinical Survey: overall trends and differences by age group. <i>Diabetology International</i> , 1	2.3	0