

# Peizeng Yang

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3663416/publications.pdf>

Version: 2024-02-01

259  
papers

7,281  
citations

108046

37  
h-index

129628

63  
g-index

280  
all docs

280  
docs citations

280  
times ranked

5576  
citing authors

#	ARTICLE	IF	CITATIONS
1	Outcome Measures for Disease Monitoring in Intraocular Inflammatory and Infectious Diseases (OCTOMERIA): Understanding the Choroid in Uveitis with Optical Coherence Tomography (OCT). <i>Ocular Immunology and Inflammation</i> , 2023, 31, 374-392.	1.0	4
2	Succinic acid exacerbates experimental autoimmune uveitis by stimulating neutrophil extracellular traps formation via SUCNR1 receptor. <i>British Journal of Ophthalmology</i> , 2023, 107, 1744-1749.	2.1	5
3	Association between temperature changes and uveitis onset in mainland China. <i>British Journal of Ophthalmology</i> , 2022, 106, 91-96.	2.1	10
4	Development of revised diagnostic criteria for Fuchs's uveitis syndrome in a Chinese population. <i>British Journal of Ophthalmology</i> , 2022, 106, 1678-1683.	2.1	7
5	Identification of differently expressed mRNAs by peripheral blood mononuclear cells in Vogt-Koyanagi-Harada disease. <i>Genes and Diseases</i> , 2022, 9, 1378-1388.	1.5	4
6	Association between Fine Particulate Air Pollution and the Onset of Uveitis in Mainland China. <i>Ocular Immunology and Inflammation</i> , 2022, 30, 1810-1815.	1.0	4
7	Identification of Novel Risk Loci for Behçet's Disease-Related Uveitis in a Chinese Population in a <scp>Genome-Wide</scp> Association Study. <i>Arthritis and Rheumatology</i> , 2022, 74, 671-681.	2.9	14
8	Surveillance of Liver Function in Uveitis with or without Chronic HBV Infection. <i>Ophthalmic Research</i> , 2022, 65, 94-103.	1.0	0
9	Evaluation of sensitivity and specificity of diagnostic criteria for Behçet's disease in the absence of a gold standard. <i>Rheumatology</i> , 2022, 61, 3667-3676.	0.9	7
10	Progranulin Suppressed Autoimmune Uveitis and Autoimmune Neuroinflammation by Inhibiting Th1/Th17 Cells and Promoting Treg Cells and M2 Macrophages. <i>Neurology: Neuroimmunology and NeuroInflammation</i> , 2022, 9, .	3.1	20
11	Visualizing lipid behind the retina in aging and age-related macular degeneration, via indocyanine green angiography (ASHS-LIA). <i>Eye</i> , 2022, 36, 1735-1746.	1.1	12
12	Genetic and Clinical Features of Blau Syndrome among Chinese Patients with Uveitis. <i>Ophthalmology</i> , 2022, 129, 821-828.	2.5	5
13	Genetically predicted fasting blood glucose level plays a causal role in intraocular pressure: A Mendelian randomisation study. <i>Clinical and Experimental Ophthalmology</i> , 2022, , .	1.3	0
14	Average corticosteroid dose and risk for HBV reactivation and hepatitis flare in patients with resolved hepatitis B infection. <i>Annals of the Rheumatic Diseases</i> , 2022, 81, 584-591.	0.5	11
15	SNP rs7130280 in lncRNA NONHSAT159216.1 confers susceptibility to Behçet's disease uveitis in a Chinese Han population. <i>Rheumatology</i> , 2022, 62, 384-396.	0.9	3
16	PD-1 Targeted Nanoparticles Inhibit Activated T Cells and Alleviate Autoimmunity via Suppression of Cellular Energy Metabolism Mediated by PKM2. <i>International Journal of Nanomedicine</i> , 2022, Volume 17, 1711-1724.	3.3	5
17	Prevalence, risk factors and management of ocular hypertension or glaucoma in patients with Vogt-Koyanagi-Harada disease. <i>British Journal of Ophthalmology</i> , 2021, 105, 1678-1682.	2.1	7
18	Prevalence and clinical features of systemic diseases in Chinese patients with uveitis. <i>British Journal of Ophthalmology</i> , 2021, 105, 75-82.	2.1	37

#	ARTICLE	IF	CITATIONS
19	Activation of the interleukin-23/interleukin-17 signalling pathway in autoinflammatory and autoimmune uveitis. <i>Progress in Retinal and Eye Research</i> , 2021, 80, 100866.	7.3	104
20	Higher 25-hydroxyvitamin D level is associated with increased risk for Behçet's disease. <i>Clinical Nutrition</i> , 2021, 40, 518-524.	2.3	12
21	Linoleic acid inhibits in vitro function of human and murine dendritic cells, CD4+T cells and retinal pigment epithelial cells. <i>Graefe's Archive for Clinical and Experimental Ophthalmology</i> , 2021, 259, 987-998.	1.0	8
22	Association of a CARD9 Gene Haplotype with Behçet's Disease in a Chinese Han Population. <i>Ocular Immunology and Inflammation</i> , 2021, 29, 219-227.	1.0	5
23	Metabolomic Analysis of Aqueous Humor Identifies Aberrant Amino Acid and Fatty Acid Metabolism in Vogt-Koyanagi-Harada and Behçet's Disease. <i>Frontiers in Immunology</i> , 2021, 12, 587393.	2.2	11
24	Identification of Urine Metabolic Biomarkers for Vogt-Koyanagi-Harada Disease. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 637489.	1.8	10
25	High Ambient Temperature Aggravates Experimental Autoimmune Uveitis Symptoms. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 629306.	1.8	8
26	Tuberculosis Exposure With Risk of Behçet Disease Among Patients With Uveitis. <i>JAMA Ophthalmology</i> , 2021, 139, 415.	1.4	12
27	Ocular involvement in extranodal natural-killer T-cell lymphoma. <i>Lancet Haematology</i> , 2021, 8, e382.	2.2	2
28	Specific sweat metabolite profile in ocular Behçet's disease. <i>International Immunopharmacology</i> , 2021, 97, 107812.	1.7	12
29	Changes in the Gut Microbiome Contribute to the Development of Behçet's Disease via Adjuvant Effects. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 716760.	1.8	9
30	Auxiliary Ocular Examinations. , 2021, , 81-138.		1
31	Meta-Analysis of miRNA Variants Associated with Susceptibility to Autoimmune Disease. <i>Disease Markers</i> , 2021, 2021, 1-21.	0.6	4
32	SNP-mediated binding of TBX1 to the enhancer element of IL-10 reduces the risk of Behçet's disease. <i>Epigenomics</i> , 2021, 13, 1523-1537.	1.0	1
33	Complications and Their Management. , 2021, , 165-168.		0
34	Acute Anterior Uveitis. , 2021, , 171-185.		0
35	Intermediate Uveitis. , 2021, , 187-193.		0
36	Uveitis-associated with Juvenile Idiopathic Arthritis. , 2021, , 307-325.		0

#	ARTICLE	IF	CITATIONS
37	Vogt-Koyanagi-Harada Disease. , 2021, , 411-537.		2
38	Retinal Vasculitis. , 2021, , 563-587.		0
39	Ocular Sarcoidosis. , 2021, , 611-625.		0
40	Ocular Examinations. , 2021, , 19-79.		0
41	Scleritis. , 2021, , 825-857.		0
42	Steroid-sparing Immunosuppressive Agents. , 2021, , 155-161.		0
43	Optical Coherence Tomographic Features and Prognostic Values of Macular Edema in Vogt-Koyanagi-Harada Disease. <i>Frontiers in Medicine</i> , 2021, 8, 772439.	1.2	1
44	A Single-Cell Transcriptome Atlas of the Human Retinal Pigment Epithelium. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 802457.	1.8	15
45	Analyses of circRNA and mRNA Profiles in Vogt-Koyanagi-Harada Disease. <i>Frontiers in Immunology</i> , 2021, 12, 738760.	2.2	7
46	Decreased Expression of TGR5 in Vogt-Koyanagi-Harada (VKH) Disease. <i>Ocular Immunology and Inflammation</i> , 2020, 28, 200-208.	1.0	5
47	Genetic aspects of idiopathic paediatric uveitis and juvenile idiopathic arthritis associated uveitis in Chinese Han. <i>British Journal of Ophthalmology</i> , 2020, 104, 443-447.	2.1	4
48	Association of toll-like receptor 10 polymorphisms with paediatric idiopathic uveitis in Han Chinese. <i>British Journal of Ophthalmology</i> , 2020, 104, 1467-1471.	2.1	5
49	Uveitis genetics. <i>Experimental Eye Research</i> , 2020, 190, 107853.	1.2	36
50	Association of apoptosis genes in PDCD1 but not PDCD1LG2, FAS, and FASLG with pediatric idiopathic uveitis in Han Chinese. <i>Pediatric Research</i> , 2020, 87, 634-638.	1.1	6
51	Altered gut microbiome composition in patients with Vogt-Koyanagi-Harada disease. <i>Gut Microbes</i> , 2020, 11, 539-555.	4.3	52
52	Aqueous cytokine levels in four common uveitis entities. <i>International Immunopharmacology</i> , 2020, 78, 106021.	1.7	18
53	The haplotypes of various TNF related genes associated with scleritis in Chinese Han. <i>Human Genomics</i> , 2020, 14, 46.	1.4	3
54	Sharing of Genetic Association Signals by Age-Related Macular Degeneration and Alzheimer's Disease at Multiple Levels. <i>Molecular Neurobiology</i> , 2020, 57, 4488-4499.	1.9	7

#	ARTICLE	IF	CITATIONS
55	Small molecules targeting ROR $\gamma$ t inhibit autoimmune disease by suppressing Th17 cell differentiation. <i>Cell Death and Disease</i> , 2020, 11, 697.	2.7	32
56	The Rs12569232 SNP Association with Vogt-Koyanagi-Harada Disease and Behcet's Disease is Probably Mediated by Regulation of Linc00467 Expression. <i>Ocular Immunology and Inflammation</i> , 2020, , 1-7.	1.0	3
57	Increased Expression of Indoleamine 2,3-Dioxygenase (IDO) in Vogt-Koyanagi-Harada (VKH) Disease May Lead to a Shift of T Cell Responses Toward a Treg Population. <i>Inflammation</i> , 2020, 43, 1780-1788.	1.7	3
58	Effect of berberine on spleen transcriptome and gut microbiota composition in experimental autoimmune uveitis. <i>International Immunopharmacology</i> , 2020, 81, 106270.	1.7	14
59	Integrated omics analysis of sweat reveals an aberrant amino acid metabolism pathway in Vogt's Koyanagi-Harada disease. <i>Clinical and Experimental Immunology</i> , 2020, 200, 250-259.	1.1	12
60	Identification of Ribosomal Protein S4, Y-Linked 1 as a cyclosporin A plus corticosteroid resistance gene. <i>Journal of Autoimmunity</i> , 2020, 112, 102465.	3.0	10
61	Plasma metabolomics study of Vogt-Koyanagi-Harada disease identifies potential diagnostic biomarkers. <i>Experimental Eye Research</i> , 2020, 196, 108070.	1.2	13
62	Analysis of the role of palmitoleic acid in acute anterior uveitis. <i>International Immunopharmacology</i> , 2020, 84, 106552.	1.7	8
63	Vogt's Koyanagi-Harada Disease. <i>Retina Atlas</i> , 2020, , 67-75.	0.0	1
64	Integrated Analysis of Key Pathways and Drug Targets Associated With Vogt-Koyanagi-Harada Disease. <i>Frontiers in Immunology</i> , 2020, 11, 587443.	2.2	11
65	Weak association of a TNFRSF1A polymorphism with Behcet's disease in Chinese Han. <i>Experimental Eye Research</i> , 2020, 196, 108045.	1.2	1
66	Macular Abnormalities in Vogt-Koyanagi-Harada Disease. <i>Ocular Immunology and Inflammation</i> , 2019, 27, 1195-1202.	1.0	15
67	Association of TLR2 Gene Polymorphisms with Presumed Viral-Induced Anterior Uveitis in male Han Chinese. <i>Experimental Eye Research</i> , 2019, 187, 107777.	1.2	1
68	Diagnosis and treatment of human sparganosis. <i>Lancet Infectious Diseases</i> , The, 2019, 19, 577-578.	4.6	3
69	Different Methylation of CpG-SNPs in Behcet's Disease. <i>BioMed Research International</i> , 2019, 2019, 1-7.	0.9	3
70	Comparison of Clinical Features and Visual Outcome between Sympathetic Ophthalmia and Vogt's Koyanagi-Harada Disease in Chinese Patients. <i>Ophthalmology</i> , 2019, 126, 1297-1305.	2.5	30
71	Epigenome-wide association study identifies Behcet's disease-associated methylation loci in Han Chinese. <i>Rheumatology</i> , 2019, 58, 1574-1584.	0.9	21
72	Identification of susceptibility SNPs in CTLA-4 and PTPN22 for scleritis in Han Chinese. <i>Clinical and Experimental Immunology</i> , 2019, 197, 230-236.	1.1	6

#	ARTICLE	IF	CITATIONS
73	A 50-year-old woman with a recurrent eyelid swelling. <i>Lancet Infectious Diseases</i> , The, 2019, 19, 338.	4.6	4
74	Replication of Genome-Wide Association Analysis Identifies New Susceptibility Loci at Long Noncoding RNA Regions for Vogt-Koyanagi-Harada Disease. , 2019, 60, 4820.		7
75	UVEOGENE: An SNP database for investigations on genetic factors associated with uveitis and their relationship with other systemic autoimmune diseases. <i>Human Mutation</i> , 2019, 40, 258-266.	1.1	12
76	Clinical features of Chinese patients with relapsing polychondritis. <i>British Journal of Ophthalmology</i> , 2019, 103, 1129-1132.	2.1	8
77	Causes of Visual Impairment and Blindness in the 2006 and 2014 Nine-Province Surveys in Rural China. <i>American Journal of Ophthalmology</i> , 2019, 197, 80-87.	1.7	32
78	Long-Term Efficacy and Safety of Interferon Alpha-2a in the Treatment of Chinese Patients with Behçet's Uveitis Not Responding to Conventional Therapy. <i>Ocular Immunology and Inflammation</i> , 2019, 27, 7-14.	1.0	20
79	Decreased interleukin(IL)-35 Expression is Associated with Active Intraocular Inflammation in Vogt-Koyanagi-Harada (VKH) Disease. <i>Ocular Immunology and Inflammation</i> , 2019, 27, 595-601.	1.0	6
80	ERAP1/ERAP2 and RUNX3 polymorphisms are not associated with ankylosing spondylitis susceptibility in Chinese Han. <i>Clinical and Experimental Immunology</i> , 2018, 193, 95-102.	1.1	13
81	Association of genetic variations in PTPN2 and CD122 with ocular Behçet's disease. <i>British Journal of Ophthalmology</i> , 2018, 102, 996-1002.	2.1	7
82	Longitudinal Study of Visual Function in Vogt-Koyanagi-Harada Disease Using Full-Field Electroretinography. <i>American Journal of Ophthalmology</i> , 2018, 191, 92-99.	1.7	6
83	Decreased expression of A20 is associated with ocular Behçet's disease (BD) but not with Vogt-Koyanagi-Harada (VKH) disease. <i>British Journal of Ophthalmology</i> , 2018, 102, 1167-1172.	2.1	11
84	Analysis of the association between Fc receptor family gene polymorphisms and ocular Behçet's disease in Han Chinese. <i>Scientific Reports</i> , 2018, 8, 4850.	1.6	5
85	Clinical Features and Complications of Scleritis in Chinese Patients. <i>Ocular Immunology and Inflammation</i> , 2018, 26, 387-396.	1.0	38
86	Clinical features of HLA-B27-positive acute anterior uveitis with or without ankylosing spondylitis in a Chinese cohort. <i>British Journal of Ophthalmology</i> , 2018, 102, 215-219.	2.1	50
87	Prevalence of Vision Impairment in Older Adults in Rural China in 2014 and Comparisons With the 2006 China Nine-Province Survey. <i>American Journal of Ophthalmology</i> , 2018, 185, 81-93.	1.7	48
88	MicroRNA-20a-5p suppresses IL-17 production by targeting OSM and CCL1 in patients with Vogt-Koyanagi-Harada disease. <i>British Journal of Ophthalmology</i> , 2018, 102, 282-290.	2.1	31
89	The Choroidal Vascularity Index Decreases and Choroidal Thickness Increases in Vogt-Koyanagi-Harada Disease Patients During a Recurrent Anterior Uveitis Attack. <i>Ocular Immunology and Inflammation</i> , 2018, 26, 1237-1243.	1.0	33
90	Multispectral image analysis in Vogt-Koyanagi-Harada disease. <i>Acta Ophthalmologica</i> , 2018, 96, 411-419.	0.6	13

#	ARTICLE	IF	CITATIONS
91	Novel treatment regimen of Vogt-Koyanagi-Harada disease with a reduced dose of corticosteroids combined with immunosuppressive agents. <i>Current Eye Research</i> , 2018, 43, 254-261.	0.7	39
92	How To Deal With Uveitis Patients?. <i>Current Molecular Medicine</i> , 2018, 17, 468-470.	0.6	30
93	Case-Control Study and Meta-Analysis Show a Weak Association between ANTXR2 Polymorphisms and Ankylosing Spondylitis in Chinese Han. <i>BioMed Research International</i> , 2018, 2018, 1-7.	0.9	1
94	Disabled-2 (DAB2) Overexpression Inhibits Monocyte-Derived Dendritic Cells' Function in Vogt-Koyanagi-Harada Disease. , 2018, 59, 4662.		10
95	Dynamic DNA Methylation Changes of <i>Tbx21</i> and <i>Rorc</i> during Experimental Autoimmune Uveitis in Mice. <i>Mediators of Inflammation</i> , 2018, 2018, 1-13.	1.4	16
96	Outcome and Prognostic Factors of Phacoemulsification Cataract Surgery in Vogt-Koyanagi-Harada Uveitis. <i>American Journal of Ophthalmology</i> , 2018, 196, 121-128.	1.7	17
97	Development and Evaluation of Diagnostic Criteria for Vogt-Koyanagi-Harada Disease. <i>JAMA Ophthalmology</i> , 2018, 136, 1025.	1.4	83
98	Association of <i>LACC1</i> , <i>CEBPB</i> , <i>PTPN1</i> , <i>RIPK2</i> and <i>ADO-EGR2</i> with ocular Behcet's disease in a Chinese Han population. <i>British Journal of Ophthalmology</i> , 2018, 102, 1308-1314.	2.1	16
99	Association of Long Noncoding RNAs Polymorphisms With Ankylosing Spondylitis, Vogt-Koyanagi-Harada Disease, and Behcet's Disease. , 2018, 59, 1158.		12
100	A metagenomic study of the gut microbiome in Behcet's disease. <i>Microbiome</i> , 2018, 6, 135.	4.9	173
101	Gut Microbiota Composition and Fecal Metabolic Phenotype in Patients With Acute Anterior Uveitis. , 2018, 59, 1523.		77
102	Aryl Hydrocarbon Receptor Regulates Apoptosis and Inflammation in a Murine Model of Experimental Autoimmune Uveitis. <i>Frontiers in Immunology</i> , 2018, 9, 1713.	2.2	43
103	Increased Expression of IL-23 Receptor (IL-23R) in Vogt-Koyanagi-Harada (VKH) Disease. <i>Current Eye Research</i> , 2018, 43, 1369-1373.	0.7	7
104	Hypermethylation of Interferon Regulatory Factor 8 (IRF8) Confers Risk to Vogt-Koyanagi-Harada Disease. <i>Scientific Reports</i> , 2017, 7, 1007.	1.6	23
105	Genetic Background of Uveitis in Chinese Population. <i>Essentials in Ophthalmology</i> , 2017, , 425-436.	0.0	0
106	Increased Complement 3a Receptor is Associated with Behcet's disease and Vogt-Koyanagi-Harada disease. <i>Scientific Reports</i> , 2017, 7, 15579.	1.6	10
107	Genetic polymorphisms of C-type lectin receptors in Behcet's disease in a Chinese Han population. <i>Scientific Reports</i> , 2017, 7, 5348.	1.6	9
108	Uveitis in Chinese Patients with Psoriasis. <i>Ocular Immunology and Inflammation</i> , 2017, 25, 855-865.	1.0	14

#	ARTICLE	IF	CITATIONS
109	Identification of susceptibility SNPs in IL10 and IL23R-IL12RB2 for Behçet's disease in Han Chinese. <i>Journal of Allergy and Clinical Immunology</i> , 2017, 139, 621-627.	1.5	36
110	miRNA Copy Number Variants Confer Susceptibility to Acute Anterior Uveitis With or Without Ankylosing Spondylitis. , 2017, 58, 1991.		11
111	Association of a PDCD1 Polymorphism With Sympathetic Ophthalmia in Han Chinese. , 2017, 58, 4218.		12
112	The Association of Chemokine Gene Polymorphisms with VKH and Behçet's Disease in a Chinese Han Population. <i>BioMed Research International</i> , 2017, 2017, 1-8.	0.9	16
113	Propofol inhibits lung cancer cell viability and induces cell apoptosis by upregulating microRNA-486 expression. <i>Brazilian Journal of Medical and Biological Research</i> , 2017, 50, e5794.	0.7	55
114	Promoter Hypermethylation of GATA3, IL-4, and TGF- $\beta$ 2 Confers Susceptibility to Vogt-Koyanagi-Harada Disease in Han Chinese. , 2017, 58, 1529.		18
115	Ocular Behçet's disease is associated with aberrant methylation of interferon regulatory factor 8 (IRF8) in monocyte-derived dendritic cells. <i>Oncotarget</i> , 2017, 8, 51277-51287.	0.8	9
116	Aberrant DNA methylation of GATA binding protein 3 (GATA3), interleukin-4 (IL-4), and transforming growth factor- $\beta$ 2 (TGF- $\beta$ 2) promoters in Behçet's disease. <i>Oncotarget</i> , 2017, 8, 64263-64272.	0.8	14
117	The Role of Mitochondria-Associated Reactive Oxygen Species in the Amyloid $\beta$ 2 Induced Production of Angiogenic Factors by ARPE-19 Cells. <i>Current Molecular Medicine</i> , 2017, 17, 140-148.	0.6	18
118	Association of T-Bet, GATA-3, RORC, and FOXP3 Copy Number Variations With Acute Anterior Uveitis With or Without Ankylosing Spondylitis in Chinese Han. , 2016, 57, 1847.		9
119	Decreased B and T lymphocyte attenuator in Behçet's disease may trigger abnormal Th17 and Th1 immune responses. <i>Scientific Reports</i> , 2016, 6, 20401.	1.6	26
120	Genetic polymorphisms of cell adhesion molecules in Behçet's disease in a Chinese Han population. <i>Scientific Reports</i> , 2016, 6, 24974.	1.6	21
121	Analysis of receptor tyrosine kinase genetics identifies two novel risk loci in GAS6 and PROS1 in Behçet's disease. <i>Scientific Reports</i> , 2016, 6, 26662.	1.6	10
122	Association of TNFSF4 Polymorphisms with Vogt-Koyanagi-Harada and Behçet's Disease in Han Chinese. <i>Scientific Reports</i> , 2016, 6, 37257.	1.6	16
123	miR-23a, miR-146a and miR-301a confer predisposition to Vogt-Koyanagi-Harada syndrome but not to Behçet's disease. <i>Scientific Reports</i> , 2016, 6, 20057.	1.6	22
124	Genetic analysis of innate immunity in Behçet's disease identifies an association with IL-37 and IL-18RAP. <i>Scientific Reports</i> , 2016, 6, 35802.	1.6	36
125	Two Genetic Variations in the IRF8 region are associated with Behçet's disease in Han Chinese. <i>Scientific Reports</i> , 2016, 6, 19651.	1.6	22
126	Genetic Variations of NLR family genes in Behçet's Disease. <i>Scientific Reports</i> , 2016, 6, 20098.	1.6	21



#	ARTICLE	IF	CITATIONS
127	Vogt-Koyanagi-Harada disease: Novel insights into pathophysiology, diagnosis and treatment. <i>Progress in Retinal and Eye Research</i> , 2016, 52, 84-111.	7.3	168
128	Investigation of the association of Vogt-Koyanagi-Harada syndrome with IL23R-C1orf141 in Han Chinese Singaporean and ADO-ZNF365-EGR2 in Thai. <i>British Journal of Ophthalmology</i> , 2016, 100, 436-442.	2.1	12
129	Higher Expression of NOD1 and NOD2 is Associated with Vogt-Koyanagi-Harada (VKH) Syndrome But Not Behcet's Disease (BD). <i>Current Molecular Medicine</i> , 2016, 16, 424-435.	0.6	13
130	Association of a NOS3 gene polymorphism with Behçet's disease but not with Vogt-Koyanagi-Harada syndrome in Han Chinese. <i>Molecular Vision</i> , 2016, 22, 311-8.	1.1	5
131	Copy Number Variants and Genetic Polymorphisms in TBX21, GATA3, Rorc, Foxp3 and Susceptibility to Behcet's Disease and Vogt-Koyanagi-Harada Syndrome. <i>Scientific Reports</i> , 2015, 5, 9511.	1.6	14
132	Copy number variations and gene polymorphisms of Complement components in ocular Behcet's disease and Vogt-Koyanagi-Harada syndrome. <i>Scientific Reports</i> , 2015, 5, 12989.	1.6	9
133	FASGene Copy Numbers are Associated with Susceptibility to Behçet Disease and VKH Syndrome in Han Chinese. <i>Human Mutation</i> , 2015, 36, 1064-1069.	1.1	15
134	Whole exome sequencing confirms the clinical diagnosis of Marfan syndrome combined with X-linked hypophosphatemia. <i>Journal of Translational Medicine</i> , 2015, 13, 179.	1.8	5
135	Association of ERAP1 Gene Polymorphisms With Behçet's Disease in Han Chinese. , 2015, 56, 6029.		22
136	Shock wave treatment enhances endothelial proliferation via autocrine vascular endothelial growth factor. <i>Genetics and Molecular Research</i> , 2015, 14, 19203-19210.	0.3	18
137	Association of ATG5 Gene Polymorphisms With Behçet's Disease and ATG10 Gene Polymorphisms With VKH Syndrome in a Chinese Han Population. , 2015, 56, 8280.		24
138	Editorial (Thematic Issue: Uveitis: Pathology, Molecular Mechanisms and Therapy). <i>Current Molecular Medicine</i> , 2015, 15, 510-510.	0.6	4
139	Genetic Variations of IL17F and IL23A Show Associations with Behçet's Disease and Vogt-Koyanagi-Harada Syndrome. <i>Ophthalmology</i> , 2015, 122, 518-523.	2.5	40
140	Association Between Copy Number Variations of TLR7 and Ocular Behcet's Disease in a Chinese Han Population. <i>Investigative Ophthalmology and Visual Science</i> , 2015, 56, 1517-1523.	3.3	18
141	A variant of CLEC16A gene confers protection for Vogt-Koyanagi-Harada syndrome but not for Behcet's disease in a Chinese Han population. <i>Experimental Eye Research</i> , 2015, 132, 225-230.	1.2	6
142	No association between Bach2 gene polymorphisms with Vogt-Koyanagi-Harada syndrome (VKH) and Behcet's disease (BD) in a Chinese Han population. <i>British Journal of Ophthalmology</i> , 2015, 99, 1150-1154.	2.1	1
143	Molecular Genetic Advances in Uveitis. <i>Progress in Molecular Biology and Translational Science</i> , 2015, 134, 283-298.	0.9	28
144	Decreased Interleukin-37 Expression in Vogt-Koyanagi-Harada Disease and Upregulation Following Immunosuppressive Treatment. <i>Journal of Interferon and Cytokine Research</i> , 2015, 35, 265-272.	0.5	19

#	ARTICLE	IF	CITATIONS
145	TLR3 and TLR4 But not TLR2 are Involved in Vogt-Koyanagi- Harada Disease by Triggering Proinflammatory Cytokines Production Through Promoting the Production of Mitochondrial Reactive Oxygen Species. <i>Current Molecular Medicine</i> , 2015, 15, 529-542.	0.6	15
146	Interleukin-10 gene polymorphisms are associated with Behcet's disease but not with Vogt-Koyanagi-Harada syndrome in the Chinese Han population. <i>Molecular Vision</i> , 2015, 21, 589-603.	1.1	25
147	TRAF5 and TRAF3IP2 Gene Polymorphisms Are Associated with Behçet's Disease and Vogt-Koyanagi-Harada Syndrome: A Case-Control Study. <i>PLoS ONE</i> , 2014, 9, e84214.	1.1	22
148	The Role of Interleukin-1 Receptor-Associated Kinases in Vogt-Koyanagi-Harada Disease. <i>PLoS ONE</i> , 2014, 9, e93214.	1.1	11
149	Association of a TNIP1 Polymorphism with Vogt-Koyanagi-Harada Syndrome but Not with Ocular Behçet's Disease in Han Chinese. <i>PLoS ONE</i> , 2014, 9, e95573.	1.1	7
150	A Functional Variant of PTPN22 Confers Risk for Vogt-Koyanagi-Harada Syndrome but Not for Ankylosing Spondylitis. <i>PLoS ONE</i> , 2014, 9, e96943.	1.1	19
151	Genetic Variations of IL-12B, IL-12RÎ1, IL-12RÎ2 in Behcet's Disease and VKH Syndrome. <i>PLoS ONE</i> , 2014, 9, e98373.	1.1	21
152	Activation of Liver X Receptor Alleviates Ocular Inflammation in Experimental Autoimmune Uveitis. , 2014, 55, 2795.		40
153	Inhibition of Proinflammatory Cytokine by IL-25 in Vogt-Koyanagi-Harada Syndrome. <i>Ocular Immunology and Inflammation</i> , 2014, 22, 294-299.	1.0	8
154	FoxO1 Gene Confers Genetic Predisposition to Acute Anterior Uveitis With Ankylosing Spondylitis. <i>Investigative Ophthalmology and Visual Science</i> , 2014, 55, 7970-7974.	3.3	29
155	High C4 gene copy numbers protects against Vogt-Koyanagi-Harada syndrome in Chinese Han. <i>British Journal of Ophthalmology</i> , 2014, 98, 1733-1737.	2.1	21
156	Activation of the aryl hydrocarbon receptor affects activation and function of human monocyte-derived dendritic cells. <i>Clinical and Experimental Immunology</i> , 2014, 177, 521-530.	1.1	66
157	<i>MicroRNA-146a</i> and <i>Ets-1</i> gene polymorphisms in ocular Behçet's disease and Vogt-Koyanagi-Harada syndrome. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 170-176.	0.5	77
158	Increased Notch pathway activation in Behçet's disease. <i>Rheumatology</i> , 2014, 53, 810-820.	0.9	33
159	Polymorphisms in Genetics of Vitamin D Metabolism Confer Susceptibility to Ocular Behçet Disease in a Chinese Han Population. <i>American Journal of Ophthalmology</i> , 2014, 157, 488-494.e6.	1.7	18
160	Genome-wide association analysis of Vogt-Koyanagi-Harada syndrome identifies two new susceptibility loci at 1p31.2 and 10q21.3. <i>Nature Genetics</i> , 2014, 46, 1007-1011.	9.4	88
161	Decreased interleukin 27 expression is associated with active uveitis in Behçet's disease. <i>Arthritis Research and Therapy</i> , 2014, 16, R117.	1.6	36
162	Predisposition to Behçet's disease and VKH syndrome by genetic variants of miR-182. <i>Journal of Molecular Medicine</i> , 2014, 92, 961-967.	1.7	56

#	ARTICLE	IF	CITATIONS
163	MicroRNA-146a and Ets-1 Gene Polymorphisms Are Associated with Pediatric Uveitis. <i>PLoS ONE</i> , 2014, 9, e91199.	1.1	28
164	A Possible Role for Interleukin 37 in the Pathogenesis of Behcet's Disease. <i>Current Molecular Medicine</i> , 2014, 14, 535-542.	0.6	53
165	Activation of the Liver X Receptor Inhibits Th17 and Th1 Responses in Behcet's Disease and Vogt-Koyanagi-Harada Disease. <i>Current Molecular Medicine</i> , 2014, 14, 712-722.	0.6	11
166	Genetic variants in the JAK1 gene confer higher risk of Behcet's disease with ocular involvement in Han Chinese. <i>Human Genetics</i> , 2013, 132, 1049-1058.	1.8	26
167	TNFAIP3 gene polymorphisms confer risk for Behcet's disease in a Chinese Han population. <i>Human Genetics</i> , 2013, 132, 293-300.	1.8	53
168	Copy Number Variations of Complement Component C4 Are Associated With Behcet's Disease but Not With Ankylosing Spondylitis Associated With Acute Anterior Uveitis. <i>Arthritis and Rheumatism</i> , 2013, 65, 2963-2970.	6.7	34
169	TNF receptor-associated factor 5 gene confers genetic predisposition to acute anterior uveitis and pediatric uveitis. <i>Arthritis Research and Therapy</i> , 2013, 15, R113.	1.6	32
170	A functional variant of pre-miRNA-196a2 confers risk for Behcet's disease but not for Vogt-Koyanagi-Harada syndrome or AAU in ankylosing spondylitis. <i>Human Genetics</i> , 2013, 132, 1395-1404.	1.8	50
171	IL-1 $\beta$ Triggered by Peptidoglycan and Lipopolysaccharide through TLR2/4 and ROS-NLRP3 Inflammasome-Dependent Pathways Is Involved in Ocular Behcet's Disease. , 2013, 54, 402.		67
172	Genetic Variant on PDGFRL Associated with Behcet Disease in Chinese Han Populations. <i>Human Mutation</i> , 2013, 34, 74-78.	1.1	15
173	MIF Gene Polymorphisms Confer Susceptibility to Vogt-Koyanagi-Harada Syndrome in a Han Chinese Population. , 2013, 54, 7734.		21
174	Association of TLR2 Gene Polymorphisms With Ocular Behcet's Disease in a Chinese Han Population. , 2013, 54, 8384.		39
175	Berberine Suppresses Th17 and Dendritic Cell Responses. , 2013, 54, 2516.		39
176	Reply. <i>Retina</i> , 2013, 33, 451-453.	1.0	0
177	Higher Expression of Toll-like Receptors 2, 3, 4, and 8 in Ocular Behcet's Disease. , 2013, 54, 6012.		40
178	JAK1, but Not JAK2 and STAT3, Confers Susceptibility to Vogt-Koyanagi-Harada (VKH) Syndrome in a Han Chinese Population. , 2013, 54, 3360.		21
179	Increased Expression of IL-22 Is Associated with Disease Activity in Behcet's Disease. <i>PLoS ONE</i> , 2013, 8, e59009.	1.1	27
180	FGFR1OP tagSNP but Not CCR6 Polymorphisms Are Associated with Vogt-Koyanagi-Harada Syndrome in Chinese Han. <i>PLoS ONE</i> , 2013, 8, e69358.	1.1	9

#	ARTICLE	IF	CITATIONS
181	TNFAIP3 Gene Polymorphisms in a Chinese Han Population with Vogt-Koyanagi-Harada Syndrome. PLoS ONE, 2013, 8, e59515.	1.1	11
182	CD40 gene polymorphisms confer risk of Behcet's disease but not of Vogt-Koyanagi-Harada syndrome in a Han Chinese population. Rheumatology, 2012, 51, 47-51.	0.9	33
183	JAK2 and STAT3 Polymorphisms in a Han Chinese Population with Behçet's Disease. , 2012, 53, 538.		40
184	Association Analysis of TGFBR3 Gene with Vogt-Koyanagi-Harada Disease and Behçet's Disease in the Chinese Han Population. Current Eye Research, 2012, 37, 312-317.	0.7	20
185	Effect of 1,25-Dihydroxyvitamin D3 on Th17 and Th1 Response in Patients with Behçet's Disease. , 2012, 53, 6434.		45
186	OCULAR MANIFESTATIONS OF SYPHILITIC UVEITIS IN CHINESE PATIENTS. Retina, 2012, 32, 1906-1914.	1.0	69
187	Identification of a susceptibility locus in STAT4 for Behçet's disease in Han Chinese in a genome-wide association study. Arthritis and Rheumatism, 2012, 64, 4104-4113.	6.7	163
188	Two-stage association study in Chinese Han identifies two independent associations in CCR1/CCR3 locus as candidate for Behçet's disease susceptibility. Human Genetics, 2012, 131, 1841-1850.	1.8	46
189	Association of Macrophage Migration Inhibitory Factor Gene Polymorphisms with Behçet's Disease in a Han Chinese Population. Ophthalmology, 2012, 119, 2514-2518.	2.5	23
190	The genetics of Behçet's disease in a Chinese population. Frontiers of Medicine, 2012, 6, 354-359.	1.5	20
191	Replication study confirms the association between UBAC2 and Behçet's disease in two independent Chinese sets of patients and controls. Arthritis Research and Therapy, 2012, 14, R70.	1.6	37
192	AAV2-Mediated Subretinal Gene Transfer of mIL-27p28 Attenuates Experimental Autoimmune Uveoretinitis in Mice. PLoS ONE, 2012, 7, e37773.	1.1	16
193	AAV2-Mediated Combined Subretinal Delivery of IFN- $\gamma$ and IL-4 Reduces the Severity of Experimental Autoimmune Uveoretinitis. PLoS ONE, 2012, 7, e37995.	1.1	19
194	IFN- $\gamma$ Inhibits the Increased Expression of IL-9 during Experimental Autoimmune Uveoretinitis. PLoS ONE, 2012, 7, e48566.	1.1	9
195	Increased IL-7 Expression in Vogt-Koyanagi-Harada Disease. , 2012, 53, 1012.		24
196	Decreased microRNA-155 Expression in Ocular Behçet's Disease but Not in Vogt Koyanagi Harada Syndrome. , 2012, 53, 5665.		69
197	Decreased IL-27 Expression in Association with an Increased Th17 Response in Vogt-Koyanagi-Harada Disease. , 2012, 53, 4668.		62
198	Alterations of color vision and central visual field in patients with Vogt-Koyanagi-Harada syndrome. Journal of Ophthalmic Inflammation and Infection, 2012, 2, 75-79.	1.2	7

#	ARTICLE	IF	CITATIONS
199	No Association of PTPN22 Polymorphisms with Susceptibility to Ocular Behcet's Disease in Two Chinese Han Populations. <i>PLoS ONE</i> , 2012, 7, e31230.	1.1	12
200	Lack of association of miR-146a and Ets-1 gene polymorphisms with Fuchs uveitis syndrome in Chinese Han patients. <i>Molecular Vision</i> , 2012, 18, 426-30.	1.1	8
201	IFN- $\gamma$ blocks IL-17 production by peripheral blood mononuclear cells in Behcet's disease. <i>Rheumatology</i> , 2011, 50, 293-298.	0.9	38
202	Behcet's disease exhibits an increased osteopontin serum level in active stage but no association with osteopontin and its receptor gene polymorphisms. <i>Human Immunology</i> , 2011, 72, 525-529.	1.2	13
203	Elevated Serum Osteopontin Levels and Genetic Polymorphisms of Osteopontin Are Associated with Vogt-Koyanagi-Harada Disease. , 2011, 52, 7084.		22
204	Label-Free Proteomics Reveals Decreased Expression of CD18 and AKNA in Peripheral CD4+ T Cells from Patients with Vogt-Koyanagi-Harada Syndrome. <i>PLoS ONE</i> , 2011, 6, e14616.	1.1	20
205	The Effects of Th17 Cytokines on the Inflammatory Mediator Production and Barrier Function of ARPE-19 Cells. <i>PLoS ONE</i> , 2011, 6, e18139.	1.1	77
206	Regulatory Effects of IFN- $\gamma$ on the Development of Experimental Autoimmune Uveoretinitis in B10R/III Mice. <i>PLoS ONE</i> , 2011, 6, e19870.	1.1	15
207	Lack of an Association of PD-1 and Its Ligand Genes with Behcet's Disease in a Chinese Han Population. <i>PLoS ONE</i> , 2011, 6, e25345.	1.1	13
208	Band-Shaped Keratopathy in Chinese Patients With Vogt-Koyanagi-Harada Syndrome. <i>Cornea</i> , 2011, 30, 1336-1340.	0.9	13
209	Vogt-Koyanagi-Harada disease presenting as acute angle closure glaucoma at onset. <i>Clinical and Experimental Ophthalmology</i> , 2011, 39, 639-647.	1.3	34
210	There is no association of CCR6 polymorphisms with susceptibility to Behcet's disease in two Chinese Han populations. <i>British Journal of Ophthalmology</i> , 2011, 95, 1603-1606.	2.1	3
211	AAV2-Mediated Subretinal Gene Transfer of hIFN- $\gamma$ Attenuates Experimental Autoimmune Uveoretinitis in Mice. <i>PLoS ONE</i> , 2011, 6, e19542.	1.1	25
212	CD4 <sup>+</sup> T cells from behcet patients produce high levels of IL-17. <i>Yan Ke Xue Bao = Eye Science</i> , 2011, 26, 65-9.	0.1	8
213	Decreased 1,25-Dihydroxyvitamin D3 level is involved in the pathogenesis of Vogt-Koyanagi-Harada (VKH) disease. <i>Molecular Vision</i> , 2011, 17, 673-9.	1.1	22
214	IL-17A stimulates the production of inflammatory mediators via Erk1/2, p38 MAPK, PI3K/Akt, and NF- $\kappa$ B pathways in ARPE-19 cells. <i>Molecular Vision</i> , 2011, 17, 3072-7.	1.1	34
215	Contribution of CD4 <sup>+</sup> CD25 <sup>+</sup> T Cells to the Regression Phase of Experimental Autoimmune Uveoretinitis. , 2010, 51, 383.		58
216	Upregulation of Interleukin 21 and Promotion of Interleukin 17 Production in Chronic or Recurrent Vogt-Koyanagi-Harada Disease. <i>JAMA Ophthalmology</i> , 2010, 128, 1449.	2.6	30

#	ARTICLE	IF	CITATIONS
217	Increased Regulatory T Cells in Spleen during Experimental Autoimmune Uveoretinitis. <i>Ocular Immunology and Inflammation</i> , 2010, 18, 38-43.	1.0	15
218	IL-23R gene confers susceptibility to Behcet's disease in a Chinese Han population. <i>Annals of the Rheumatic Diseases</i> , 2010, 69, 1325-1328.	0.5	91
219	Monocyte chemoattractant protein-1 $\alpha$ 2518 A/G single nucleotide polymorphism in Chinese Han patients with ocular Behcet's disease. <i>Human Immunology</i> , 2010, 71, 79-82.	1.2	22
220	Polymorphisms of IL23R and Vogt-Koyanagi-Harada syndrome in a Chinese Han population. <i>Human Immunology</i> , 2010, 71, 414-417.	1.2	19
221	STAT4 polymorphism in a Chinese Han population with Vogt-Koyanagi-Harada syndrome and Behcet's disease. <i>Human Immunology</i> , 2010, 71, 723-726.	1.2	41
222	Interleukin-17 gene polymorphism is associated with Vogt-Koyanagi-Harada syndrome but not with Behcet's disease in a Chinese Han population. <i>Human Immunology</i> , 2010, 71, 988-991.	1.2	58
223	Production of interleukin-17 in Behcet's disease is inhibited by cyclosporin A. <i>Molecular Vision</i> , 2010, 16, 880-6.	1.1	35
224	Inhibitory effect of rapamycin and dexamethasone on production of IL-17 and IFN- $\gamma$ in Vogt-Koyanagi-Harada patients. <i>British Journal of Ophthalmology</i> , 2009, 93, 249-253.	2.1	22
225	No association of CTLA-4 polymorphisms with susceptibility to Behcet disease. <i>British Journal of Ophthalmology</i> , 2009, 93, 1378-1381.	2.1	13
226	Inhibitory effect of Cyclosporin A and corticosteroids on the production of IFN- $\gamma$ and IL-17 by T cells in Vogt-Koyanagi-Harada syndrome. <i>Clinical Immunology</i> , 2009, 131, 333-342.	1.4	73
227	Immune Response Genes in Uveitis. <i>Ocular Immunology and Inflammation</i> , 2009, 17, 249-256.	1.0	57
228	PDCD1 genes may protect against extraocular manifestations in Chinese Han patients with Vogt-Koyanagi-Harada syndrome. <i>Molecular Vision</i> , 2009, 15, 386-92.	1.1	32
229	Polymorphisms of FCRL3 in a Chinese population with Vogt-Koyanagi-Harada (VKH) syndrome. <i>Molecular Vision</i> , 2009, 15, 955-61.	1.1	10
230	Lack of association of two polymorphisms of IRF5 with Behcet's disease. <i>Molecular Vision</i> , 2009, 15, 2018-21.	1.1	4
231	Association of the CTLA-4 gene with Vogt-Koyanagi-Harada syndrome. <i>Clinical Immunology</i> , 2008, 127, 43-48.	1.4	50
232	Vogt-Koyanagi-Harada Syndrome. <i>Current Eye Research</i> , 2008, 33, 517-523.	0.7	114
233	Study of Macular Function by Multifocal Electroretinography in Patients With Vogt-Koyanagi-Harada Syndrome. <i>American Journal of Ophthalmology</i> , 2008, 146, 767-771.e2.	1.7	36
234	Clinical Features of Chinese Patients with Behcet's Disease. <i>Ophthalmology</i> , 2008, 115, 312-318.e4.	2.5	175

#	ARTICLE	IF	CITATIONS
235	SUMO4 gene polymorphisms in Chinese Han patients with Behcet's disease. <i>Clinical Immunology</i> , 2008, 129, 170-175.	1.4	63
236	Longitudinal quantification of aqueous flare and cells in Vogt-Koyanagi-Harada disease. <i>British Journal of Ophthalmology</i> , 2008, 92, 182-185.	2.1	44
237	Leptin increases in Vogt-Koyanagi-Harada (VKH) disease and promotes cell proliferation and inflammatory cytokine secretion. <i>British Journal of Ophthalmology</i> , 2008, 92, 557-561.	2.1	24
238	Diminished Frequency and Function of CD4+CD25highRegulatory T Cells Associated with Active Uveitis in Vogt-Koyanagi-Harada Syndrome. , 2008, 49, 3475.		109
239	Upregulated IL-23 and IL-17 in Behçet Patients with Active Uveitis. , 2008, 49, 3058.		296
240	Association between polymorphisms of FCRL3, a non-HLA gene, and Behçet's disease in a Chinese population with ophthalmic manifestations. <i>Molecular Vision</i> , 2008, 14, 2136-42.	1.1	27
241	Small ubiquitin-like modifier 4 (SUMO4) polymorphisms and Vogt-Koyanagi-Harada (VKH) syndrome in the Chinese Han population. <i>Molecular Vision</i> , 2008, 14, 2597-603.	1.1	20
242	Clinical Characteristics of Vogt-Koyanagi-Harada Syndrome in Chinese Patients. <i>Ophthalmology</i> , 2007, 114, 606-614.e3.	2.5	257
243	IL-23 promotes CD4+ T cells to produce IL-17 in Vogt-Koyanagi-Harada disease. <i>Journal of Allergy and Clinical Immunology</i> , 2007, 119, 1218-1224.	1.5	190
244	T-bet expression in the iris and spleen parallels disease expression during endotoxin-induced uveitis. <i>Graefe's Archive for Clinical and Experimental Ophthalmology</i> , 2007, 245, 407-413.	1.0	7
245	Phenotypes, distribution, and morphological features of antigen-presenting cells in the murine cornea following intravitreal injection. <i>Molecular Vision</i> , 2007, 13, 475-86.	1.1	4
246	Clinical Features of Chinese Patients with Fuchs' Syndrome. <i>Ophthalmology</i> , 2006, 113, 473-480.	2.5	74
247	Indoleamine 2,3-dioxygenase (IDO) is involved in promoting the development of anterior chamber-associated immune deviation. <i>Immunology Letters</i> , 2006, 107, 140-147.	1.1	21
248	Upregulation of T-bet expression in peripheral blood mononuclear cells during Vogt-Koyanagi-Harada disease. <i>British Journal of Ophthalmology</i> , 2005, 89, 1410-1412.	2.1	48
249	Clinical Patterns and Characteristics of Uveitis in a Tertiary Center for Uveitis in China. <i>Current Eye Research</i> , 2005, 30, 943-948.	0.7	307
250	Macrophages and MHC class II positive dendritiform cells in the iris and choroid of the pig. <i>Current Eye Research</i> , 2003, 26, 291-296.	0.7	16
251	T-bet expression is upregulated in active Behcet's disease. <i>British Journal of Ophthalmology</i> , 2003, 87, 1264-1267.	2.1	42
252	Resistance of lymphocytes to Fas-mediated apoptosis in Behçet's disease and Vogt-Koyangi-Harada syndrome. <i>Ocular Immunology and Inflammation</i> , 2002, 10, 47-52.	1.0	30



#	ARTICLE	IF	CITATIONS
253	Immune cells in the porcine retina: distribution, characterization and morphological features. <i>Investigative Ophthalmology and Visual Science</i> , 2002, 43, 1488-92.	3.3	29
254	Disturbed expression of Fas/FasL on CD4+ and CD8+T cells in Behcet's disease, Vogt-Koyanagi-Harada syndrome, and idiopathic anterior uveitis. <i>Ocular Immunology and Inflammation</i> , 2001, 9, 185-191.	1.0	15
255	Localization and characterization of immunocompetent cells in the human retina. <i>Ocular Immunology and Inflammation</i> , 2000, 8, 149-157.	1.0	34
256	Localization and characterization of immunocompetent cells in the human retina. <i>Ocular Immunology and Inflammation</i> , 2000, 8, 149-57.	1.0	9
257	A case report of frosted branch angiitis and its visual electrophysiology. <i>Documenta Ophthalmologica</i> , 1998, 97, 135-142.	1.0	9
258	Immunohistochemical studies on the endotoxin-induced uveitis. <i>Chinese Medical Journal</i> , 1998, 111, 252-6.	0.9	2
259	Genetics in Behcet's Disease: An Update Review. <i>Frontiers in Ophthalmology</i> , 0, 2, .	0.2	0