

Stephen M Twigg

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3679704/stephen-m-twigg-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

137
papers

4,358
citations

36
h-index

63
g-index

141
ext. papers

5,000
ext. citations

5.6
avg. IF

5.28
L-index

#	Paper	IF	Citations
137	Sitagliptin Is More Effective Than Gliclazide in Preventing Pro-Fibrotic and Pro-Inflammatory Changes in a Rodent Model of Diet-Induced Non-Alcoholic Fatty Liver Disease.. <i>Molecules</i> , 2022 , 27,	4.8	1
136	Targeting CCN2 protects against progressive non-alcoholic steatohepatitis in a preclinical model induced by high-fat feeding and type 2 diabetes.. <i>Journal of Cell Communication and Signaling</i> , 2022 , 1	5.2	1
135	Alterations of CD163 expression in the complications of diabetes: A systematic review.. <i>Journal of Diabetes and Its Complications</i> , 2022 , 108150	3.2	0
134	Guidelines development protocol and findings: part of the 2021 Australian evidence-based guidelines for diabetes-related foot disease.. <i>Journal of Foot and Ankle Research</i> , 2022 , 15, 28	3.2	1
133	The effect of acute aerobic exercise on central arterial stiffness, wave reflections, and hemodynamics in adults with diabetes: A randomized cross-over design. <i>Journal of Sport and Health Science</i> , 2021 , 10, 499-506	8.2	5
132	Australian guideline on wound classification of diabetes-related foot ulcers: part of the 2021 Australian evidence-based guidelines for diabetes-related foot disease. <i>Journal of Foot and Ankle Research</i> , 2021 , 14, 60	3.2	2
131	A Randomized Trial Comparing Weekly With Every Second Week Sharp Debridement in People With Diabetes-Related Foot Ulcers Shows Similar Healing Outcomes: Potential Benefit to Resource Utilization. <i>Diabetes Care</i> , 2021 , 44, e203-e205	14.6	
130	Managing arterial health in adults with metabolic diseases: Is high-intensity interval exercise the answer? Response to the commentary by Lopes et al. <i>Journal of Sport and Health Science</i> , 2021 , 10, 510-512	8.2	
129	Impact of adiposity on clinical outcomes in people living with a Fontan circulation. <i>International Journal of Cardiology</i> , 2021 , 329, 82-88	3.2	4
128	The association between cardiorespiratory fitness, liver fat and insulin resistance in adults with or without type 2 diabetes: a cross-sectional analysis. <i>BMC Sports Science, Medicine and Rehabilitation</i> , 2021 , 13, 40	2.4	1
127	Monocyte phenotype as a predictive marker for wound healing in diabetes-related foot ulcers. <i>Journal of Diabetes and Its Complications</i> , 2021 , 35, 107889	3.2	2
126	Prevalence, causes and associated mortality of hypercalcaemia in modern hospital care. <i>Internal Medicine Journal</i> , 2021 ,	1.6	1
125	Degree of adiposity and obesity severity is associated with cutaneous microvascular dysfunction in type 2 diabetes. <i>Microvascular Research</i> , 2021 , 136, 104149	3.7	0
124	Estimating the diagnostic accuracy of the ankle-brachial pressure index for detecting peripheral arterial disease in people with diabetes: A systematic review and meta-analysis. <i>Diabetic Medicine</i> , 2021 , 38, e14379	3.5	3
123	Metabolic syndrome in type 1 diabetes and its association with diabetes complications. <i>Diabetic Medicine</i> , 2021 , 38, e14376	3.5	3
122	U-Shaped Relationship of Leukocyte Telomere Length With All-Cause and Cancer-Related Mortality in Older Men. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2021 , 76, 164-171	6.4	2
121	Effect of exercise on hepatic steatosis: Are benefits seen without dietary intervention? A systematic review and meta-analysis. <i>Journal of Diabetes</i> , 2021 , 13, 63-77	3.8	10

120	An Enhanced SMS Text Message-Based Support and Reminder Program for Young Adults With Type 2 Diabetes (TEXT2U): Randomized Controlled Trial. <i>Journal of Medical Internet Research</i> , 2021 , 23, e27263	7.6	0
119	Exercise induces favorable metabolic changes in white adipose tissue preventing high-fat diet obesity. <i>Physiological Reports</i> , 2021 , 9, e14929	2.6	1
118	Frequency of sharp wound debridement in the management of diabetes-related foot ulcers: exploring current practice. <i>Journal of Foot and Ankle Research</i> , 2021 , 14, 52	3.2	0
117	Young adult onset type 2 diabetes versus type 1 diabetes: Progression to and survival on renal replacement therapy. <i>Journal of Diabetes and Its Complications</i> , 2021 , 35, 108023	3.2	1
116	The Effect of a Sustained High-Fat Diet on the Metabolism of White and Brown Adipose Tissue and Its Impact on Insulin Resistance: A Selected Time Point Cross-Sectional Study.. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	3
115	The association of periodontal disease with the complications of diabetes mellitus. A systematic review. <i>Diabetes Research and Clinical Practice</i> , 2020 , 165, 108244	7.4	19
114	The effect of low-volume high-intensity interval training on cardiovascular health outcomes in type 2 diabetes: A randomised controlled trial. <i>International Journal of Cardiology</i> , 2020 , 320, 148-154	3.2	17
113	Effect of High-Intensity Interval Training on Glycemic Control in Adults With Type 1 Diabetes and Overweight or Obesity: A Randomized Controlled Trial With Partial Crossover. <i>Diabetes Care</i> , 2020 , 43, 2281-2288	14.6	6
112	Young-onset type 2 diabetes and younger current age: increased susceptibility to retinopathy in contrast to other complications. <i>Diabetic Medicine</i> , 2020 , 37, 991-999	3.5	11
111	Associations of plasma IGF1, IGFBP3 and estradiol with leucocyte telomere length, a marker of biological age, in men. <i>European Journal of Endocrinology</i> , 2020 , 182, 23-33	6.5	5
110	Secular Trends in Information Communications Technology: Access, Use, and Attitudes of Young and Older Patients With Diabetes. <i>Diabetes Spectrum</i> , 2020 , 33, 66-73	1.9	2
109	The effect of TGF β 1 on thermogenic markers is dependent on the degree of adipocyte differentiation. <i>Bioscience Reports</i> , 2020 , 40,	4.1	2
108	Skeletal muscle adiponectin induction in obesity and exercise. <i>Metabolism: Clinical and Experimental</i> , 2020 , 102, 154008	12.7	34
107	Apolipoprotein-AI mimetic peptides D-4F and L-5F decrease hepatic inflammation and increase insulin sensitivity in C57BL/6 mice. <i>PLoS ONE</i> , 2020 , 15, e0226931	3.7	7
106	High-intensity interval exercise and hypoglycaemia minimisation in adults with type 1 diabetes: A randomised cross-over trial. <i>Journal of Diabetes and Its Complications</i> , 2020 , 34, 107514	3.2	6
105	The Effect of a Novel Low-Volume Aerobic Exercise Intervention on Liver Fat in Type 2 Diabetes: A Randomized Controlled Trial. <i>Diabetes Care</i> , 2020 , 43, 2371-2378	14.6	14
104	Constant-moderate versus high-intensity interval training on heart adiponectin levels in high-fat fed mice: a preventive and treatment approach. <i>Archives of Physiology and Biochemistry</i> , 2020 , 1-5	2.2	0
103	Blockade of High-Fat Diet Proteomic Phenotypes Using Exercise as Prevention or Treatment. <i>Molecular and Cellular Proteomics</i> , 2020 , 20, 100027	7.6	2

102	Apolipoprotein-AI mimetic peptides D-4F and L-5F decrease hepatic inflammation and increase insulin sensitivity in C57BL/6 mice 2020 , 15, e0226931		
101	Apolipoprotein-AI mimetic peptides D-4F and L-5F decrease hepatic inflammation and increase insulin sensitivity in C57BL/6 mice 2020 , 15, e0226931		
100	Apolipoprotein-AI mimetic peptides D-4F and L-5F decrease hepatic inflammation and increase insulin sensitivity in C57BL/6 mice 2020 , 15, e0226931		
99	Apolipoprotein-AI mimetic peptides D-4F and L-5F decrease hepatic inflammation and increase insulin sensitivity in C57BL/6 mice 2020 , 15, e0226931		
98	Constant-Moderate and High-Intensity Interval Training Have Differential Benefits on Insulin Sensitive Tissues in High-Fat Fed Mice. <i>Frontiers in Physiology</i> , 2019 , 10, 459	4.6	14
97	Prevalence and risk factors for low bone density in adults with a Fontan circulation. <i>Congenital Heart Disease</i> , 2019 , 14, 987-995	3.1	6
96	An on-line support tool to reduce exercise-related hypoglycaemia and improve confidence to exercise in type 1 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2019 , 33, 682-689	3.2	2
95	OR18-2 Higher Plasma Estradiol Concentration Is Independently Associated with Lower Biological Age, Assessed as Leucocyte Telomere Length, in Older Men. <i>Journal of the Endocrine Society</i> , 2019 , 3,	0.4	78
94	Opioid-induced hypoadrenalism resulting in fasting hypoglycaemia. <i>BMJ Case Reports</i> , 2019 , 12,	0.9	3
93	Cross-sectional associations of sex hormones with leucocyte telomere length, a marker of biological age, in a community-based cohort of older men. <i>Clinical Endocrinology</i> , 2019 , 90, 562-569	3.4	4
92	Contrasting effects of IGF binding protein-3 expression in mammary tumor cells and the tumor microenvironment. <i>Experimental Cell Research</i> , 2019 , 374, 38-45	4.2	5
91	Self-reported physical activity in community-dwelling adults with diabetes and its association with diabetes complications. <i>Journal of Diabetes and Its Complications</i> , 2019 , 33, 33-38	3.2	8
90	Identification of Novel Natural Substrates of Fibroblast Activation Protein-alpha by Differential Degradomics and Proteomics. <i>Molecular and Cellular Proteomics</i> , 2019 , 18, 65-85	7.6	18
89	Differential metabolic effects of constant moderate versus high intensity interval training in high-fat fed mice: possible role of muscle adiponectin. <i>Physiological Reports</i> , 2018 , 6, e13599	2.6	23
88	Regulation and bioactivity of the CCN family of genes and proteins in obesity and diabetes. <i>Journal of Cell Communication and Signaling</i> , 2018 , 12, 359-368	5.2	10
87	Differing clinical phenotype for higher alanine-aminotransferase (ALT) compared with high-risk NAFLD fibrosis score in type 2 diabetes mellitus. <i>Journal of Diabetes and Its Complications</i> , 2018 , 32, 321-324	3.2	5
86	Enhancement of mammary tumour growth by IGFBP-3 involves impaired T cell accumulation. <i>Endocrine-Related Cancer</i> , 2018 , 25, 111-122	5.7	9
85	Skeletal muscle adiponectin induction depends on diet, muscle type/activity, and exercise modality in C57BL/6 mice. <i>Physiological Reports</i> , 2018 , 6, e13848	2.6	5

84	Association of Patient Profile with Glycemic Control and Hypoglycemia with Insulin Glargine 300 U/mL in Type 2 Diabetes: A Post Hoc Patient-Level Meta-Analysis. <i>Diabetes Therapy</i> , 2018 , 9, 2043-2053	3.6	6
83	Cardiac Effects of Sulfonylurea-Related Hypoglycemia. <i>Diabetes Care</i> , 2017 , 40, 663-670	14.6	21
82	Data collection on retinopathy as a public health tool: The Hubble telescope equivalent of looking back in time. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 721-725	3.2	2
81	Utility and reliability of non-invasive muscle function tests in high-fat-fed mice. <i>Experimental Physiology</i> , 2017 , 102, 773-778	2.4	8
80	The emerging role of skeletal muscle extracellular matrix remodelling in obesity and exercise. <i>Obesity Reviews</i> , 2017 , 18, 776-790	10.6	26
79	Post-occlusive reactive hyperaemia of skin microvasculature and foot complications in type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 1305-1310	3.2	12
78	Insulin treatment prevents wounding associated changes in tissue and circulating neutrophil MMP-9 and NGAL in diabetic rats. <i>PLoS ONE</i> , 2017 , 12, e0170951	3.7	11
77	Method for Analysis of Matrix Degradation by CCN2 Through the MMP/TIMP System. <i>Methods in Molecular Biology</i> , 2017 , 1489, 523-532	1.4	
76	Non-invasive lower limb small arterial measures co-segregate strongly with foot complications in people with diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 589-593	3.2	2
75	Changing trends for diagnostic methods in Graves disease in Australia: an immunological diagnosis as the emerging preference. <i>Internal Medicine Journal</i> , 2017 , 47, 1464-1465	1.6	
74	Painful ovulation in a 46,XX SRY -ve adult male with duplication. <i>Endocrinology, Diabetes and Metabolism Case Reports</i> , 2017 , 2017,	1.4	4
73	Low alanine aminotransferase levels and higher number of cardiovascular events in people with Type 2 diabetes: analysis of the Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) study. <i>Diabetic Medicine</i> , 2016 , 33, 356-64	3.5	3
72	Hypoglycaemia and QT interval prolongation: Detection by simultaneous Holter and continuous glucose monitoring. <i>Diabetes Research and Clinical Practice</i> , 2016 , 113, 211-4	7.4	16
71	Opposite associations between alanine aminotransferase and γ -glutamyl transferase levels and all-cause mortality in type 2 diabetes: Analysis of the Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) study. <i>Metabolism: Clinical and Experimental</i> , 2016 , 65, 783-793	12.7	13
70	Insulin-like growth factor binding protein-3 links obesity and breast cancer progression. <i>Oncotarget</i> , 2016 , 7, 55491-55505	3.3	14
69	Shorter telomeres in adults with Type 1 diabetes correlate with diabetes duration, but only weakly with vascular function and risk factors. <i>Diabetes Research and Clinical Practice</i> , 2016 , 117, 4-11	7.4	12
68	An Inverse Relationship Between Age of Type 2 Diabetes Onset and Complication Risk and Mortality: The Impact of Youth-Onset Type 2 Diabetes. <i>Diabetes Care</i> , 2016 , 39, 823-9	14.6	111
67	Monocyte CD163 is altered in association with diabetic complications: possible protective role. <i>Journal of Leukocyte Biology</i> , 2016 , 100, 1375-1383	6.5	19

66	Once-daily liraglutide (1.2 mg) compared with twice-daily exenatide (10 µg) in the treatment of type 2 diabetes patients: An indirect treatment comparison meta-analysis. <i>Journal of Diabetes</i> , 2016 , 8, 866-876	3.8	4
65	Screening for Celiac Disease in Type 1 Diabetes: A Systematic Review. <i>Pediatrics</i> , 2015 , 136, e170-6	7.4	84
64	Lower serum fibroblast activation protein shows promise in the exclusion of clinically significant liver fibrosis due to non-alcoholic fatty liver disease in diabetes and obesity. <i>Diabetes Research and Clinical Practice</i> , 2015 , 108, 466-72	7.4	15
63	Congestive heart failure presence predicts delayed healing of foot ulcers in diabetes: An audit from a multidisciplinary high-risk foot clinic. <i>Journal of Diabetes and Its Complications</i> , 2015 , 29, 556-62	3.2	17
62	Chronic erythropoietin treatment improves diet-induced glucose intolerance in rats. <i>Journal of Endocrinology</i> , 2015 , 225, 77-88	4.7	14
61	An association of large-fibre peripheral nerve dysfunction with non-invasive measures of liver fibrosis secondary to non-alcoholic fatty liver disease in diabetes. <i>Journal of Diabetes and Its Complications</i> , 2015 , 29, 1240-7	3.2	19
60	CCN2 requires TGF-β signalling to regulate CCAAT/enhancer binding proteins and inhibit fat cell differentiation. <i>Journal of Cell Communication and Signaling</i> , 2015 , 9, 27-36	5.2	6
59	The effects of high-fat feeding on physical function and skeletal muscle extracellular matrix. <i>Nutrition and Diabetes</i> , 2015 , 5, e187	4.7	18
58	Circulating dipeptidyl peptidase-4 activity correlates with measures of hepatocyte apoptosis and fibrosis in non-alcoholic fatty liver disease in type 2 diabetes mellitus and obesity: A dual cohort cross-sectional study. <i>Journal of Diabetes</i> , 2015 , 7, 809-19	3.8	37
57	The imperative to prevent diabetes complications: a broadening spectrum and an increasing burden despite improved outcomes. <i>Medical Journal of Australia</i> , 2015 , 202, 300-4	4	6
56	Topically applied connective tissue growth factor/CCN2 improves diabetic preclinical cutaneous wound healing: potential role for CTGF in human diabetic foot ulcer healing. <i>Journal of Diabetes Research</i> , 2015 , 2015, 236238	3.9	32
55	A mitotic cause of Whipple's triad: non-islet cell tumour hypoglycaemia in incurable low-grade malignancy. <i>BMJ Case Reports</i> , 2015 , 2015,	0.9	0
54	Ethnic specific differences in survival of patients with type 2 diabetes: analysis of data collected from an Australian multi-ethnic cohort over a 25 year period. <i>Diabetes Research and Clinical Practice</i> , 2015 , 107, 130-8	7.4	3
53	Opioid-induced secondary adrenal insufficiency presenting as hypercalcaemia. <i>Endocrinology, Diabetes and Metabolism Case Reports</i> , 2015 , 2015, 150035	1.4	7
52	Topical application of the bee hive protectant propolis is well tolerated and improves human diabetic foot ulcer healing in a prospective feasibility study. <i>Journal of Diabetes and Its Complications</i> , 2014 , 28, 850-7	3.2	44
51	Reduction of ARNT in myeloid cells causes immune suppression and delayed wound healing. <i>American Journal of Physiology - Cell Physiology</i> , 2014 , 307, C349-57	5.4	13
50	Medication Safety: an audit of medication discrepancies in transferring type 2 diabetes mellitus (T2DM) patients from Australian primary care to tertiary ambulatory care. <i>International Journal for Quality in Health Care</i> , 2014 , 26, 397-403	1.9	14
49	Age of diabetes diagnosis and diabetes duration associate with glycated haemoglobin. <i>Diabetes Research and Clinical Practice</i> , 2014 , 104, e1-4	7.4	5

48	Quantitation of fibroblast activation protein (FAP)-specific protease activity in mouse, baboon and human fluids and organs. <i>FEBS Open Bio</i> , 2013 , 4, 43-54	2.7	65
47	CCN2 plays a key role in extracellular matrix gene expression in severe hypertrophic cardiomyopathy and heart failure. <i>Journal of Molecular and Cellular Cardiology</i> , 2013 , 62, 164-78	5.8	35
46	Diabetes and nonalcoholic Fatty liver disease: a pathogenic duo. <i>Endocrine Reviews</i> , 2013 , 34, 84-129	27.2	157
45	Long-term complications and mortality in young-onset diabetes: type 2 diabetes is more hazardous and lethal than type 1 diabetes. <i>Diabetes Care</i> , 2013 , 36, 3863-9	14.6	255
44	Connective tissue growth factor/CCN-2 is upregulated in epididymal and subcutaneous fat depots in a dietary-induced obesity model. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2013 , 304, E1291-302	6	18
43	Interaction between IGF binding protein-3 and TGF β in the regulation of adipocyte differentiation. <i>Endocrinology</i> , 2012 , 153, 4799-807	4.8	14
42	Diabetes is a progression factor for hepatic fibrosis in a high fat fed mouse obesity model of non-alcoholic steatohepatitis. <i>Journal of Hepatology</i> , 2011 , 55, 435-44	13.4	72
41	Report on the 6th international workshop of the CCN family of genes. <i>Journal of Cell Communication and Signaling</i> , 2011 , 5, 1-3	5.2	
40	CCN-2 is up-regulated by and mediates effects of matrix bound advanced glycated end-products in human renal mesangial cells. <i>Journal of Cell Communication and Signaling</i> , 2011 , 5, 193-200	5.2	10
39	Suboptimal performance of blood glucose meters in an antenatal diabetes clinic. <i>Diabetes Care</i> , 2011 , 34, 335-7	14.6	19
38	Post-mortem pathologic and genetic studies in "dead in bed syndrome" cases in type 1 diabetes mellitus. <i>Human Pathology</i> , 2010 , 41, 392-400	3.7	21
37	Sudden death in type 1 diabetes: the mystery of the dead in bed syndrome. <i>International Journal of Cardiology</i> , 2010 , 138, 91-3	3.2	38
36	A novel primate model of delayed wound healing in diabetes: dysregulation of connective tissue growth factor. <i>Diabetologia</i> , 2010 , 53, 572-83	10.3	28
35	Regulation of pro-inflammatory and pro-fibrotic factors by CCN2/CTGF in H9c2 cardiomyocytes. <i>Journal of Cell Communication and Signaling</i> , 2010 , 4, 15-23	5.2	53
34	Mastering a mediator: blockade of CCN-2 shows early promise in human diabetic kidney disease. <i>Journal of Cell Communication and Signaling</i> , 2010 , 4, 189-96	5.2	8
33	Adverse effects of high glucose and free fatty acid on cardiomyocytes are mediated by connective tissue growth factor. <i>American Journal of Physiology - Cell Physiology</i> , 2009 , 297, C1490-500	5.4	51
32	Inhibition of adipocyte differentiation by insulin-like growth factor-binding protein-3. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2009 , 296, E654-63	6	73
31	Increased matrix metalloproteinase-9 predicts poor wound healing in diabetic foot ulcers. <i>Diabetes Care</i> , 2009 , 32, 117-9	14.6	239

30	Actions of IGF binding proteins and related proteins in adipose tissue. <i>Trends in Endocrinology and Metabolism</i> , 2009 , 20, 499-505	8.8	41
29	Position statement of the Australian Diabetes Society: individualisation of glycated haemoglobin targets for adults with diabetes mellitus. <i>Medical Journal of Australia</i> , 2009 , 191, 339-44	4	51
28	The anti-inflammatory agent Propolis improves wound healing in a rodent model of experimental diabetes. <i>Wound Repair and Regeneration</i> , 2008 , 16, 706-13	3.6	53
27	Diastolic dysfunction and abnormalities of the microcirculation in type 2 diabetes. <i>Diabetes, Obesity and Metabolism</i> , 2008 , 10, 739-46	6.7	53
26	The metabolic syndrome in type 1 diabetes: does it exist and does it matter?. <i>Journal of Diabetes and Its Complications</i> , 2008 , 22, 18-23	3.2	58
25	Renal connective tissue growth factor correlates with glomerular basement membrane thickness and prospective albuminuria in a non-human primate model of diabetes: possible predictive marker for incipient diabetic nephropathy. <i>Journal of Diabetes and Its Complications</i> , 2008 , 22, 284-94	3.2	50
24	Connective tissue growth factor inhibits adipocyte differentiation. <i>American Journal of Physiology - Cell Physiology</i> , 2008 , 295, C740-51	5.4	76
23	Timing is everything: age of onset influences long-term retinopathy risk in type 2 diabetes, independent of traditional risk factors. <i>Diabetes Care</i> , 2008 , 31, 1985-90	14.6	97
22	Causes of death in young Australians with type 1 diabetes: a review of coronial postmortem examinations. <i>Medical Journal of Australia</i> , 2008 , 188, 699-702	4	36
21	Prediabetes: a position statement from the Australian Diabetes Society and Australian Diabetes Educators Association. <i>Medical Journal of Australia</i> , 2007 , 186, 461-5	4	94
20	Insulin levels in insulin resistance: phantom of the metabolic opera?. <i>Medical Journal of Australia</i> , 2007 , 186, 271-272	4	
19	Alterations in liver sinusoidal endothelium in a baboon model of type 1 diabetes. <i>Diabetologia</i> , 2007 , 50, 1969-1976	10.3	19
18	Bacterial load predicts healing rate in neuropathic diabetic foot ulcers. <i>Diabetes Care</i> , 2007 , 30, 378-80	14.6	74
17	Improving wound-healing outcomes in diabetic foot ulcers. <i>Expert Review of Endocrinology and Metabolism</i> , 2007 , 2, 205-213	4.1	1
16	Connective tissue growth factor plays an important role in advanced glycation end product-induced tubular epithelial-to-mesenchymal transition: implications for diabetic renal disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2006 , 17, 2484-94	12.7	218
15	The metabolic syndrome in type 2 diabetes: When does it matter?. <i>Diabetes, Obesity and Metabolism</i> , 2006 , 8, 690-7	6.7	10
14	Connective tissue growth factor mediates high glucose effects on matrix degradation through tissue inhibitor of matrix metalloproteinase type 1: implications for diabetic nephropathy. <i>Endocrinology</i> , 2004 , 145, 5646-55	4.8	90
13	Connective tissue growth factor is up-regulated in the diabetic retina: amelioration by angiotensin-converting enzyme inhibition. <i>Endocrinology</i> , 2004 , 145, 860-6	4.8	64

12	Monocyte adhesion to decidual endothelial cells is increased in pregnancies complicated by type 1 diabetes but not by gestational diabetes. <i>Diabetes Care</i> , 2004 , 27, 2514-5	14.6	4
11	The time has come to target connective tissue growth factor in diabetic complications. <i>Diabetologia</i> , 2004 , 47, 965-8	10.3	43
10	A breaker of advanced glycation end products attenuates diabetes-induced myocardial structural changes. <i>Circulation Research</i> , 2003 , 92, 785-92	15.7	363
9	Connective tissue growth factor/IGF-binding protein-related protein-2 is a mediator in the induction of fibronectin by advanced glycosylation end-products in human dermal fibroblasts. <i>Endocrinology</i> , 2002 , 143, 1260-9	4.8	86
8	Prevention of accelerated atherosclerosis by angiotensin-converting enzyme inhibition in diabetic apolipoprotein E-deficient mice. <i>Circulation</i> , 2002 , 106, 246-53	16.7	236
7	Renal connective tissue growth factor induction in experimental diabetes is prevented by aminoguanidine. <i>Endocrinology</i> , 2002 , 143, 4907-15	4.8	127
6	Advanced glycosylation end products up-regulate connective tissue growth factor (insulin-like growth factor-binding protein-related protein 2) in human fibroblasts: a potential mechanism for expansion of extracellular matrix in diabetes mellitus. <i>Endocrinology</i> , 2001 , 142, 1760-9	4.8	135
5	Loss of heterozygosity in sporadic parathyroid tumours: involvement of chromosome 1 and the MEN1 gene locus in 11q13. <i>Clinical Endocrinology</i> , 2000 , 53, 85-92	3.4	25
4	A purified bovine serum albumin preparation contains an insulin-like growth factor (IGF) binding protein-3 fragment that forms ternary complexes selectively with IGF-II and the acid-labile subunit. <i>Growth Hormone and IGF Research</i> , 2000 , 10, 215-23	2	4
3	Evaluation of retinoblastoma and Ki-67 immunostaining as diagnostic markers of benign and malignant parathyroid disease. <i>World Journal of Surgery</i> , 1999 , 23, 68-74	3.3	70
2	Precipitated insulin: a potentially life-threatening problem. <i>Australian and New Zealand Journal of Medicine</i> , 1994 , 24, 574		
1	Advanced Glycosylation End Products Up-Regulate Connective Tissue Growth Factor (Insulin-Like Growth Factor-Binding Protein-Related Protein 2) in Human Fibroblasts: A Potential Mechanism for Expansion of Extracellular Matrix in Diabetes Mellitus		37