

Ying Xiao

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/368105/publications.pdf>

Version: 2024-02-01

24
papers

432
citations

840776

11
h-index

794594

19
g-index

27
all docs

27
docs citations

27
times ranked

490
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 1 | Atorvastatin Restores PPAR α Inhibition of Lipid Metabolism Disorders by Downregulating miR-21 Expression to Improve Mitochondrial Function and Alleviate Diabetic Nephropathy Progression. <i>Frontiers in Pharmacology</i> , 2022, 13, 819787. | 3.5 | 7 |
| 2 | BMP-7 ameliorates partial epithelial-mesenchymal transition by restoring SnoN protein level via Smad1/5 pathway in diabetic kidney disease. <i>Cell Death and Disease</i> , 2022, 13, 254. | 6.3 | 14 |
| 3 | Autophagy-related protein EI24 delays the development of pulmonary fibrosis by promoting autophagy. <i>Life Sciences</i> , 2021, 264, 118664. | 4.3 | 6 |
| 4 | The role of CDX2 in renal tubular lesions during diabetic kidney disease. <i>Aging</i> , 2021, 13, 6782-6803. | 3.1 | 10 |
| 5 | Outer membrane protein A inhibits the degradation of caspase-1 to regulate NLRP3 inflammasome activation and exacerbate the <i>Acinetobacter baumannii</i> pulmonary inflammation. <i>Microbial Pathogenesis</i> , 2021, 153, 104788. | 2.9 | 10 |
| 6 | Identification of YAP1 as a novel downstream effector of the FGF2/STAT3 pathway in the pathogenesis of renal tubulointerstitial fibrosis. <i>Journal of Cellular Physiology</i> , 2021, 236, 7655-7671. | 4.1 | 10 |
| 7 | YAP1 Overexpression Is Associated with Kidney Dysfunction in Lupus Nephritis. <i>Pathobiology</i> , 2021, 88, 412-423. | 3.8 | 4 |
| 8 | Blood glucose control contributes to protein stability of Ski α -related novel protein β in a rat model of diabetes. <i>Experimental and Therapeutic Medicine</i> , 2021, 22, 1341. | 1.8 | 1 |
| 9 | SAA1 is transcriptionally activated by STAT3 and accelerates renal interstitial fibrosis by inducing endoplasmic reticulum stress. <i>Experimental Cell Research</i> , 2021, 408, 112856. | 2.6 | 13 |
| 10 | EI24 alleviates renal interstitial fibrosis through inhibition of epithelial \rightarrow mesenchymal transition and fibroblast activation. <i>FASEB Journal</i> , 2021, 35, e21239. | 0.5 | 2 |
| 11 | Smad2 and Smad3 play antagonistic roles in high glucose-induced renal tubular fibrosis via the regulation of SnoN. <i>Experimental and Molecular Pathology</i> , 2020, 113, 104375. | 2.1 | 14 |
| 12 | Oxymatrine Inhibits Twist-Mediated Renal Tubulointerstitial Fibrosis by Upregulating Id2 Expression. <i>Frontiers in Physiology</i> , 2020, 11, 599. | 2.8 | 13 |
| 13 | MicroRNA-27a targets Sfrp1 to induce renal fibrosis in diabetic nephropathy by activating Wnt/ β -Catenin signalling. <i>Bioscience Reports</i> , 2020, 40, . | 2.4 | 18 |
| 14 | BMP-7/Smads-induced inhibitor of differentiation 2 (Id2) upregulation and Id2/Twist interaction was involved in attenuating diabetic renal tubulointerstitial fibrosis. <i>International Journal of Biochemistry and Cell Biology</i> , 2019, 116, 105613. | 2.8 | 11 |
| 15 | BMP-7 inhibits renal fibrosis in diabetic nephropathy via miR-21 downregulation. <i>Life Sciences</i> , 2019, 238, 116957. | 4.3 | 57 |
| 16 | Regulation of PTEN/AKT/FAK pathways by PPAR δ impacts on fibrosis in diabetic nephropathy. <i>Journal of Cellular Biochemistry</i> , 2019, 120, 6998-7014. | 2.6 | 19 |
| 17 | TAK1 may promote the development of diabetic nephropathy by reducing the stability of SnoN protein. <i>Life Sciences</i> , 2019, 228, 1-10. | 4.3 | 6 |
| 18 | Ski α -related novel protein suppresses the development of diabetic nephropathy by modulating transforming growth factor β signaling and microRNA α 21 expression. <i>Journal of Cellular Physiology</i> , 2019, 234, 17925-17936. | 4.1 | 22 |

| # | ARTICLE | IF | CITATIONS |
|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 19 | Notch1 regulates PTEN expression to exacerbate renal tubulointerstitial fibrosis in diabetic nephropathy by inhibiting autophagy via interactions with Hes1. <i>Biochemical and Biophysical Research Communications</i> , 2018, 497, 1110-1116. | 2.1 | 33 |
| 20 | MicroRNA-22 Promotes Renal Tubulointerstitial Fibrosis by Targeting PTEN and Suppressing Autophagy in Diabetic Nephropathy. <i>Journal of Diabetes Research</i> , 2018, 2018, 1-11. | 2.3 | 69 |
| 21 | BMP-7 enhances SnoN mRNA expression in renal tubular epithelial cells under high-glucose conditions. <i>Molecular Medicine Reports</i> , 2017, 16, 3308-3314. | 2.4 | 12 |
| 22 | SnoN upregulation ameliorates renal fibrosis in diabetic nephropathy. <i>PLoS ONE</i> , 2017, 12, e0174471. | 2.5 | 18 |
| 23 | Suberoylanilide hydroxamic acid attenuates paraquat-induced pulmonary fibrosis by preventing Smad7 from deacetylation in rats. <i>Journal of Thoracic Disease</i> , 2016, 8, 2485-2494. | 1.4 | 22 |
| 24 | Oxymatrine Inhibits Renal Tubular EMT Induced by High Glucose via Upregulation of SnoN and Inhibition of TGF- β 1/Smad Signaling Pathway. <i>PLoS ONE</i> , 2016, 11, e0151986. | 2.5 | 40 |