

Peter Junker

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3681241/peter-junker-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

76
papers

1,186
citations

18
h-index

32
g-index

82
ext. papers

1,400
ext. citations

3.3
avg, IF

3.56
L-index

| # | Paper | IF | Citations |
|----|---|-----|-----------|
| 76 | Increased synovial galectin-3 induce inflammatory fibroblast activation and osteoclastogenesis in patients with rheumatoid arthritis.. <i>Scandinavian Journal of Rheumatology</i> , 2022 , 1-9 | 1.9 | 1 |
| 75 | Erratum corrige: Increased serum levels of microfibrillar-associated protein 4 (MFAP4) are not associated with clinical synovitis in rheumatoid arthritis but may reflect underlying cardiovascular comorbidity. <i>Clinical and Experimental Rheumatology</i> , 2022 , 40, 198-198 | 2.2 | |
| 74 | The Programmed Death-1 Pathway Counter-Regulates Inflammation-Induced Osteoclast Activity in Clinical and Experimental Settings.. <i>Frontiers in Immunology</i> , 2022 , 13, 773946 | 8.4 | 2 |
| 73 | Altered Antibody Response to Epstein-Barr Virus in Patients With Rheumatoid Arthritis and Healthy Subjects Predisposed to the Disease. A Twin Study. <i>Frontiers in Immunology</i> , 2021 , 12, 650713 | 8.4 | 1 |
| 72 | Circadian, Week-to-Week, and Physical Exercise-Induced Variation of Serum Microfibrillar-Associated Protein 4. <i>Biomarker Insights</i> , 2021 , 16, 11772719211016359 | 3.5 | 0 |
| 71 | Similar lipid level changes in early rheumatoid arthritis patients following 1-year treat-to-target strategy with adalimumab plus methotrexate versus placebo plus methotrexate: secondary analyses from the randomised controlled OPERA trial. <i>Rheumatology International</i> , 2021 , 41, 543-549 | 3.6 | 2 |
| 70 | Impact of season on the association between vitamin D levels at diagnosis and one-year remission in early Rheumatoid Arthritis. <i>Scientific Reports</i> , 2020 , 10, 7371 | 4.9 | 1 |
| 69 | AB0194 VITAMIN D TRAJECTORIES IN EARLY DIAGNOSED, AGGRESSIVELY TREATED RHEUMATOID ARTHRITIS PATIENTS: A 10 YEAR LONGITUDINAL COHORT STUDY BASED ON THE DANISH CIMESTRA TRIAL. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 1397-1397 | 2.4 | |
| 68 | Programmed death ligand 2 - A link between inflammation and bone loss in rheumatoid arthritis. <i>Journal of Translational Autoimmunity</i> , 2020 , 3, 100028 | 4.1 | 5 |
| 67 | Increased serum levels of microfibrillar-associated protein 4 (MFAP4) are not associated with clinical synovitis in rheumatoid arthritis but may reflect underlying cardiovascular comorbidity. <i>Clinical and Experimental Rheumatology</i> , 2020 , 38, 122-128 | 2.2 | 0 |
| 66 | Site-specific absence of microfibrillar-associated protein 4 (MFAP4) from the internal elastic membrane of arterioles in the rheumatoid arthritis synovial membrane: an immunohistochemical study in patients with advanced rheumatoid arthritis versus osteoarthritis. <i>Apmis</i> , 2019 , 127, 588-593 | 3.4 | 6 |
| 65 | IgG4-related disease in patients with newly diagnosed idiopathic retroperitoneal fibrosis: a population-based Danish study. <i>Scandinavian Journal of Rheumatology</i> , 2019 , 48, 320-325 | 1.9 | 4 |
| 64 | Validity and Responsiveness of Combined Inflammation and Combined Joint Damage Scores Based on the OMERACT Rheumatoid Arthritis MRI Scoring System (RAMRIS). <i>Journal of Rheumatology</i> , 2019 , 46, 1222-1227 | 4.1 | 5 |
| 63 | Anti-cyclic citrullinated peptide antibodies, 28-joint Disease Activity Score, and magnetic resonance imaging bone oedema at baseline predict 11 years' functional and radiographic outcome in early rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2019 , 48, 1-8 | 1.9 | 15 |
| 62 | Predictive value of a multi-biomarker disease activity score for clinical remission and radiographic progression in patients with early rheumatoid arthritis: a post-hoc study of the OPERA trial. <i>Scandinavian Journal of Rheumatology</i> , 2019 , 48, 9-16 | 1.9 | 14 |
| 61 | Immune thrombocytopenia is associated with persistently deranged fibrosis-related seromarker profiles but low bone marrow fibrosis grades: A 2-year observational study on thrombopoietin receptor agonist treatment. <i>Platelets</i> , 2019 , 30, 222-228 | 3.6 | 1 |
| 60 | Circulating serum interleukin-6, serum chitinase-3-like protein-1, and plasma vascular endothelial growth factor are not predictive for remission and radiographic progression in patients with early rheumatoid arthritis: post-hoc explorative and validation studies based on the CIMESTRA and OPERA trials. <i>Scandinavian Journal of Rheumatology</i> , 2019 , 47, 259-269 | 1.9 | 2 |

| | | | |
|----|---|-----|----|
| 59 | The D-vitamin metabolite 1,25(OH) D in serum is associated with disease activity and Anti-Citrullinated Protein Antibodies in active and treatment naïve, early Rheumatoid Arthritis Patients. <i>Scandinavian Journal of Immunology</i> , 2018 , 88, e12704 | 3.4 | 8 |
| 58 | Surfactant protein-D, a potential mediator of inflammation in axial spondyloarthritis. <i>Rheumatology</i> , 2018 , 57, 1861-1865 | 3.9 | 3 |
| 57 | Plasma MicroRNA Profiles in Patients with Early Rheumatoid Arthritis Responding to Adalimumab plus Methotrexate vs Methotrexate Alone: A Placebo-controlled Clinical Trial. <i>Journal of Rheumatology</i> , 2018 , 45, 53-61 | 4.1 | 23 |
| 56 | Localization of surfactant protein-D in the rheumatoid synovial membrane. <i>Apmis</i> , 2018 , 126, 9-13 | 3.4 | 4 |
| 55 | Type IV collagen metabolism is associated with disease activity, radiographic progression and response to tocilizumab in rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36, 829-832 | 3.2 | 11 |
| 54 | Association between baseline vitamin D metabolite levels and long-term cardiovascular events in patients with rheumatoid arthritis from the CIMESTRA trial: protocol for a cohort study with patient-record evaluated outcomes. <i>BMJ Open</i> , 2017 , 7, e014816 | 3 | 2 |
| 53 | Hepcidin plasma levels are not associated with changes in haemoglobin in early rheumatoid arthritis patients. <i>Scandinavian Journal of Rheumatology</i> , 2017 , 46, 441-445 | 1.9 | 6 |
| 52 | Soluble CD206 plasma levels in rheumatoid arthritis reflect decrease in disease activity. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2017 , 77, 385-389 | 2 | 6 |
| 51 | Magnetic resonance imaging assessed inflammation in the wrist is associated with patient-reported physical impairment, global assessment of disease activity and pain in early rheumatoid arthritis: longitudinal results from two randomised controlled trials. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 1707-1715 | 2.4 | 21 |
| 50 | Expression of soluble CD83 in plasma from early-stage rheumatoid arthritis patients is not modified by anti-TNF- α therapy. <i>Cytokine</i> , 2017 , 96, 1-7 | 4 | 16 |
| 49 | Hand bone loss in early rheumatoid arthritis during a methotrexate-based treat-to-target strategy with or without adalimumab-a substudy of the optimized treatment algorithm in early RA (OPERA) trial. <i>Clinical Rheumatology</i> , 2017 , 36, 781-789 | 3.9 | 7 |
| 48 | Galectin-3 is Persistently Increased in Early Rheumatoid Arthritis (RA) and Associates with Anti-CCP Seropositivity and MRI Bone Lesions, While Early Fibrosis Markers Correlate with Disease Activity. <i>Scandinavian Journal of Immunology</i> , 2017 , 86, 471-478 | 3.4 | 14 |
| 47 | Increased galectin-3 may serve as a serologic signature of pre-rheumatoid arthritis while markers of synovitis and cartilage do not differ between early undifferentiated arthritis subsets. <i>Arthritis Research and Therapy</i> , 2017 , 19, 80 | 5.7 | 10 |
| 46 | Effect of a treat-to-target strategy based on methotrexate and intra-articular betamethasone with or without additional cyclosporin on MRI-assessed synovitis, osteitis, tenosynovitis, bone erosion, and joint space narrowing in early rheumatoid arthritis: results from a 2-year randomized controlled trial. <i>Scandinavian Journal of Immunology</i> , 2017 , 86, 335-345 | 1.9 | 7 |
| 45 | No diagnostic utility of antibody patterns against <i>Klebsiella pneumoniae</i> capsular serotypes in patients with axial spondyloarthritis vs. patients with non-specific low back pain: a cross-sectional study. <i>Scandinavian Journal of Rheumatology</i> , 2017 , 46, 296-302 | 1.9 | 6 |
| 44 | Incidence of Chronic Persistent Rheumatoid Arthritis and the Impact of Smoking: A Historical Twin Cohort Study. <i>Arthritis Care and Research</i> , 2017 , 69, 616-624 | 4.7 | 18 |
| 43 | Type I and III collagen turnover is increased in axial spondyloarthritis and psoriatic arthritis. Associations with disease activity and diagnostic capacity. <i>Clinical and Experimental Rheumatology</i> , 2017 , 35, 653-659 | 2.2 | 6 |
| 42 | Clinical and radiographic outcome of a treat-to-target strategy using methotrexate and intra-articular glucocorticoids with or without adalimumab induction: a 2-year investigator-initiated, double-blinded, randomised, controlled trial (OPERA). <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1645-53 | 2.4 | 30 |

| | | | |
|----|--|-----|----|
| 41 | Prediction of treatment response to adalimumab: a double-blind placebo-controlled study of circulating microRNA in patients with early rheumatoid arthritis. <i>Pharmacogenomics Journal</i> , 2016 , 16, 141-6 | 3.5 | 44 |
| 40 | FRI0535 Hand Bone Loss in Early Rheumatoid Arthritis Is Independent of Adalimumab Treatment. A Substudy of The Optimized Treatment Algorithm in Early RA (Opera) Trial. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 633.1-633 | 2.4 | |
| 39 | The interleukin-20 receptor axis in early rheumatoid arthritis: novel links between disease-associated autoantibodies and radiographic progression. <i>Arthritis Research and Therapy</i> , 2016 , 18, 61 | 5.7 | 24 |
| 38 | Single test isolated lupus anticoagulant positivity is associated with increased plasma levels of inflammatory markers and dyslipidemia. <i>Lupus</i> , 2016 , 25, 241-7 | 2.6 | 5 |
| 37 | Cartilage collagen type II seromarker patterns in axial spondyloarthritis and psoriatic arthritis: associations with disease activity, smoking and HLA-B27. <i>Rheumatology International</i> , 2016 , 36, 541-9 | 3.6 | 8 |
| 36 | Are human endogenous retroviruses triggers of autoimmune diseases? Unveiling associations of three diseases and viral loci. <i>Immunologic Research</i> , 2016 , 64, 55-63 | 4.3 | 29 |
| 35 | Differentially Methylated DNA Regions in Monozygotic Twin Pairs Discordant for Rheumatoid Arthritis: An Epigenome-Wide Study. <i>Frontiers in Immunology</i> , 2016 , 7, 510 | 8.4 | 18 |
| 34 | Chondrocyte activity is increased in psoriatic arthritis and axial spondyloarthritis. <i>Arthritis Research and Therapy</i> , 2016 , 18, 141 | 5.7 | 15 |
| 33 | Changes in Soluble CD18 in Murine Autoimmune Arthritis and Rheumatoid Arthritis Reflect Disease Establishment and Treatment Response. <i>PLoS ONE</i> , 2016 , 11, e0148486 | 3.7 | 10 |
| 32 | A disintegrin and metalloprotease-17 and galectin-9 are important regulators of local 4-1BB activity and disease outcome in rheumatoid arthritis. <i>Rheumatology</i> , 2016 , 55, 1871-9 | 3.9 | 9 |
| 31 | Cardiac Abnormalities in Adult Patients With Polymyositis or Dermatomyositis as Assessed by Noninvasive Modalities. <i>Arthritis Care and Research</i> , 2016 , 68, 1012-20 | 4.7 | 31 |
| 30 | Within-day variation and influence of physical exercise on circulating Galectin-3 in patients with rheumatoid arthritis and healthy individuals. <i>Scandinavian Journal of Immunology</i> , 2015 , 82, 70-5 | 3.4 | 13 |
| 29 | Traditional cardiovascular risk factors and coronary artery calcification in adults with polymyositis and dermatomyositis: a Danish multicenter study. <i>Arthritis Care and Research</i> , 2015 , 67, 848-54 | 4.7 | 29 |
| 28 | AB0648 Disease Mapping in Anca Associated Vasculitis (AAV). Yield of FDG-PET/CT Imaging Versus a Conventional Clinico-Pathological Approach,. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1115.3-1116 | 2.4 | |
| 27 | OP0073 Polymorphisms in the FCN1 Gene Coding for M-Ficolin are Associated with Disease Activity, Radiographic Damage and are the Strongest Predictors of DAS28 Remission in 180 DMARD Naive Early Rheumatoid Arthritis Patients. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 96.1-96 | 2.4 | |
| 26 | A treat-to-target strategy with methotrexate and intra-articular triamcinolone with or without adalimumab effectively reduces MRI synovitis, osteitis and tenosynovitis and halts structural damage progression in early rheumatoid arthritis: results from the OPERA randomised controlled trial. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 917-7 | 2.4 | 44 |
| 25 | Macrophage activity assessed by soluble CD163 in early rheumatoid arthritis: association with disease activity but different response patterns to synthetic and biologic DMARDs. <i>Clinical and Experimental Rheumatology</i> , 2015 , 33, 498-502 | 2.2 | 14 |
| 24 | Cardiac abnormalities assessed by non-invasive techniques in patients with newly diagnosed idiopathic inflammatory myopathies. <i>Clinical and Experimental Rheumatology</i> , 2015 , 33, 706-14 | 2.2 | 16 |

| | | | |
|----|--|-----|-----|
| 23 | Newborn infant characteristics and risk of future rheumatoid arthritis: a twin-control study. <i>Rheumatology International</i> , 2014 , 34, 523-8 | 3.6 | 5 |
| 22 | Heritability assessment of cartilage metabolism. A twin study on circulating procollagen IIA N-terminal propeptide (PIIANP). <i>Osteoarthritis and Cartilage</i> , 2014 , 22, 1142-7 | 6.2 | 2 |
| 21 | THU0550 Changes in Soluble CD18 Reflect Latency in Restoration of the Immune System after Early Treatment of Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 373.1-373 | 2.4 | |
| 20 | Adalimumab added to a treat-to-target strategy with methotrexate and intra-articular triamcinolone in early rheumatoid arthritis increased remission rates, function and quality of life. The OPERA Study: an investigator-initiated, randomised, double-blind, parallel-group, placebo-controlled trial. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 654-61 | 2.4 | 84 |
| 19 | Risk of lymphoma and solid cancer among patients with rheumatoid arthritis in a primary care setting. <i>PLoS ONE</i> , 2014 , 9, e99388 | 3.7 | 11 |
| 18 | Palpable purpura with an unexpected outcome. <i>Case Reports in Rheumatology</i> , 2013 , 2013, 678427 | 0.8 | 1 |
| 17 | AB0316 Patient's global assessment of general health by vas at baseline predicts ACR/EULAR remission after 3, 6 and 12 months of efficient treatment in DMARD- and steroid naïve early rheumatoid arthritis patients. The danish cimestra-study. <i>Annals of the Rheumatic Diseases</i> , 2013 , 71, 655.12-655 | 2.4 | |
| 16 | OP0223 M-ficolin, an activator of the complement system, predicts DAS28 remission in early DMARD naïve rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 71, 131.1-131 | 2.4 | 1 |
| 15 | On the origin of rheumatoid arthritis: the impact of environment and genes--a population based twin study. <i>PLoS ONE</i> , 2013 , 8, e57304 | 3.7 | 57 |
| 14 | Uncoupling of collagen II metabolism in newly diagnosed, untreated rheumatoid arthritis is linked to inflammation and antibodies against cyclic citrullinated peptides. <i>Journal of Rheumatology</i> , 2010 , 37, 1113-20 | 4.1 | 12 |
| 13 | Radiographic progression and remission rates in early rheumatoid arthritis - MRI bone oedema and anti-CCP predicted radiographic progression in the 5-year extension of the double-blind randomised CIMESTRA trial. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 1789-95 | 2.4 | 136 |
| 12 | FORMATION OF GRANULATION TISSUE IN SUBCUTANEOUSLY IMPLANTED SPONGES IN RATS. <i>Acta Pathologica Et Microbiologica Scandinavica Section A, Pathology</i> , 2009 , 87A, 367-374 | | |
| 11 | Ankylosing spondylitis in Danish and Norwegian twins: occurrence and the relative importance of genetic vs. environmental effectors in disease causation. <i>Scandinavian Journal of Rheumatology</i> , 2008 , 37, 120-6 | 1.9 | 76 |
| 10 | Combination treatment with methotrexate, cyclosporine, and intraarticular betamethasone compared with methotrexate and intraarticular betamethasone in early active rheumatoid arthritis: an investigator-initiated, multicenter, randomized, double-blind, parallel-group, placebo-controlled study. <i>Arthritis and Rheumatism</i> , 2006 , 54, 1481-9 | | 129 |
| 9 | Relative importance of genetic effects in rheumatoid arthritis: historical cohort study of Danish nationwide twin population. <i>BMJ, The</i> , 2002 , 324, 264-6 | 5.9 | 34 |
| 8 | Serum hyaluronan is increased in malignant lymphoma. <i>American Journal of Hematology</i> , 1995 , 50, 231-37.1 | | 12 |
| 7 | Serum aminoterminal procollagen type III peptide in acute viral hepatitis. A long-term follow-up study. <i>Liver</i> , 1987 , 7, 96-105 | | 20 |
| 6 | Effect of D-penicillamine pre- and post-implantation treatment on formation of sponge-induced granulation tissue in rats. <i>Scandinavian Journal of Rheumatology</i> , 1984 , 13, 222-8 | 1.9 | 2 |

- 5 Third Euler Workshop on Rheumatology Research. *Clinical Rheumatology*, **1983**, 2, 81-106 3.9
- 4 The effect of D-penicillamine and methylprednisolone on the morphology of experimental granulation tissue in rats. *Acta Pathologica, Microbiologica, Et Immunologica Scandinavica Section A, Pathology*, **1982**, 90, 51-5 1
- 3 Fibronectin in experimental granulation tissue. *Acta Pathologica, Microbiologica, Et Immunologica Scandinavica Section A, Pathology*, **1982**, 90, 159-65 20
- 2 Effect of different doses of D-penicillamine and combined administration of D-penicillamine and methylprednisolone on collagen, glycosaminoglycans, DNA and RNA of granulation tissue, skin, bone and aorta in rats. *Acta Pharmacologica Et Toxicologica*, **1981**, 49, 366-80 7
- 1 Effect of D-penicillamine on collagen, glycosaminoglycans, DNA and RNA of granulation tissue and connective tissue of skin, bone and aorta in rats. *Acta Pharmacologica Et Toxicologica*, **1981**, 48, 300-10 9