

Oscar Fernandez-Capetillo

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3684011/oscar-fernandez-capetillo-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

109
papers

14,560
citations

51
h-index

117
g-index

117
ext. papers

16,228
ext. citations

14
avg, IF

6.06
L-index

#	Paper	IF	Citations
109	A bispecific monomeric nanobody induces spike trimer dimers and neutralizes SARS-CoV-2 in vivo.. <i>Nature Communications</i> , 2022 , 13, 155	17.4	6
108	USP7 and VCP define the SUMO/Ubiquitin landscape at the DNA replication fork. <i>Cell Reports</i> , 2021 , 37, 109819	10.6	1
107	USP7 limits CDK1 activity throughout the cell cycle. <i>EMBO Journal</i> , 2021 , 40, e99692	13	1
106	A chemical screen for modulators of mRNA translation identifies a distinct mechanism of toxicity for sphingosine kinase inhibitors. <i>PLoS Biology</i> , 2021 , 19, e3001263	9.7	1
105	Widespread displacement of DNA- and RNA-binding factors underlies toxicity of arginine-rich cell-penetrating peptides. <i>EMBO Journal</i> , 2021 , 40, e103311	13	7
104	Overview of virus and cancer relationships. Position paper. <i>Revista Espanola De Quimioterapia</i> , 2021 , 34, 525-555	1.6	0
103	Distinct roles for PARP-1 and PARP-2 in c-Myc-driven B-cell lymphoma in mice. <i>Blood</i> , 2021 ,	2.2	4
102	Prolonged estrogen deprivation triggers a broad immunosuppressive phenotype in breast cancer cells. <i>Molecular Oncology</i> , 2021 ,	7.9	6
101	Supraphysiological protection from replication stress does not extend mammalian lifespan. <i>Aging</i> , 2020 , 12, 5612-5624	5.6	
100	ATR expands embryonic stem cell fate potential in response to replication stress. <i>ELife</i> , 2020 , 9,	8.9	15
99	A Chemical Screen Identifies Compounds Capable of Selecting for Haploidy in Mammalian Cells. <i>Cell Reports</i> , 2019 , 28, 597-604.e4	10.6	10
98	TIAR marks nuclear G2/M transition granules and restricts CDK1 activity under replication stress. <i>EMBO Reports</i> , 2019 , 20,	6.5	6
97	A Chemical Screen Identifies Compounds Limiting the Toxicity of C9ORF72 Dipeptide Repeats. <i>Cell Chemical Biology</i> , 2019 , 26, 235-243.e5	8.2	8
96	ERF deletion rescues RAS deficiency in mouse embryonic stem cells. <i>Genes and Development</i> , 2018 , 32, 568-576	12.6	7
95	The RNA Polymerase II Factor RPAP1 Is Critical for Mediator-Driven Transcription and Cell Identity. <i>Cell Reports</i> , 2018 , 22, 396-410	10.6	15
94	ATR is required to complete meiotic recombination in mice. <i>Nature Communications</i> , 2018 , 9, 2622	17.4	29
93	Targeting ATR in cancer. <i>Nature Reviews Cancer</i> , 2018 , 18, 586-595	31.3	136

92	Control of DNA Replication by ATR. <i>Cancer Drug Discovery and Development</i> , 2018 , 35-61	0.3	
91	A p53-dependent response limits the viability of mammalian haploid cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017 , 114, 9367-9372	11.5	33
90	Trap: An RNA Sequencing-Based Pipeline for the Identification of Gene-Trap Insertions in Mammalian Cells. <i>Journal of Molecular Biology</i> , 2017 , 429, 2780-2789	6.5	6
89	POLD3 Is Haploinsufficient for DNA Replication in Mice. <i>Molecular Cell</i> , 2016 , 63, 877-83	17.6	25
88	Replication fork stability confers chemoresistance in BRCA-deficient cells. <i>Nature</i> , 2016 , 535, 382-7	50.4	456
87	Targeting the kinase activities of ATR and ATM exhibits antitumoral activity in mouse models of MLL-rearranged AML. <i>Science Signaling</i> , 2016 , 9, ra91	8.8	50
86	The (elusive) role of the SMC5/6 complex. <i>Cell Cycle</i> , 2016 , 15, 775-6	4.7	7
85	USP7 is a SUMO deubiquitinase essential for DNA replication. <i>Nature Structural and Molecular Biology</i> , 2016 , 23, 270-7	17.6	82
84	Efficacy of ATR inhibitors as single agents in Ewing sarcoma. <i>Oncotarget</i> , 2016 , 7, 58759-58767	3.3	40
83	A Genome-wide CRISPR Screen Identifies CDC25A as a Determinant of Sensitivity to ATR Inhibitors. <i>Molecular Cell</i> , 2016 , 62, 307-313	17.6	110
82	A SUMO and ubiquitin code coordinates protein traffic at replication factories. <i>BioEssays</i> , 2016 , 38, 1209-1217	12.7	7
81	PARP-2 sustains erythropoiesis in mice by limiting replicative stress in erythroid progenitors. <i>Cell Death and Differentiation</i> , 2015 , 22, 1144-57	12.7	47
80	Increased Rrm2 gene dosage reduces fragile site breakage and prolongs survival of ATR mutant mice. <i>Genes and Development</i> , 2015 , 29, 690-5	12.6	43
79	Replication stress caused by low MCM expression limits fetal erythropoiesis and hematopoietic stem cell functionality. <i>Nature Communications</i> , 2015 , 6, 8548	17.4	64
78	A Single Conserved Residue Mediates Binding of the Ribonucleotide Reductase Catalytic Subunit RRM1 to RRM2 and Is Essential for Mouse Development. <i>Molecular and Cellular Biology</i> , 2015 , 35, 2910-7	4.8	6
77	Limiting replication stress during somatic cell reprogramming reduces genomic instability in induced pluripotent stem cells. <i>Nature Communications</i> , 2015 , 6, 8036	17.4	65
76	NSMCE2 suppresses cancer and aging in mice independently of its SUMO ligase activity. <i>EMBO Journal</i> , 2015 , 34, 2604-19	13	34
75	Modeling the study of DNA damage responses in mice. <i>Methods in Molecular Biology</i> , 2015 , 1267, 413-37	1.4	8

74	Replication stress and cancer: it takes two to tango. <i>Experimental Cell Research</i> , 2014 , 329, 26-34	4.2	99
73	p21 suppresses inflammation and tumorigenesis on pRB-deficient stratified epithelia. <i>Oncogene</i> , 2014 , 33, 4599-4612	9.2	11
72	The solute carrier SLC35F2 enables YM155-mediated DNA damage toxicity. <i>Nature Chemical Biology</i> , 2014 , 10, 768-773	11.7	125
71	The maternal side of Fanconi Anemia. <i>Molecular Cell</i> , 2014 , 55, 803-804	17.6	2
70	A synthetic lethal interaction between APC/C and topoisomerase poisons uncovered by proteomic screens. <i>Cell Reports</i> , 2014 , 6, 670-83	10.6	38
69	Emergence and evolutionary analysis of the human DDR network: implications in comparative genomics and downstream analyses. <i>Molecular Biology and Evolution</i> , 2014 , 31, 940-61	8.3	15
68	Fos-dependent induction of Chk1 protects osteoblasts from replication stress. <i>Cell Cycle</i> , 2014 , 13, 1980-67	4.7	10
67	Naked replication forks break apRPA. <i>Cell</i> , 2013 , 155, 979-80	56.2	9
66	Cyclin-dependent kinase inhibitor p21 controls adult neural stem cell expansion by regulating Sox2 gene expression. <i>Cell Stem Cell</i> , 2013 , 12, 88-100	18	142
65	Identification of early replicating fragile sites that contribute to genome instability. <i>Cell</i> , 2013 , 152, 620-32	32.2	280
64	A proteomic characterization of factors enriched at nascent DNA molecules. <i>Cell Reports</i> , 2013 , 3, 1105-16	16.6	87
63	53BP1 mediates productive and mutagenic DNA repair through distinct phosphoprotein interactions. <i>Cell</i> , 2013 , 153, 1266-80	56.2	247
62	Late-replicating CNVs as a source of new genes. <i>Biology Open</i> , 2013 , 2, 1402-11	2.2	6
61	Polycomb protein SCML2 regulates the cell cycle by binding and modulating CDK/CYCLIN/p21 complexes. <i>PLoS Biology</i> , 2013 , 11, e1001737	9.7	22
60	INK4a/ARF limits the expansion of cells suffering from replication stress. <i>Cell Cycle</i> , 2013 , 12, 1948-54	4.7	16
59	BRCA1 functions independently of homologous recombination in DNA interstrand crosslink repair. <i>Molecular Cell</i> , 2012 , 46, 125-35	17.6	212
58	An extra allele of Chk1 limits oncogene-induced replicative stress and promotes transformation. <i>Journal of Experimental Medicine</i> , 2012 , 209, 455-61	16.6	80
57	Oncogenic stress sensitizes murine cancers to hypomorphic suppression of ATR. <i>Journal of Clinical Investigation</i> , 2012 , 122, 241-52	15.9	136

56	An extra allele of Chk1 limits oncogene-induced replicative stress and promotes transformation. <i>Journal of Cell Biology</i> , 2012 , 196, i7-i7	7.3	
55	Targeting ATR and Chk1 kinases for cancer treatment: a new model for new (and old) drugs. <i>Molecular Oncology</i> , 2011 , 5, 368-73	7.9	141
54	Exploiting oncogene-induced replicative stress for the selective killing of Myc-driven tumors. <i>Nature Structural and Molecular Biology</i> , 2011 , 18, 1331-1335	17.6	281
53	Lac operator repeats generate a traceable fragile site in mammalian cells. <i>EMBO Reports</i> , 2011 , 12, 1032-1035	6.5	37
52	A cell-based screen identifies ATR inhibitors with synthetic lethal properties for cancer-associated mutations. <i>Nature Structural and Molecular Biology</i> , 2011 , 18, 721-7	17.6	358
51	Genomic instability in iPS: time for a break. <i>EMBO Journal</i> , 2011 , 30, 991-3	13	40
50	CGK733 does not inhibit ATM or ATR kinase activity in H460 human lung cancer cells. <i>DNA Repair</i> , 2011 , 10, 1000-1; author reply 1002	4.3	9
49	Intrauterine programming of ageing. <i>EMBO Reports</i> , 2010 , 11, 32-6	6.5	27
48	Cdk2 suppresses cellular senescence induced by the c-myc oncogene. <i>Nature Cell Biology</i> , 2010 , 12, 54-9; sup pp 1-14	23.4	184
47	ATR suppresses telomere fragility and recombination but is dispensable for elongation of short telomeres by telomerase. <i>Journal of Cell Biology</i> , 2010 , 188, 639-52	7.3	51
46	Class switching and meiotic defects in mice lacking the E3 ubiquitin ligase RNF8. <i>Journal of Experimental Medicine</i> , 2010 , 207, 973-81	16.6	83
45	Nuclear phosphoinositide 3-kinase beta controls double-strand break DNA repair. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 7491-6	11.5	131
44	Histone H2A C-terminus regulates chromatin dynamics, remodeling, and histone H1 binding. <i>PLoS Genetics</i> , 2010 , 6, e1001234	6	58
43	Sirt1 improves healthy ageing and protects from metabolic syndrome-associated cancer. <i>Nature Communications</i> , 2010 , 1, 3	17.4	456
42	53BP1 inhibits homologous recombination in Brca1-deficient cells by blocking resection of DNA breaks. <i>Cell</i> , 2010 , 141, 243-54	56.2	1147
41	The ATR barrier to replication-born DNA damage. <i>DNA Repair</i> , 2010 , 9, 1249-55	4.3	106
40	Class switching and meiotic defects in mice lacking the E3 ubiquitin ligase RNF8. <i>Journal of Cell Biology</i> , 2010 , 189, i5-i5	7.3	
39	Limited role of murine ATM in oncogene-induced senescence and p53-dependent tumor suppression. <i>PLoS ONE</i> , 2009 , 4, e5475	3.7	43

38	p38 Mitogen-activated protein kinase- and HuR-dependent stabilization of p21(Cip1) mRNA mediates the G(1)/S checkpoint. <i>Molecular and Cellular Biology</i> , 2009 , 29, 4341-51	4.8	183
37	p27Kip1 stabilization is essential for the maintenance of cell cycle arrest in response to DNA damage. <i>Cancer Research</i> , 2009 , 69, 8726-32	10.1	43
36	A p53-mediated DNA damage response limits reprogramming to ensure iPS cell genomic integrity. <i>Nature</i> , 2009 , 460, 1149-53	50.4	842
35	A mouse model of ATR-Seckel shows embryonic replicative stress and accelerated aging. <i>Nature Genetics</i> , 2009 , 41, 891-8	36.3	281
34	Increased telomere fragility and fusions resulting from TRF1 deficiency lead to degenerative pathologies and increased cancer in mice. <i>Genes and Development</i> , 2009 , 23, 2060-75	12.6	275
33	Overall Cdk activity modulates the DNA damage response in mammalian cells. <i>Journal of Cell Biology</i> , 2009 , 187, 773-80	7.3	48
32	ATM breaks into heterochromatin. <i>Molecular Cell</i> , 2008 , 31, 303-4	17.6	9
31	Why cells respond differently to DNA damage: a chromatin perspective. <i>Cell Cycle</i> , 2008 , 7, 980-3	4.7	4
30	ATR signaling can drive cells into senescence in the absence of DNA breaks. <i>Genes and Development</i> , 2008 , 22, 297-302	12.6	126
29	Genomic instability: on the birth and death of cancer. <i>Clinical and Translational Oncology</i> , 2007 , 9, 216-20	3.6	13
28	RAD51C deficiency in mice results in early prophase I arrest in males and sister chromatid separation at metaphase II in females. <i>Journal of Cell Biology</i> , 2007 , 176, 581-92	7.3	96
27	Global chromatin compaction limits the strength of the DNA damage response. <i>Journal of Cell Biology</i> , 2007 , 178, 1101-8	7.3	197
26	ATM regulates ATR chromatin loading in response to DNA double-strand breaks. <i>Journal of Experimental Medicine</i> , 2006 , 203, 297-303	16.6	193
25	Changes in chromatin structure and mobility in living cells at sites of DNA double-strand breaks. <i>Journal of Cell Biology</i> , 2006 , 172, 823-34	7.3	405
24	"ATR activation in response to ionizing radiation: still ATM territory". <i>Cell Division</i> , 2006 , 1, 7	2.8	36
23	ATM regulates ATR chromatin loading in response to DNA double-strand breaks. <i>Journal of Cell Biology</i> , 2006 , 172, i9-i9	7.3	
22	Neuropilin-1 regulates attachment in human endothelial cells independently of vascular endothelial growth factor receptor-2. <i>Blood</i> , 2005 , 105, 1992-9	2.2	102
21	Role of Nbs1 in the activation of the Atm kinase revealed in humanized mouse models. <i>Nature Cell Biology</i> , 2005 , 7, 675-85	23.4	194

20	Silencing of unsynapsed meiotic chromosomes in the mouse. <i>Nature Genetics</i> , 2005 , 37, 41-7	36.3	444
19	Specific association of mouse MDC1/NFBD1 with NBS1 at sites of DNA-damage. <i>Cell Cycle</i> , 2005 , 4, 177-82	4.7	29
18	Linking histone deacetylation with the repair of DNA breaks. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004 , 101, 1427-8	11.5	33
17	Phosphorylation of histone H2B at DNA double-strand breaks. <i>Journal of Experimental Medicine</i> , 2004 , 199, 1671-7	16.6	148
16	H2AX: the histone guardian of the genome. <i>DNA Repair</i> , 2004 , 3, 959-67	4.3	743
15	Focusing on Foci: H2AX and the Recruitment of DNA-Damage Response Factors. <i>Cell Cycle</i> , 2003 , 2, 425-426	4.7	137
14	Histone H2AX phosphorylation is dispensable for the initial recognition of DNA breaks. <i>Nature Cell Biology</i> , 2003 , 5, 675-9	23.4	795
13	H2AX haploinsufficiency modifies genomic stability and tumor susceptibility. <i>Cell</i> , 2003 , 114, 371-383	56.2	475
12	H2AX is required for chromatin remodeling and inactivation of sex chromosomes in male mouse meiosis. <i>Developmental Cell</i> , 2003 , 4, 497-508	10.2	461
11	H2AX regulates meiotic telomere clustering. <i>Journal of Cell Biology</i> , 2003 , 163, 15-20	7.3	54
10	Focusing on foci: H2AX and the recruitment of DNA-damage response factors. <i>Cell Cycle</i> , 2003 , 2, 426-7	4.7	83
9	DNA damage-induced G2-M checkpoint activation by histone H2AX and 53BP1. <i>Nature Cell Biology</i> , 2002 , 4, 993-7	23.4	538
8	Genomic instability in mice lacking histone H2AX. <i>Science</i> , 2002 , 296, 922-7	33.3	1124
7	Extracellular signal-regulated protein kinase signaling pathway negatively regulates the phenotypic and functional maturation of monocyte-derived human dendritic cells. <i>Blood</i> , 2001 , 98, 2175-82	2.2	174
6	Immunoreactivity for the group III metabotropic glutamate receptor subtype mGluR4a in the superficial laminae of the rat spinal dorsal horn. <i>Journal of Comparative Neurology</i> , 2001 , 430, 448-57	3.4	61
5	Mutation of E2F2 in mice causes enhanced T lymphocyte proliferation, leading to the development of autoimmunity. <i>Immunity</i> , 2001 , 15, 959-70	32.3	133
4	Immunocytochemical localization of the metabotropic glutamate receptor mGluR4a in the piriform cortex of the rat. <i>Journal of Comparative Neurology</i> , 2000 , 417, 263-274	3.4	16
3	ATR is required to complete meiotic recombination in mice		2

- 2 A chemical screen identifies a link between lipid metabolism and mRNA translation 1
- 1 USP7 couples DNA replication termination to mitotic entry 1