Rachel Codd

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/3690269/rachel-codd-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

82 26 48 2,540 g-index h-index citations papers 5.16 2,743 5.1 90 L-index avg, IF ext. citations ext. papers

| # | Paper | IF | Citations |
|----|--|------------------|-----------|
| 82 | Adding a diazo-transfer reagent to culture to generate secondary metabolite probes for click chemistry <i>Methods in Enzymology</i> , 2022 , 665, 49-71 | 1.7 | |
| 81 | Siderophores and iron transport 2021 , | | O |
| 80 | Directing macrocyclic architecture using iron(III)-, gallium(III)-, or zirconium(IV)-assisted ring closure of linear dimeric endo-hydroxamic acid ligands. <i>Journal of Inorganic Biochemistry</i> , 2021 , 216, 111337 | 4.2 | 1 |
| 79 | Azido-Desferrioxamine Siderophores as Functional Click-Chemistry Probes Generated in Culture upon Adding a Diazo-Transfer Reagent. <i>ChemBioChem</i> , 2020 , 21, 1433-1445 | 3.8 | 3 |
| 78 | Improved Access to Linear Tetrameric Hydroxamic Acids with Potential as Radiochemical Ligands for Zirconium(iv)-89 PET Imaging. <i>Australian Journal of Chemistry</i> , 2020 , | 1.2 | 4 |
| 77 | Immobilized Metal Affinity Chromatography as a Drug Discovery Platform for Metalloenzyme Inhibitors. <i>Journal of Medicinal Chemistry</i> , 2020 , 63, 12116-12127 | 8.3 | 1 |
| 76 | -Hydroxamic Acid Monomers for the Assembly of a Suite of Non-native Dimeric Macrocyclic Siderophores Using Metal-Templated Synthesis. <i>Inorganic Chemistry</i> , 2019 , 58, 13591-13603 | 5.1 | 4 |
| 75 | Preparation of a Dithiol-Reactive Probe for PET Imaging of Cell Death. <i>Methods in Molecular Biology</i> , 2019 , 1967, 295-304 | 1.4 | 3 |
| 74 | Analogues of desferrioxamine B (DFOB) with new properties and new functions generated using precursor-directed biosynthesis. <i>BioMetals</i> , 2019 , 32, 395-408 | 3.4 | 2 |
| 73 | Exploring hydroxamic acid inhibitors of HDAC1 and HDAC2 using small molecule tools and molecular or homology modelling. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2019 , 29, 2581-2586 | 2.9 | 1 |
| 72 | Engineering a cleavable disulfide bond into a natural product siderophore using precursor-directed biosynthesis. <i>Chemical Communications</i> , 2018 , 54, 9813-9816 | 5.8 | 8 |
| 71 | Fluorinated Analogues of Desferrioxamine B from Precursor-Directed Biosynthesis Provide New Insight into the Capacity of DesBCD. <i>ACS Chemical Biology</i> , 2018 , 13, 2456-2471 | 4.9 | 8 |
| 70 | Advances in the Chemical Biology of Desferrioxamine B. ACS Chemical Biology, 2018, 13, 11-25 | 4.9 | 38 |
| 69 | Rubik Cube of Siderophore Assembly Established from Mixed-Substrate Precursor-Directed Biosynthesis. <i>ACS Omega</i> , 2018 , 3, 18160-18169 | 3.9 | 4 |
| 68 | The chemical biology and coordination chemistry of putrebactin, avaroferrin, bisucaberin, and alcaligin. <i>Journal of Biological Inorganic Chemistry</i> , 2018 , 23, 969-982 | 3.7 | 10 |
| 67 | Octadentate Zirconium(IV)-Loaded Macrocycles with Varied Stoichiometry Assembled From Hydroxamic Acid Monomers using Metal-Templated Synthesis. <i>Inorganic Chemistry</i> , 2017 , 56, 3719-372 | 8 ^{5.1} | 21 |
| 66 | Adamantyl- and other polycyclic cage-based conjugates of desferrioxamine B (DFOB) for treating iron-mediated toxicity in cell models of Parkinson's disease. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2017 , 27, 1698-1704 | 2.9 | 9 |

(2012-2017)

| 65 | Analogues of desferrioxamine B designed to attenuate iron-mediated neurodegeneration: synthesis, characterisation and activity in the MPTP-mouse model of Parkinson's disease. Metallomics, 2017, 9, 852-864 | 4.5 | 20 | |
|----|---|--------------------|----|--|
| 64 | Exploiting the biosynthetic machinery of Streptomyces pilosus to engineer a water-soluble zirconium(iv) chelator. <i>Organic and Biomolecular Chemistry</i> , 2017 , 15, 5719-5730 | 3.9 | 25 | |
| 63 | Reverse Biosynthesis: Generating Combinatorial Pools of Drug Leads from Enzyme-Mediated Fragmentation of Natural Products. <i>ChemBioChem</i> , 2017 , 18, 368-373 | 3.8 | 4 | |
| 62 | Dimeric and trimeric homo- and heteroleptic hydroxamic acid macrocycles formed using mixed-ligand Fe(III)-based metal-templated synthesis. <i>Journal of Inorganic Biochemistry</i> , 2017 , 177, 344 | -3 1 57 | 6 | |
| 61 | Simultaneous biosynthesis of putrebactin, avaroferrin and bisucaberin by Shewanella putrefaciens and characterisation of complexes with iron(III), molybdenum(VI) or chromium(V). <i>Journal of Inorganic Biochemistry</i> , 2016 , 162, 207-215 | 4.2 | 23 | |
| 60 | Mixing Up the Pieces of the Desferrioxamine B Jigsaw Defines the Biosynthetic Sequence Catalyzed by DesD. <i>ACS Chemical Biology</i> , 2016 , 11, 1452-62 | 4.9 | 21 | |
| 59 | The BET bromodomain inhibitor exerts the most potent synergistic anticancer effects with quinone-containing compounds and anti-microtubule drugs. <i>Oncotarget</i> , 2016 , 7, 79217-79232 | 3.3 | 14 | |
| 58 | Hands-On Approach to Structure Activity Relationships: The Synthesis, Testing, and Hansch Analysis of a Series of Acetylcholineesterase Inhibitors. <i>Journal of Chemical Education</i> , 2015 , 92, 1745-1750 | 2.4 | 3 | |
| 57 | Forward and reverse (retro) iron(III) or gallium(III) desferrioxamine E and ring-expanded analogues prepared using metal-templated synthesis from endo-hydroxamic acid monomers. <i>Inorganic Chemistry</i> , 2015 , 54, 3573-83 | 5.1 | 12 | |
| 56 | The resolution of two clinical agents, bleomycin and desferrioxamine B, from a Streptomyces verticillus fermentation mixture using multi-dimensional immobilised metal ion affinity chromatography. <i>RSC Advances</i> , 2015 , 5, 3443-3453 | 3.7 | 8 | |
| 55 | Isolation of doxorubicin from a bacterial culture using immobilised metal ion affinity chromatography. <i>RSC Advances</i> , 2015 , 5, 46437-46442 | 3.7 | 3 | |
| 54 | Unsaturated macrocyclic dihydroxamic acid siderophores produced by Shewanella putrefaciens using precursor-directed biosynthesis. <i>ACS Chemical Biology</i> , 2014 , 9, 945-56 | 4.9 | 32 | |
| 53 | Dinuclear [(V(V)O(putrebactin))2(EOCH3)2] formed in solution as established from LC-MS measurements using 50V-enriched V2O5. <i>Inorganic Chemistry</i> , 2014 , 53, 5852-61 | 5.1 | 10 | |
| 52 | Solution species of Fe(III), Ga(III), In(III) or Ln(III) and suberodihydroxamic acid from electrospray ionization mass spectrometry. <i>RSC Advances</i> , 2013 , 3, 16051 | 3.7 | 2 | |
| 51 | The variable hydroxamic acid siderophore metabolome of the marine actinomycete Salinispora tropica CNB-440. <i>Metallomics</i> , 2013 , 5, 1519-28 | 4.5 | 36 | |
| 50 | Lipophilic adamantyl- or deferasirox-based conjugates of desferrioxamine B have enhanced neuroprotective capacity: implications for Parkinson disease. <i>Free Radical Biology and Medicine</i> , 2013 , 60, 147-56 | 7.8 | 26 | |
| 49 | Amide-based derivatives of Falanine hydroxamic acid as histone deacetylase inhibitors: attenuation of potency through resonance effects. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2012 , 22, 6200-4 | 2.9 | 5 | |
| 48 | Coordinate-bond-dependent solid-phase organic synthesis of biotinylated desferrioxamine B: a new route for metal-specific probes. <i>Chemical Communications</i> , 2012 , 48, 2003-5 | 5.8 | 5 | |

| 47 | Proteomics of Pseudomonas aeruginosa Australian epidemic strain 1 (AES-1) cultured under conditions mimicking the cystic fibrosis lung reveals increased iron acquisition via the siderophore pyochelin. <i>Journal of Proteome Research</i> , 2012 , 11, 776-95 | 5.6 | 33 |
|----|---|---------------------|-----|
| 46 | Directing the biosynthesis of putrebactin or desferrioxamine B in Shewanella putrefaciens through the upstream inhibition of ornithine decarboxylase. <i>Chemistry and Biodiversity</i> , 2012 , 9, 1880-90 | 2.5 | 25 |
| 45 | Copper(II)-based metal affinity chromatography for the isolation of the anticancer agent bleomycin from Streptomyces verticillus culture. <i>Journal of Inorganic Biochemistry</i> , 2012 , 115, 198-203 | 4.2 | 25 |
| 44 | Analytical-scale purification of trichostatin A from bacterial culture in a single step and with high selectivity using immobilised metal affinity chromatography. <i>RSC Advances</i> , 2012 , 2, 333-337 | 3.7 | 10 |
| 43 | Complexes formed in solution between vanadium(IV)/(V) and the cyclic dihydroxamic acid putrebactin or linear suberodihydroxamic acid. <i>Inorganic Chemistry</i> , 2011 , 50, 5978-89 | 5.1 | 18 |
| 42 | Cold adaptation of the mononuclear molybdoenzyme periplasmic nitrate reductase from the Antarctic bacterium Shewanella gelidimarina. <i>Biochemical and Biophysical Research Communications</i> , 2011 , 414, 783-8 | 3.4 | 10 |
| 41 | Comparing the potential renal protective activity of desferrioxamine B and the novel chelator desferrioxamine B-N-(3-hydroxyadamant-1-yl)carboxamide in a cell model of myoglobinuria. <i>Biochemical Journal</i> , 2011 , 435, 669-77 | 3.8 | 11 |
| 40 | The many faces of the adamantyl group in drug design. <i>European Journal of Medicinal Chemistry</i> , 2011 , 46, 1949-63 | 6.8 | 198 |
| 39 | The periplasmic nitrate reductase in Shewanella: the resolution, distribution and functional implications of two NAP isoforms, NapEDABC and NapDAGHB. <i>Microbiology (United Kingdom)</i> , 2010 , 156, 302-312 | 2.9 | 59 |
| 38 | Resolution of two native monomeric 90kDa nitrate reductase active proteins from Shewanella gelidimarina and the sequence of two napA genes. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 398, 13-8 | 3.4 | 15 |
| 37 | Structural diversity of complexes between Cu(II) or Ni(II) and endocyclic oxygen- or nitrogen-containing ligands: synthesis, X-ray structure determinations and circular dichroism spectra. <i>CrystEngComm</i> , 2010 , 12, 4217 | 3.3 | 10 |
| 36 | Conjugates of desferrioxamine B (DFOB) with derivatives of adamantane or with orally available chelators as potential agents for treating iron overload. <i>Journal of Medicinal Chemistry</i> , 2010 , 53, 1370-6 | 3 <mark>8</mark> .3 | 41 |
| 35 | Cloning, expression, purification and crystallization of dihydrodipicolinate synthase from the psychrophile Shewanella benthica. <i>Acta Crystallographica Section F: Structural Biology Communications</i> , 2010 , 66, 1511-6 | | 9 |
| 34 | Zn(II)-dependent histone deacetylase inhibitors: suberoylanilide hydroxamic acid and trichostatin A. <i>International Journal of Biochemistry and Cell Biology</i> , 2009 , 41, 736-9 | 5.6 | 98 |
| 33 | 2,5-Dioxopyrrolidin-1-yl adamantane-1-carboxyl-ate. <i>Acta Crystallographica Section E: Structure Reports Online</i> , 2009 , 65, o1740-1 | | 3 |
| 32 | Methyl 3-[(1-adamantylcarbon-yloxy)amino-carbon-yl]propanoate. <i>Acta Crystallographica Section E: Structure Reports Online</i> , 2009 , 65, o1742-3 | | 2 |
| 31 | Chromium in Cancer and Dietary Supplements. <i>Biological Magnetic Resonance</i> , 2009 , 551-579 | 0.5 | 6 |
| 30 | Immobilised metal affinity chromatography for the capture of hydroxamate-containing siderophores and other Fe(III)-binding metabolites directly from bacterial culture supernatants. <i>Analyst, The</i> , 2008 , 133, 877-80 | 5 | 25 |

(1999-2008)

| 29 | Studies of iron-uptake mechanisms in two bacterial species of the shewanella genus adapted to middle-range (Shewanella putrefaciens) or antarctic (Shewanella gelidimarina) temperatures. <i>Chemistry and Biodiversity</i> , 2008 , 5, 2113-23 | 2.5 | 16 |
|----|--|------|-----|
| 28 | Traversing the coordination chemistry and chemical biology of hydroxamic acids. <i>Coordination Chemistry Reviews</i> , 2008 , 252, 1387-1408 | 23.2 | 209 |
| 27 | Life on Earth. Extremophiles Continue to Move the Goal Posts. Environmental Chemistry, 2006, 3, 77 | 3.2 | 39 |
| 26 | Aqua[bis(2-pyridylmethyl)amine][chelidonato(1.5]]copper(II) chelidonate(0.5]]monohydrate. <i>Acta Crystallographica Section E: Structure Reports Online</i> , 2006 , 62, m3582-m3584 | | 9 |
| 25 | Chromium(V) complexes generated in Arthrobacter oxydans by simulation analysis of EPR spectra. <i>Journal of Inorganic Biochemistry</i> , 2006 , 100, 1827-33 | 4.2 | 22 |
| 24 | Chromium(V) complexes of hydroxamic acids: formation, structures, and reactivities. <i>Inorganic Chemistry</i> , 2005 , 44, 2934-43 | 5.1 | 34 |
| 23 | Coordination modes between copper(II) and N-acetylneuraminic (sialic) acid from a 2D-simulation analysis of EPR spectra. Implications for copper mediation of sialoglycoconjugate chemistry relevant to human biology. <i>Inorganic Chemistry</i> , 2005 , 44, 2531-43 | 5.1 | 14 |
| 22 | Metalloglycomics: a new perspective upon competitive metal-carbohydrate binding using EPR spectroscopy. <i>Chemical Communications</i> , 2004 , 2653-5 | 5.8 | 9 |
| 21 | X-ray absorption spectroscopic studies of chromium(V/IV/III)- 2-ethyl-2-hydroxybutanoato(2-/1-) complexes. <i>Inorganic Chemistry</i> , 2004 , 43, 1046-55 | 5.1 | 32 |
| 20 | Sialoglycoprotein and carbohydrate complexes in chromium toxicity. <i>Current Opinion in Chemical Biology</i> , 2003 , 7, 213-9 | 9.7 | 59 |
| 19 | Oxochromium(V) species formed with 2,3-dehydro-2-deoxy-N-acetylneuraminic or N-acetylneuraminic (sialic) acids: an in vitro model system of oxochromium(V) species potentially stabilized in the respiratory tract upon inhalation of carcinogenic chromium(VI) compounds. | 4 | 23 |
| 18 | Chemical Research in Toxicology, 2003, 16, 881-92 Pulsed ELDOR spectroscopy of the Mo(V)/Fe(III) state of sulfite oxidase prepared by one-electron reduction with Ti(III) citrate. <i>Journal of Biological Inorganic Chemistry</i> , 2002, 7, 338-50 | 3.7 | 45 |
| 17 | A potential synthon for models of vanadium haloperoxidase: (3,5-dimethylpyrazole)bis[2-hydroxy-2-methylbutanoato(1月oxovanadium(IV). <i>Acta Crystallographica Section E: Structure Reports Online</i> , 2002 , 58, m737-m739 | | 3 |
| 16 | Studies on the genotoxicity of chromium: from the test tube to the cell. <i>Coordination Chemistry Reviews</i> , 2001 , 216-217, 537-582 | 23.2 | 162 |
| 15 | Chromium(V)-sialic (neuraminic) acid species are formed from mixtures of chromium(VI) and saliva. <i>Journal of the American Chemical Society</i> , 2001 , 123, 11799-800 | 16.4 | 29 |
| 14 | Characterization and X-ray absorption spectroscopic studies of bis[quinato(2-)]oxochromate(V). <i>Inorganic Chemistry</i> , 2000 , 39, 990-7 | 5.1 | 32 |
| 13 | Competition between 1,2-Diol and 2-Hydroxy Acid Coordination in Cr(V)-Quinic Acid Complexes: Implications for Stabilization of Cr(V) Intermediates of Relevance to Cr(VI)-Induced Carcinogenesis. <i>Journal of the American Chemical Society</i> , 1999 , 121, 7864-7876 | 16.4 | 46 |
| 12 | In vitro plasmid DNA cleavage by chromium(V) and -(IV) 2-hydroxycarboxylato complexes. <i>Chemical Research in Toxicology</i> , 1999 , 12, 371-81 | 4 | 51 |

| 11 | Stability and Ligand Exchange Reactions of Chromium(IV) Carboxylato Complexes in Aqueous Solutions1. <i>Inorganic Chemistry</i> , 1997 , 36, 5440-5448 | 5.1 | 57 |
|----|---|------|-----|
| 10 | Synthesis, Characterization, and Crystal Structure of a Vanadium(V)-Vanadium(IV)-Vanadium(V) Mixed-Valence Trinuclear Complex with a Tridentate Quinato Ligand, (NH4)2{[VV(O)2]2[VIV(O)](.mu(-)-quinato(3-))2}.cntdot.H2O. <i>Inorganic Chemistry</i> , 1995 , 34, 877-882 | 5.1 | 46 |
| 9 | EPR characterisation of the CrV intermediates in the CrVI/V oxidations of organic substrates and of relevance to Cr-induced cancers. <i>Journal of the Chemical Society, Faraday Transactions</i> , 1995 , 91, 1207 | | 81 |
| 8 | Electrochemical investigations of poplar, spinach, cucumber, and parsley plastocyanins at conventional and microsized carbon electrodes. <i>Inorganic Chemistry</i> , 1992 , 31, 5007-5014 | 5.1 | 22 |
| 7 | The expression, purification and crystallization of the epsilon subunit of the F1 portion of the ATPase of Escherichia coli. <i>Journal of Molecular Biology</i> , 1992 , 228, 306-9 | 6.5 | 4 |
| 6 | Measurement of (1 -r3),(1 -r4)-£D-glucan in barley and oats: A streamlined enzymic procedure. <i>Journal of the Science of Food and Agriculture</i> , 1991 , 55, 303-312 | 4.3 | 313 |
| 5 | Resonance Raman spectra of plastocyanin and pseudoazurin: evidence for conserved cysteine ligand conformations in cupredoxins (blue copper proteins). <i>Biochemistry</i> , 1991 , 30, 10904-13 | 3.2 | 72 |
| 4 | Measurement of ⊞mylase in cereal flours and commercial enzyme preparations. <i>Journal of Cereal Science</i> , 1989 , 9, 17-33 | 3.8 | 128 |
| 3 | Pyrolysis and mesophase formation from sucrose. <i>Carbon</i> , 1988 , 26, 553-558 | 10.4 | 5 |
| 2 | Chromium in Biology: Toxicology and Nutritional Aspects145-250 | | 4 |
| 1 | New Applications of Immobilized Metal Ion Affinity Chromatography in Chemical Biology1-35 | | 2 |