

Ronald L Chandler

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/370925/publications.pdf>

Version: 2024-02-01

15
papers

648
citations

1040018

9
h-index

996954

15
g-index

16
all docs

16
docs citations

16
times ranked

1189
citing authors

#	ARTICLE	IF	CITATIONS
1	SWI/SNF Antagonism of PRC2 Mediates Estrogen-Induced Progesterone Receptor Expression. <i>Cells</i> , 2022, 11, 1000.	4.1	12
2	AP-1 Subunit JUNB Promotes Invasive Phenotypes in Endometriosis. <i>Reproductive Sciences</i> , 2022, 29, 3266-3277.	2.5	5
3	Endometrial Epithelial ARID1A Is Required for Uterine Immune Homeostasis during Early Pregnancy. <i>International Journal of Molecular Sciences</i> , 2022, 23, 6067.	4.1	1
4	Co-existing TP53 and ARID1A mutations promote aggressive endometrial tumorigenesis. <i>PLoS Genetics</i> , 2021, 17, e1009986.	3.5	24
5	A mouse model of endometriosis mimicking the natural spread of invasive endometrium. <i>Human Reproduction</i> , 2020, 35, 58-69.	0.9	24
6	SWI/SNF inactivation in the endometrial epithelium leads to loss of epithelial integrity. <i>Human Molecular Genetics</i> , 2020, 29, 3412-3430.	2.9	11
7	ARID1A Mutations Promote P300-Dependent Endometrial Invasion through Super-Enhancer Hyperacetylation. <i>Cell Reports</i> , 2020, 33, 108366.	6.4	36
8	Lgr5-positive endothelial progenitor cells occupy a tumor and injury prone niche in the kidney vasa recta. <i>Stem Cell Research</i> , 2020, 46, 101849.	0.7	2
9	ATAC-seq normalization method can significantly affect differential accessibility analysis and interpretation. <i>Epigenetics and Chromatin</i> , 2020, 13, 22.	3.9	49
10	ARID1A and PI3-kinase pathway mutations in the endometrium drive epithelial transdifferentiation and collective invasion. <i>Nature Communications</i> , 2019, 10, 3554.	12.8	96
11	The SWI/SNF BAF-A complex is essential for neural crest development. <i>Developmental Biology</i> , 2016, 411, 15-24.	2.0	39
12	Global gene expression profiling of a mouse model of ovarian clear cell carcinoma caused by ARID1A and PIK3CA mutations implicates a role for inflammatory cytokine signaling. <i>Genomics Data</i> , 2015, 5, 329-332.	1.3	4
13	Coexistent ARID1A and PIK3CA mutations promote ovarian clear-cell tumorigenesis through pro-tumorigenic inflammatory cytokine signalling. <i>Nature Communications</i> , 2015, 6, 6118.	12.8	247
14	Characterization of a Brg1 hypomorphic allele demonstrates that genetic and biochemical activity are tightly correlated. <i>Epigenetics</i> , 2014, 9, 249-256.	2.7	1
15	ARID1a-DNA Interactions Are Required for Promoter Occupancy by SWI/SNF. <i>Molecular and Cellular Biology</i> , 2013, 33, 265-280.	2.3	97