
Simon J Tabrett

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3711317/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Mixed culture purple phototrophic bacteria is an effective fishmeal replacement in aquaculture. Water Research X, 2019, 4, 100031.	2.8	80
2	Comparison of faecal collection methods and diet acclimation times for the measurement of digestibility coefficients in barramundi (<i>Lates calcarifer</i>). Aquaculture Nutrition, 2015, 21, 248-255.	1.1	28
3	An analysis of the effect of diet and genotype on protein and energy utilization by the black tiger shrimp, <i>Penaeus monodon</i> - why do genetically selected shrimp grow faster?. Aquaculture Nutrition, 2013, 19, 128-138.	1.1	25
4	The use of passive acoustics to measure feed consumption by Penaeus monodon (giant tiger prawn) in cultured systems. Aquacultural Engineering, 2013, 57, 38-47.	1.4	40
5	The purine nucleotides guanine, adenine and inosine are a dietary requirement for optimal growth of black tiger prawn, P. monodon. Aquaculture, 2013, 408-409, 100-105.	1.7	13

6 Dietary nucleotides are semi-essential nutrients for optimal growth of black tiger shrimp (Penaeus) Tj ETQq0 0 0 rgBT/Overlock 10 Tf 50

7	Digestibility of Lupinus albus lupin meals in barramundi (Lates calcarifer). Aquaculture, 2012, 364-365, 1-5.	1.7	14
8	An assessment of cereal grains and other starch sources in diets for barramundi (Lates calcarifer) - implications for nutritional and functional qualities of extruded feeds. Aquaculture Nutrition, 2012, 18, 388-399.	1.1	60
9	Do formulated feeds for juvenile Panulirus ornatus lobsters require dietary cholesterol supplementation?. Aquaculture, 2010, 307, 241-246.	1.7	14

 $10 \qquad \text{Effect of background colour on the distribution of astaxanthin in black tiger prawn (Penaeus) Tj ETQq0 0 0 rgBT /Overlock 10, Tf 50 382 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 = 1.7 =$

11	Digestibility of lupin kernel meals in feeds for the black tiger shrimp, Penaeus monodon. Aquaculture, 2007, 264, 353-362.	1.7	31
12	Growth response of the black tiger shrimp, Penaeus monodon fed diets containing different lupin cultivars. Aquaculture, 2007, 269, 436-446.	1.7	19
13	Response of the black tiger shrimp, Penaeus monodon to feed containing the lupin alkaloid, gramine. Aquaculture, 2007, 272, 556-563.	1.7	13
14	The efficacy of ingredients included in shrimp feeds to stimulate intake. Aquaculture Nutrition, 2005, 11, 263-272.	1.1	69
15	Water immersion time reduces the preference of juvenile tropical spiny lobster Panulirus ornatus for pelleted dry feeds and fresh mussel. Aquaculture Nutrition, 2005, 11, 415-426.	1.1	31
16	A novel method of collecting fecal samples from spiny lobsters. Aquaculture, 2005, 243, 269-272.	1.7	26
17	Evidence of a growth factor in some crustacean-based feed ingredients in diets for the giant tiger shrimp Penaeus monodon. Aquaculture, 2005, 250, 377-390.	1.7	34
18	The effect of dietary protein on the growth and survival of the shrimp, Penaeus monodon in outdoor tanks. Aquaculture Nutrition, 2004, 10, 15-23.	1.1	19

Simon J Tabrett

#	Article	IF	CITATIONS
19	Accurate measurement of in vivo digestibility of shrimp feeds. Aquaculture, 2004, 232, 563-580.	1.7	49
20	Development of a pelleted feed for juvenile tropical spiny lobster (Panulirus ornatus): response to dietary protein and lipid. Aquaculture Nutrition, 2003, 9, 231-237.	1.1	34
21	The effect of feeding frequency on water quality and growth of the black tiger shrimp (Penaeus) Tj ETQq1 1 0.784	314 rgBT	Qverlock 10
22	Cholesterol requirement of subadult black tiger shrimpPenaeus monodon(Fabricius). Aquaculture Research, 2001, 32, 399-405.	0.9	24
23	Relationships between the weight and chemical composition of exuvia and whole body of the black tiger prawn, Penaeus monodon. Aquaculture, 1994, 119, 249-258.	1.7	10