

# Anne Winther-Larsen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3718795/publications.pdf>

Version: 2024-02-01

25  
papers

375  
citations

858243

12  
h-index

939365

18  
g-index

25  
all docs

25  
docs citations

25  
times ranked

687  
citing authors

#	ARTICLE	IF	CITATIONS
1	Biological variation of serum neurofilament light chain. <i>Clinical Chemistry and Laboratory Medicine</i> , 2022, 60, 569-575.	1.4	19
2	New Insights in Coagulation and Fibrinolysis in Patients with Primary Brain Cancer: A Systematic Review. <i>Seminars in Thrombosis and Hemostasis</i> , 2022, 48, 323-337.	1.5	3
3	Inflammation-scores as prognostic markers of overall survival in lung cancer: a register-based study of 6,210 Danish lung cancer patients. <i>BMC Cancer</i> , 2022, 22, 63.	1.1	14
4	Hyperfibrinolysis in Patients with Solid Malignant Neoplasms: A Systematic Review. <i>Seminars in Thrombosis and Hemostasis</i> , 2021, 47, 581-588.	1.5	11
5	Hyponatremia as a prognostic factor in non-small cell lung cancer: a systematic review and meta-analysis. <i>Translational Lung Cancer Research</i> , 2021, 10, 651-661.	1.3	8
6	Hyponatremia in lung cancer: Incidence and prognostic value in a Danish population-based cohort study. <i>Lung Cancer</i> , 2021, 153, 42-48.	0.9	8
7	Pre-Treatment C-Reactive Protein Predicts Survival in Small Cell Lung Cancer Patients. <i>Onco</i> , 2021, 1, 114-123.	0.2	0
8	Pretreatment Albumin-to-Alkaline Phosphatase Ratio Is a Prognostic Marker in Lung Cancer Patients: A Registry-Based Study of 7077 Lung Cancer Patients. <i>Cancers</i> , 2021, 13, 6133.	1.7	9
9	Protein C deficiency; PROC gene variants in a Danish population. <i>Thrombosis Research</i> , 2020, 185, 153-159.	0.8	5
10	Neurofilament Light Chain as A Biomarker for Brain Metastases. <i>Cancers</i> , 2020, 12, 2852.	1.7	20
11	The ABO Locus is Associated with Increased Fibrin Network Formation in Patients with Stable Coronary Artery Disease. <i>Thrombosis and Haemostasis</i> , 2020, 120, 1248-1256.	1.8	7
12	Genomic Profiling of Circulating Tumor DNA Predicts Outcome and Demonstrates Tumor Evolution in ALK-Positive Non-Small Cell Lung Cancer Patients. <i>Cancers</i> , 2020, 12, 947.	1.7	20
13	Circulating miR-30b and miR-30c predict erlotinib response in EGFR-mutated non-small cell lung cancer patients. <i>Lung Cancer</i> , 2019, 135, 92-96.	0.9	22
14	Day-to-day and within-day biological variation of cell-free DNA. <i>EBioMedicine</i> , 2019, 49, 284-290.	2.7	49
15	Clinical impact of direct oral anticoagulant measuring in a real-life setting. <i>Thrombosis Research</i> , 2019, 175, 40-45.	0.8	8
16	EGFR Gene Polymorphism Predicts Improved Outcome in Patients With EGFR Mutation-positive Non-small cell Lung Cancer Treated With Erlotinib. <i>Clinical Lung Cancer</i> , 2019, 20, 161-166.e1.	1.1	13
17	Reference limits for GAD65 and IA-2 autoantibodies by chemiluminescence immunoassay in Northern European adults and children. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2019, 79, 123-125.	0.6	0
18	A method for treatment monitoring using circulating tumour DNA in cancer patients without targetable mutations. <i>Oncotarget</i> , 2018, 9, 31066-31076.	0.8	18

#	ARTICLE	IF	CITATIONS
19	<sup>18</sup> F-FDG PET/CT for Very Early Response Evaluation Predicts CT Response in Erlotinib-Treated Non-Small Cell Lung Cancer Patients: A Comparison of Assessment Methods. <i>Journal of Nuclear Medicine</i> , 2017, 58, 1931-1937.	2.8	16
20	Correlation between circulating mutant DNA and metabolic tumour burden in advanced non-small cell lung cancer patients. <i>British Journal of Cancer</i> , 2017, 117, 704-709.	2.9	45
21	Early Change in FDG-PET Signal and Plasma Cell-Free DNA Level Predicts Erlotinib Response in EGFR Wild-Type NSCLC Patients. <i>Translational Oncology</i> , 2016, 9, 505-511.	1.7	13
22	Metabolic tumor burden as marker of outcome in advanced EGFR wild-type NSCLC patients treated with erlotinib. <i>Lung Cancer</i> , 2016, 94, 81-87.	0.9	34
23	Evaluation of factors associated with loco-regional failure and survival in limited disease small cell lung cancer patients treated with chemoradiotherapy. <i>Acta Oncologica</i> , 2015, 54, 1574-1581.	0.8	9
24	Genetic polymorphism in the epidermal growth factor receptor gene predicts outcome in advanced non-small cell lung cancer patients treated with erlotinib. <i>Lung Cancer</i> , 2015, 90, 314-320.	0.9	13
25	EGFR CA repeat polymorphism predict clinical outcome in EGFR mutation positive NSCLC patients treated with erlotinib. <i>Lung Cancer</i> , 2014, 85, 435-441.	0.9	11