

# Sandra PÃ©rez-Torras

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3729316/publications.pdf>

Version: 2024-02-01

22  
papers

606  
citations

758635

12  
h-index

713013

21  
g-index

22  
all docs

22  
docs citations

22  
times ranked

1177  
citing authors

#	ARTICLE	IF	CITATIONS
1	Emerging Roles of Nucleoside Transporters. <i>Frontiers in Pharmacology</i> , 2018, 9, 606.	1.6	105
2	Who Is Who in Adenosine Transport. <i>Frontiers in Pharmacology</i> , 2018, 9, 627.	1.6	85
3	Nucleoside transporter proteins as biomarkers of drug responsiveness and drug targets. <i>Frontiers in Pharmacology</i> , 2015, 6, 13.	1.6	84
4	Ribonucleotide reductase is an effective target to overcome gemcitabine resistance in gemcitabine-resistant pancreatic cancer cells with dual resistant factors. <i>Journal of Pharmacological Sciences</i> , 2015, 127, 319-325.	1.1	45
5	Adenoviral-mediated overexpression of human equilibrative nucleoside transporter 1 (hENT1) enhances gemcitabine response in human pancreatic cancer. <i>Biochemical Pharmacology</i> , 2008, 76, 322-329.	2.0	40
6	Intestinal Nucleoside Transporters: Function, Expression, and Regulation. , 2018, 8, 1003-1017.		35
7	Connexin-26 Is a Key Factor Mediating Gemcitabine Bystander Effect. <i>Molecular Cancer Therapeutics</i> , 2011, 10, 505-517.	1.9	33
8	Aquaporin 3 (AQP3) participates in the cytotoxic response to nucleoside-derived drugs. <i>BMC Cancer</i> , 2012, 12, 434.	1.1	28
9	Concentrative nucleoside transporter 1 (hCNT1) promotes phenotypic changes relevant to tumor biology in a translocation-independent manner. <i>Cell Death and Disease</i> , 2013, 4, e648-e648.	2.7	26
10	Characterization of human pancreatic orthotopic tumor xenografts suitable for drug screening. <i>Cellular Oncology (Dordrecht)</i> , 2011, 34, 511-521.	2.1	23
11	Transportome Profiling Identifies Profound Alterations in Crohn's Disease Partially Restored by Commensal Bacteria. <i>Journal of Crohn's and Colitis</i> , 2016, 10, 850-859.	0.6	21
12	Human pancreatic cancer stem cells are sensitive to dual inhibition of IGF-IR and ErbB receptors. <i>BMC Cancer</i> , 2015, 15, 223.	1.1	16
13	New role of the human equilibrative nucleoside transporter 1 (hENT1) in Epithelial-mesenchymal transition in renal tubular cells. <i>Journal of Cellular Physiology</i> , 2012, 227, 1521-1528.	2.0	15
14	Role of the Transporter Regulator Protein (RS1) in the Modulation of Concentrative Nucleoside Transporters (CNTs) in Epithelia. <i>Molecular Pharmacology</i> , 2012, 82, 59-67.	1.0	12
15	Deficiency of perforin and hCNT1, a novel inborn error of pyrimidine metabolism, associated with a rapidly developing lethal phenotype due to multi-organ failure. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2019, 1865, 1182-1191.	1.8	8
16	Dexamethasone-Loaded Lipomers: Development, Characterization, and Skin Biodistribution Studies. <i>Pharmaceutics</i> , 2021, 13, 533.	2.0	7
17	The Physicochemical, Biopharmaceutical, and In Vitro Efficacy Properties of Freeze-Dried Dexamethasone-Loaded Lipomers. <i>Pharmaceutics</i> , 2021, 13, 1322.	2.0	7
18	Role of drug-dependent transporter modulation on the chemosensitivity of cholangiocarcinoma. <i>Oncotarget</i> , 2017, 8, 90185-90196.	0.8	6

#	ARTICLE	IF	CITATIONS
19	Pharmacogenomic analyzis of the responsiveness of gastrointestinal tumor cell lines to drug therapy: A transportome approach. Pharmacological Research, 2016, 113, 364-375.	3.1	4
20	FMS-like tyrosine kinase 3 (FLT3) modulates key enzymes of nucleotide metabolism implicated in cytarabine responsiveness in pediatric acute leukemia. Pharmacological Research, 2020, 151, 104556.	3.1	3
21	OncomiRs miR-106a and miR-17 negatively regulate the nucleoside-derived drug transporter hCNT1. Cellular and Molecular Life Sciences, 2021, 78, 7505-7518.	2.4	2
22	An Escape-Room about Krebs cycle prepared for Chemical Students. International Journal on Engineering, Science and Technology, 2022, 3, 155-164.	0.2	1