Greg E Gaski

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3734022/publications.pdf

Version: 2024-02-01

1307594 1281871 19 148 7 11 citations g-index h-index papers 20 20 20 203 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Computational evidence for an early, amplified systemic inflammation program in polytrauma patients with severe extremity injuries. PLoS ONE, 2019, 14, e0217577.	2.5	26
2	Early Immunologic Response in Multiply Injured Patients With Orthopaedic Injuries Is Associated With Organ Dysfunction. Journal of Orthopaedic Trauma, 2019, 33, 220-228.	1.4	21
3	Precision Medicine Applications to Manage Multiply Injured Patients With Orthopaedic Trauma. Journal of Orthopaedic Trauma, 2019, 33, S25-S29.	1.4	13
4	Tissue damage volume predicts organ dysfunction and inflammation after injury. Journal of Surgical Research, 2016, 202, 188-195.	1.6	11
5	The effect of pH versus base deficit on organ failure in trauma patients. Journal of Surgical Research, 2016, 200, 260-265.	1.6	11
6	Fixation Using Alternative Implants for the Treatment of Hip Fractures (FAITH-2): The Clinical Outcomes of a Multicenter 2 × 2 Factorial Randomized Controlled Pilot Trial in Young Femoral Neck Fracture Patients. Journal of Orthopaedic Trauma, 2020, 34, 524-532.	1.4	9
7	Early dynamic orchestration of immunologic mediators identifies multiply injured patients who are tolerant or sensitive to hemorrhage. Journal of Trauma and Acute Care Surgery, 2021, 90, 441-450.	2.1	8
8	Large-magnitude Pelvic and Retroperitoneal Tissue Damage Predicts Organ Failure. Clinical Orthopaedics and Related Research, 2016, 474, 1410-1416.	1.5	7
9	Base Deficit ≥Â6 within 24Âh of Injury is a risk factor for fracture nonunion in the polytraumatized patient. Injury, 2021, 52, 3271-3276.	1.7	6
10	Diagnosis and Management of Polytraumatized Patients With Severe Extremity Trauma. Journal of Orthopaedic Trauma, 2018, 32, S1-S6.	1.4	5
11	Shock volume: Patient-specific cumulative hypoperfusion predicts organ dysfunction in a prospective cohort of multiply injured patients. Journal of Trauma and Acute Care Surgery, 2018, 85, S84-S91.	2.1	5
12	Abdominopelvic bleed rate on admission CT correlates with mortality and transfusion needs in the setting of blunt pelvic fractures: a single institution pilot study. Emergency Radiology, 2019, 26, 37-44.	1.8	5
13	Retrograde intramedullary nailing below a hip arthroplasty prosthesis: a viable fixation option for periprosthetic and interprosthetic femur fractures. International Orthopaedics, 2020, 44, 2283-2289.	1.9	5
14	Cluster identification, selection, and description in cluster randomized crossover trials: the PREP-IT trials. Trials, 2020, 21, 712.	1.6	5
15	Fixation using Alternative Implants for the Treatment of Hip Fractures (FAITH-2): The Exploratory Health-Related Quality of Life and Patient-Reported Functional Outcomes of a Multi-Centre 2—Â2 Factorial Randomized Controlled Pilot Trial in Young Femoral Neck Fracture Patients. Injury, 2021, 52, 3051-3059.	1.7	5
16	Patient-Specific Precision Injury Signatures to Optimize Orthopaedic Interventions in Multiply Injured Patients (PRECISE STUDY). Journal of Orthopaedic Trauma, 2022, 36, S14-S20.	1.4	3
17	Fixation Using Alternative Implants for the Treatment of Hip Fractures. OTA International the Open Access Journal of Orthopaedic Trauma, 2020, 3, e66.	1.0	2
18	Risk factors for complications within 30 days of operatively fixed periprosthetic femur fractures. Journal of Clinical Orthopaedics and Trauma, 2022, , 101925.	1.5	1

#	Article	IF	CITATIONS
19	Polytraumatized patient lower extremity nonunion development: Raw data. Data in Brief, 2021, 37, 107244.	1.0	0