Zheng-Gang Ren

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3735932/publications.pdf

Version: 2024-02-01

43 papers

2,494 citations

257101 24 h-index 243296 44 g-index

45 all docs

45 docs citations

45 times ranked

3512 citing authors

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Guidelines for Diagnosis and Treatment of Primary Liver Cancer in China (2017 Edition). Liver Cancer, 2018, 7, 235-260. | 4.2 | 426 |
| 2 | Camrelizumab in patients with previously treated advanced hepatocellular carcinoma: a multicentre, open-label, parallel-group, randomised, phase 2 trial. Lancet Oncology, The, 2020, 21, 571-580. | 5.1 | 373 |
| 3 | 2019 Chinese clinical guidelines for the management of hepatocellular carcinoma: updates and insights. Hepatobiliary Surgery and Nutrition, 2020, 9, 452-463. | 0.7 | 267 |
| 4 | Coexpression of gene Oct4 and Nanog initiates stem cell characteristics in hepatocellular carcinoma and promotes epithelial-mesenchymal transition through activation of Stat3/Snail signaling. Journal of Hematology and Oncology, 2015, 8, 23. | 6.9 | 136 |
| 5 | Prognostic Significance of the Neutrophil-to-Lymphocyte Ratio in Primary Liver Cancer: A Meta-Analysis. PLoS ONE, 2014, 9, e96072. | 1.1 | 101 |
| 6 | Postoperative adjuvant arterial chemoembolization improves survival of hepatocellular carcinoma patients with risk factors for residual tumor: A retrospective control study. World Journal of Gastroenterology, 2004, 10, 2791. | 1.4 | 85 |
| 7 | Reactive cutaneous capillary endothelial proliferation in advanced hepatocellular carcinoma patients treated with camrelizumab: data derived from a multicenter phase 2 trial. Journal of Hematology and Oncology, 2020, 13, 47. | 6.9 | 84 |
| 8 | ID1 promotes hepatocellular carcinoma proliferation and confers chemoresistance to oxaliplatin by activating pentose phosphate pathway. Journal of Experimental and Clinical Cancer Research, 2017, 36, 166. | 3.5 | 79 |
| 9 | Transcatheter arterial chemoembolization combined with radiofrequency ablation delays tumor progression and prolongs overall survival in patients with intermediate (BCLC B) hepatocellular carcinoma. BMC Cancer, 2014, 14, 849. | 1.1 | 74 |
| 10 | Incomplete Radiofrequency Ablation Enhances Invasiveness and Metastasis of Residual Cancer of Hepatocellular Carcinoma Cell HCCLM3 via Activating \hat{l}^2 -Catenin Signaling. PLoS ONE, 2014, 9, e115949. | 1.1 | 58 |
| 11 | Critical appraisal of Chinese 2017 guideline on the management of hepatocellular carcinoma. Hepatobiliary Surgery and Nutrition, 2017, 6, 387-396. | 0.7 | 54 |
| 12 | Extracellular matrix collagen I promotes the tumor progression of residual hepatocellular carcinoma after heat treatment. BMC Cancer, 2018, 18, 901. | 1.1 | 49 |
| 13 | Suppressing DRP1-mediated mitochondrial fission and mitophagy increases mitochondrial apoptosis of hepatocellular carcinoma cells in the setting of hypoxia. Oncogenesis, 2020, 9, 67. | 2.1 | 46 |
| 14 | Astragaloside IV inhibits metastasis in hepatoma cells through the suppression of epithelial-mesenchymal transition via the Akt/GSK-3β/β-catenin pathway. Oncology Reports, 2017, 37, 1725-1735. | 1.2 | 40 |
| 15 | Increased matrix stiffness promotes tumor progression of residual hepatocellular carcinoma after insufficient heat treatment. Cancer Science, 2017, 108, 1778-1786. | 1.7 | 39 |
| 16 | Albumin to gamma-glutamyltransferase ratio as a prognostic indicator in intrahepatic cholangiocarcinoma after curative resection. Oncotarget, 2017, 8, 13293-13303. | 0.8 | 39 |
| 17 | Identifying Clonal Origin of Multifocal Hepatocellular Carcinoma and Its Clinical Implications. Clinical and Translational Gastroenterology, 2019, 10, e00006. | 1.3 | 36 |
| 18 | Maintenance of Stemness in Oxaliplatin-Resistant Hepatocellular Carcinoma Is Associated with Increased Autocrine of IGF1. PLoS ONE, 2014, 9, e89686. | 1.1 | 33 |

| # | Article | IF | Citations |
|----|---|-----|-----------|
| 19 | Activated hepatic stellate cells secrete periostin to induce stem cell-like phenotype of residual hepatocellular carcinoma cells after heat treatment. Scientific Reports, 2017, 7, 2164. | 1.6 | 33 |
| 20 | Camrelizumab plus oxaliplatinâ€based chemotherapy as firstâ€line therapy for advanced biliary tract cancer: A multicenter, phase 2 trial. International Journal of Cancer, 2021, 149, 1944-1954. | 2.3 | 32 |
| 21 | Dynamic Expression Patterns of Differential Proteins during Early Invasion of Hepatocellular Carcinoma. PLoS ONE, 2014, 9, e88543. | 1.1 | 29 |
| 22 | Herbal Compound <i>Songyou Yin</i> and Moderate Swimming Suppress Growth and Metastasis of Liver Cancer by Enhancing Immune Function. Integrative Cancer Therapies, 2016, 15, 368-375. | 0.8 | 29 |
| 23 | Identification of long noncoding RNA expression profile in oxaliplatin-resistant hepatocellular carcinoma cells. Gene, 2017, 596, 53-88. | 1.0 | 27 |
| 24 | Angiogenesis enhanced by treatment damage to hepatocellular carcinoma through the release of <scp>GDF</scp> 15. Cancer Medicine, 2018, 7, 820-830. | 1.3 | 27 |
| 25 | Goosecoid Promotes the Metastasis of Hepatocellular Carcinoma by Modulating the Epithelial-Mesenchymal Transition. PLoS ONE, 2014, 9, e109695. | 1.1 | 27 |
| 26 | Prognostic Nomogram Based on Histological Characteristics of Fibrotic Tumor Stroma in Patients Who Underwent Curative Resection for Intrahepatic Cholangiocarcinoma. Oncologist, 2018, 23, 1482-1493. | 1.9 | 26 |
| 27 | Intermittent hypoxia alleviates increased VEGF and pro†angiogenic potential in liver cancer cells. Oncology Letters, 2019, 18, 1831-1839. | 0.8 | 26 |
| 28 | Radiofrequency ablation following first-line transarterial chemoembolization for patients with unresectable hepatocellular carcinoma beyond the Milan criteria. BMC Gastroenterology, 2014, 14, 11. | 0.8 | 24 |
| 29 | An SCD1-dependent mechanoresponsive pathway promotes HCC invasion and metastasis through lipid metabolic reprogramming. Molecular Therapy, 2022, 30, 2554-2567. | 3.7 | 24 |
| 30 | Oxaliplatin and 5-fluorouracil hepatic infusion with lipiodolized chemoembolization in large hepatocellular carcinoma. World Journal of Gastroenterology, 2015, 21, 3970. | 1.4 | 19 |
| 31 | Aspirin in combination with TACE in treatment of unresectable HCC: a matched-pairs analysis. American Journal of Cancer Research, 2016, 6, 2109-2116. | 1.4 | 19 |
| 32 | The Rho GTPase Rnd1 inhibits epithelial–mesenchymal transition in hepatocellular carcinoma and is a favorable anti-metastasis target. Cell Death and Disease, 2018, 9, 486. | 2.7 | 18 |
| 33 | Hepatic Arterial Infusion Chemotherapy with Modified FOLFOX as an Alternative Treatment Option in Advanced Hepatocellular Carcinoma Patients with Failed or Unsuitability for Transarterial Chemoembolization. Academic Radiology, 2021, 28, S157-S166. | 1.3 | 16 |
| 34 | Albumin-to-alkaline phosphatase ratio as a predictor of tumor recurrence and prognosis in patients with early-stage hepatocellular carcinoma undergoing radiofrequency ablation as initial therapy. International Journal of Hyperthermia, 2021, 38, 1-10. | 1.1 | 16 |
| 35 | Comparison of transarterial chemoembolization with radiofrequency ablation for unresectable Barcelona Clinic Liver Cancer stage 0/A hepatocellular carcinoma: a propensity score matching. Journal of Gastroenterology and Hepatology (Australia), 2016, 31, 442-449. | 1.4 | 14 |
| 36 | <i>Clostridium perfringens</i> infection after transarterial chemoembolization for large hepatocellular carcinoma. World Journal of Gastroenterology, 2015, 21, 4397. | 1.4 | 14 |

| # | Article | IF | Citations |
|----|--|-----|-----------|
| 37 | The herbal compound Songyou Yin (SYY) inhibits hepatocellular carcinoma growth and improves survival in models of chronic fibrosis via paracrine inhibition of activated hepatic stellate cells. Oncotarget, 2015, 6, 40068-40080. | 0.8 | 12 |
| 38 | Norepinephrine-stimulated HSCs secrete sFRP1 to promote HCC progression following chronic stress via augmentation of a Wnt16B/ \hat{l}^2 -catenin positive feedback loop. Journal of Experimental and Clinical Cancer Research, 2020, 39, 64. | 3.5 | 9 |
| 39 | A new substage classification strategy for Barcelona Clinic Liver Cancer stage B patients with hepatocellular carcinoma. Journal of Gastroenterology and Hepatology (Australia), 2019, 34, 1984-1991. | 1.4 | 7 |
| 40 | A new prediction model for prognosis of patients with intermediate-stage HCC after conventional transarterial chemoembolization: an internally validated study. Journal of Cancer, 2019, 10, 6535-6542. | 1.2 | 6 |
| 41 | Over expression of hyaluronan promotes progression of HCC via CD44-mediated pyruvate kinase M2 nuclear translocation. American Journal of Cancer Research, 2016, 6, 509-21. | 1.4 | 6 |
| 42 | Prognostic significance of preoperative systemic immune-inflammation index in combined hepatocellular-cholangiocarcinoma. Cancer Biomarkers, 2021, 31, 1-15. | 0.8 | 5 |
| 43 | Development of a Prognostic Scoring System for Hepatocellular Carcinoma Patients With Main Portal Vein Tumor Thrombus Undergoing Conventional Transarterial Chemoembolization: An Analysis of 173 Patients. Frontiers in Oncology, 2021, 11, 671171. | 1.3 | 4 |