
José Pedreira

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3735947/publications.pdf Version: 2024-02-01



IOSÃO DEDDEIDA

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Integrating the control of helminths in dairy cattle: Deworming, rotational grazing and nutritional pellets with parasiticide fungi. Veterinary Parasitology, 2020, 278, 109038. | 1.8 | 14 |
| 2 | Effect of the Filamentous Fungus <i>Mucor circinelloides</i> On The Development of Eggs of the Rumen Fluke <i>Calicophoron daubneyi</i> (Paramphistomidae). Journal of Parasitology, 2017, 103, 199-206. | 0.7 | 6 |
| 3 | Feeding horses with industrially manufactured pellets with fungal spores to promote nematode integrated control. Veterinary Parasitology, 2016, 229, 37-44. | 1.8 | 22 |
| 4 | Prevalence of mixed trematode infections in an abattoir receiving cattle from northern Portugal and northâ€west Spain. Veterinary Record, 2011, 168, 408-408. | 0.3 | 50 |
| 5 | Field Evaluation for Anthelmintic-Resistant Ovine Gastrointestinal Nematodes by In Vitro and In Vivo Assays. Journal of Parasitology, 2008, 94, 925-928. | 0.7 | 27 |
| 6 | Immunodiagnosis of current fasciolosis in sheep naturally exposed to Fasciola hepatica by using a 2.9kDa recombinant protein. Veterinary Parasitology, 2007, 146, 46-49. | 1.8 | 14 |
| 7 | Assessment of climatic and orographic conditions on the infection by Calicophoron daubneyi and Dicrocoelium dendriticum in grazing beef cattle (NW Spain). Veterinary Parasitology, 2007, 149, 285-289. | 1.8 | 30 |
| 8 | Risk periods of infection by Calicophoron daubneyi (Digenea:Paramphistomidae) in cattle from oceanic climate areas. Parasitology Research, 2007, 101, 339-342. | 1.6 | 28 |
| 9 | A 2.9kDa Fasciola hepatica-recombinant protein based ELISA test for the detection of current-ovine fasciolosis trickle infected. Veterinary Parasitology, 2006, 137, 67-73. | 1.8 | 15 |
| 10 | Analysis of the IgG antibody response against Paramphistomidae trematoda in naturally infected cattle. Veterinary Parasitology, 2006, 140, 281-288. | 1.8 | 26 |
| 11 | Toxocara canis larvae viability after disinfectant—exposition. Parasitology Research, 2006, 99, 558-561. | 1.6 | 18 |
| 12 | Prevalences of gastrointestinal parasites in sheep and parasite-control practices in NW Spain. Preventive Veterinary Medicine, 2006, 75, 56-62. | 1.9 | 23 |
| 13 | Analysis of the humoral immune response to Oestrus ovis in ovine. Veterinary Parasitology, 2005, 134, 153-158. | 1.8 | 23 |
| 14 | Prevalence of natural ovine fasciolosis shown by demonstrating the presence of serum circulating antigens. Parasitology Research, 2003, 91, 328-331. | 1.6 | 20 |
| 15 | Time-course analysis of coproantigens in rats infected and challenged with Fasciola hepatica. Parasitology Research, 2002, 88, 568-573. | 1.6 | 12 |
| 16 | Influence of age and breed on natural bovine fasciolosis in an endemic area (Galicia, NW Spain). Veterinary Research Communications, 2002, 26, 361-370. | 1.6 | 34 |
| 17 | Effect of fasciolicides on the antigenaemia in sheep naturally infected with Fasciola hepatica. Parasitology Research, 2001, 87, 609-614. | 1.6 | 34 |