

Yutaka Takahashi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3737431/publications.pdf>

Version: 2024-02-01

14
papers

205
citations

1163117

8
h-index

1125743

13
g-index

16
all docs

16
docs citations

16
times ranked

272
citing authors

#	ARTICLE	IF	CITATIONS
1	A Pilot Study of Individualized Maximum Repeatable Dose (iMRD), a New Dose Finding System, of Weekly Gemcitabine for Patients With Metastatic Pancreas Cancer. <i>Pancreas</i> , 2005, 30, 206-210.	1.1	32
2	The angiogenic switch of human colon cancer occurs simultaneous to initiation of invasion. <i>Oncology Reports</i> , 2003, 10, 9-13.	2.6	31
3	Î±-Difluoromethylornithine induces apoptosis as well as anti-angiogenesis in the inhibition of tumor growth and metastasis in a human gastric cancer model. , 2000, 85, 243-247.		30
4	Angiogenesis of AFP producing gastric carcinoma: correlation with frequent liver metastasis and its inhibition by anti-AFP antibody. <i>Oncology Reports</i> , 2004, 11, 809-13.	2.6	18
5	Randomized phase II trial of first-line treatment with tailored irinotecan and S-1 therapy versus S-1 monotherapy for advanced or recurrent gastric carcinoma (JFMC31-0301). <i>Anti-Cancer Drugs</i> , 2011, 22, 576-583.	1.4	16
6	Î±-Difluoromethylornithine induces apoptosis as well as anti-angiogenesis in the inhibition of tumor growth and metastasis in a human gastric cancer model. <i>International Journal of Cancer</i> , 2000, 85, 243-247.	5.1	16
7	Multiparametric in situ mRNA Hybridization Analysis of Gastric Biopsies Predicts Lymph Node Metastasis in Patients with Gastric Carcinoma. <i>Japanese Journal of Cancer Research</i> , 2002, 93, 1258-1265.	1.7	12
8	Preoperative CEA and PPD values as prognostic factors for immunochemotherapy using PSK and 5-FU. <i>Anticancer Research</i> , 2005, 25, 1377-84.	1.1	10
9	Therapeutic approaches targeting tumor vasculature in gastrointestinal cancers. <i>Frontiers in Bioscience - Elite</i> , 2011, E3, 541-548.	1.8	9
10	A Randomized Phase II Clinical Trial of Tailored CPT-11 + S-1 vs S-1 in Patients with Advanced or Recurrent Gastric Carcinoma as the First Line Chemotherapy. <i>Japanese Journal of Clinical Oncology</i> , 2004, 34, 342-345.	1.3	8
11	Pilot Study of Low-Dose, Divided Maximum Tolerated Dose of CPT-11 in 21 Consecutive Patients with Metastatic Colorectal or Gastric Cancer. <i>Surgery Today</i> , 2004, 34, 246-250.	1.5	8
12	Chemotherapy under cachectic conditions and the possibility of cachexia-controlled chemotherapy. <i>Oncology Reports</i> , 2005, 14, 135-40.	2.6	6
13	Oral Administration of Uracil-Tegafur (UFT) Inhibits Liver Micrometastasis of Human Colon Cancer in an Orthotopic Nude Mouse Model and Its Early Detection System. <i>Surgery Today</i> , 2004, 34, 855-9.	1.5	4
14	Tailored-dose chemotherapy with gemcitabine and irinotecan in patients with platinum-refractory/resistant ovarian or primary peritoneal cancer: a phase II trial. <i>Journal of Gynecologic Oncology</i> , 2021, 32, e8.	2.2	3