

# Michael MÄGller

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3745291/publications.pdf>

Version: 2024-02-01

102  
papers

3,002  
citations

172207  
h-index

189595  
g-index

103  
all docs

103  
docs citations

103  
times ranked

2416  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Lectotypifications of four names in the family Gesneriaceae. <i>Adansonia</i> , 2022, 44, .   | 0.1 | 0         |
| 2  | One transfer to <i>Primulina</i> (Gesneriaceae) and amended descriptions for <i>P. crassifolia</i> and <i>P. quanbaensis</i> from northern Vietnam. <i>Nordic Journal of Botany</i> , 2022, 2022, .                               | 0.2 | 1         |
| 3  | The first genome for the Cape Primrose <i>Streptocarpus rexii</i> (Gesneriaceae), a model plant for studying meristem-driven shoot diversity. <i>Plant Direct</i> , 2022, 6, e388.  | 0.8 | 4         |
| 4  | Testing the Complete Plastome for Species Discrimination, Cryptic Species Discovery and Phylogenetic Resolution in Cephalotaxus (Cephalotaxaceae). <i>Frontiers in Plant Science</i> , 2022, 13, .                                | 1.7 | 16        |
| 5  | Mix and match: Patchwork domain evolution of the land plant-specific Ca <sup>2+</sup> -permeable mechanosensitive channel MCA. <i>PLoS ONE</i> , 2021, 16, e0249735.  | 1.1 | 10        |
| 6  | (2809) Proposal to conserve the name <i>Tetraphyllum</i> Griff. ex C.B. Clarke (Gesneriaceae) against <i>Tetraphyllum</i> Hosius & Marck (published as fossil <i>Magnoliophyta</i> ). <i>Taxon</i> , 2021, 70, 437-438.           | 0.4 | 1         |
| 7  | Phylogenomics of Gesneriaceae using targeted capture of nuclear genes. <i>Molecular Phylogenetics and Evolution</i> , 2021, 157, 107068.  | 1.2 | 46        |
| 8  | Sectional polyphyly and morphological homoplasy in Southeast Asian Cyrtandra (Gesneriaceae): consequences for the taxonomy of a mega-diverse genus. <i>Plant Systematics and Evolution</i> , 2021, 307, 1.                        | 0.3 | 4         |
| 9  | Repeated intercontinental migrations and recurring hybridizations characterise the evolutionary history of yew ( <i>Taxus</i> L.). <i>Molecular Phylogenetics and Evolution</i> , 2020, 153, 106952.                              | 1.2 | 10        |
| 10 | QTL dissection of floral traits in <i>Streptocarpus</i> (Gesneriaceae). <i>Euphytica</i> , 2020, 216, 1.  | 0.6 | 4         |
| 11 | Virus-induced Gene Silencing in <i>Streptocarpus rexii</i> (Gesneriaceae). <i>Molecular Biotechnology</i> , 2020, 62, 317-325.  | 1.3 | 6         |
| 12 | The Family Placement of Cyrtandromoea. <i>Systematic Botany</i> , 2019, 44, 616-630.  | 0.2 | 7         |
| 13 | <i>Boeica konchurangensis</i> sp. nov. (Gesneriaceae) from Gia Lai plateau, Vietnam. <i>Nordic Journal of Botany</i> , 2019, 37, .  | 0.2 | 1         |
| 14 | Gene duplication and relaxation from selective constraints of GCYC genes correlated with various floral symmetry patterns in Asiatic Gesneriaceae tribe Trichosporeae. <i>PLoS ONE</i> , 2019, 14, e0210054.                      | 1.1 | 5         |
| 15 | The Living Collection at the Royal Botanic Garden Edinburgh Illustrates the Floral Diversity in <i>Streptocarpus</i> (Gesneriaceae). <i>Sibbaldia the International Journal of Botanic Garden Horticulture</i> , 2019, , 155-177. | 0.1 | 2         |
| 16 | Low genetic diversity in small leading edge populations of the European paleoendemic <i>Ramonda serbica</i> (Gesneriaceae) in Bulgaria. <i>Nordic Journal of Botany</i> , 2018, 36, njb-01655.                                    | 0.2 | 3         |
| 17 | <i>Streptocarpus peltatus</i> (Gesneriaceae), a Distinctive New Species from Southeastern Madagascar. <i>Novon</i> , 2018, 26, 22-27.   | 0.3 | 1         |
| 18 | Integrating a comprehensive DNA barcode reference library with a global map of yews ( <i>Taxus</i> L.) for forensic identification. <i>Molecular Ecology Resources</i> , 2018, 18, 1115-1131.                                     | 2.2 | 38        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Two new species of <i>Oreocharis</i> (Gesneriaceae) from Fan Si Pan, the highest mountain in Vietnam. <i>PhytoKeys</i> , 2018, 94, 95-106.   | 0.4 | 17        |
| 20 | DNA barcoding of East Asian <i>Amentotaxus</i> (Taxaceae): Potential new species and implications for conservation. <i>Journal of Systematics and Evolution</i> , 2017, 55, 16-24.   | 1.6 | 25        |
| 21 | From shoot to leaf: step-wise shifts in meristem and KNOX1 activity correlate with the evolution of a unifoliate body plan in Gesneriaceae. <i>Development Genes and Evolution</i> , 2017, 227, 41-60.   | 0.4 | 10        |
| 22 | Secondary contact, hybridization and polyploidization add to the biodiversity in the Hengduan Mountains, exemplified by the widespread <i>Corallodiscus lanuginosus</i> (Gesneriaceae). <i>Plant Systematics and Evolution</i> , 2017, 303, 587-602.                                 | 0.3 | 5         |
| 23 | Both temperature fluctuations and East Asian monsoons have driven plant diversification in the karst ecosystems from southern China. <i>Molecular Ecology</i> , 2017, 26, 6414-6429.   | 2.0 | 74        |
| 24 | Multiple origins and a narrow gene pool characterise the African tea germplasm: concordant patterns revealed by nuclear and plastid DNA markers. <i>Scientific Reports</i> , 2017, 7, 4053.  | 1.6 | 22        |
| 25 | Domestication Origin and Breeding History of the Tea Plant ( <i>Camellia sinensis</i> ) in China and India Based on Nuclear Microsatellites and cpDNA Sequence Data. <i>Frontiers in Plant Science</i> , 2017, 8, 2270.  | 1.7 | 71        |
| 26 | Insights into the Genetic Relationships and Breeding Patterns of the African Tea Germplasm Based on nSSR Markers and cpDNA Sequences. <i>Frontiers in Plant Science</i> , 2016, 7, 1244.   | 1.7 | 39        |
| 27 | < i> <i>Henckelia wijesundarae</i> (< i>Gesneriaceae</ i>), a new endemic species from Sri Lanka, and lectotypification of < i> <i>Chirita walkerae</i> </ i> and < i> <i>C. walkerae</i> </ i> var. < i> <i>parviflora</i> </ i>. <i>Willdenowia</i> , 2016, 46, 213-224.           | 0.5 | 5         |
| 28 | The first phylogenetic hypothesis for the southern African endemic genus < i> <i>Tulbaghia</i> </ i> (Amaryllidaceae, Allioideae) based on plastid and nuclear DNA sequences. <i>Botanical Journal of the Linnean Society</i> , 2016, 181, 156-170.                                  | 0.8 | 12        |
| 29 | <i>Glabrella leiophylla</i> (Gesneriaceae), a new combination for a former <i>Briggsia</i> species from Guizhou, China. <i>Phytotaxa</i> , 2015, 218, 193.   | 0.1 | 1         |
| 30 | < p class="HeadingRunIn">< strong>< em>Oreocharis</em> Å— < em>heterandra</em> (Gesneriaceae): a natural hybrid from the Shengtangshan Mountains, Guangxi, China</strong></p>. <i>Phytotaxa</i> , 2015, 38, 1.   | 0.1 | 16        |
| 31 | Rearranging Specimens on Herbarium Type Sheets of <i>Streptocarpus betsiliensis</i> Humbert (Gesneriaceae). <i>Candollea</i> , 2015, 70, 145.  | 0.1 | 0         |
| 32 | < i> <i>Streptocarpus</i> </ i> redefined to include all Afroâ€“Malagasy Gesneriaceae: Molecular phylogenies prove congruent with geographical distribution and basic chromosome numbers and uncover remarkable morphological homoplasies. <i>Taxon</i> , 2015, 64, 1243-1274.       | 0.4 | 25        |
| 33 | DNA barcoding of < i> <i>Rhododendron</i> </ i> (Ericaceae), the largest Chinese plant genus in biodiversity hotspots of the Himalayaâ€“Hengduan Mountains. <i>Molecular Ecology Resources</i> , 2015, 15, 932-944.  | 2.2 | 101       |
| 34 | Chayamaritia (Gesneriaceae: Didymocarpoideae), a new genus from Southeast Asia. <i>Plant Systematics and Evolution</i> , 2015, 301, 1947-1966.   | 0.3 | 9         |
| 35 | The European Paleoendemic < i> <i>Haberlea rhodopensis</i> </ i> (Gesneriaceae) Has an Oligocene Origin and a Pleistocene Diversification and Occurs in a Long-Persisting Refugial Area in Southeastern Europe. <i>International Journal of Plant Sciences</i> , 2015, 176, 499-514. | 0.6 | 25        |
| 36 | Nuclear Ribosomal ITS Functional Paralogs Resolve the Phylogenetic Relationships of a Late-Miocene Radiation Cycad <i>Cycas</i> (Cycadaceae). <i>PLoS ONE</i> , 2015, 10, e0117971.  | 1.1 | 32        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | Low genetic diversity and high inbreeding of the endangered yews in Central Himalaya: implications for conservation of their highly fragmented populations. <i>Diversity and Distributions</i> , 2014, 20, 1270-1284.  | 1.9 | 27        |
| 38 | Three New Species of &lt;I&gt;Petrocodon&lt;/I&gt; (Gesneriaceae), Endemic to the Limestone Areas of Southwest China, and Preliminary Insights into the Diversification Patterns of the Genus. <i>Systematic Botany</i> , 2014, 39, 316-330.                   | 0.2 | 18        |
| 39 | GA2 and GA20-oxidase expressions are associated with the meristem position in <i>Streptocarpus rexii</i> (Gesneriaceae). <i>Plant Growth Regulation</i> , 2014, 72, 123-140.   | 1.8 | 6         |
| 40 | Genetic diversity, demographical history and conservation aspects of the endangered yew tree <i>Taxus contorta</i> (syn. <i>Taxus fuana</i> ) in Pakistan. <i>Tree Genetics and Genomes</i> , 2014, 10, 653-665.   | 0.6 | 24        |
| 41 | Morphological variation, genetic diversity and genome size of critically endangered <i>Haberlea</i> (Gesneriaceae) populations in Bulgaria do not support the recognition of two different species. <i>Plant Systematics and Evolution</i> , 2014, 300, 29-41. | 0.3 | 15        |
| 42 | Additional notes on <i>Oreocaris yunnanensis</i> , a species of Gesneriaceae from southern Yunnan, China, including morphological and molecular data. <i>Phytotaxa</i> , 2014, 167, 283.   | 0.1 | 7         |
| 43 | <i>Billolivia</i> , a new genus of Gesneriaceae from Vietnam with five new species. <i>Phytotaxa</i> , 2014, 161, 241.   | 0.1 | 19        |
| 44 | Taxonomic Status, Phylogenetic Affinities and Genetic Diversity of a Presumed Extinct Genus, <i>Paraisometrum</i> W.T. Wang (Gesneriaceae) from the Karst Regions of Southwest China. <i>PLoS ONE</i> , 2014, 9, e107967.                                      | 1.1 | 25        |
| 45 | <i>Oreocaris jinpingensis</i> (Gesneriaceae), a New Species from Yunnan, China. <i>Annales Botanici Fennici</i> , 2013, 50, 312-316.   | 0.0 | 10        |
| 46 | Yews ( <i>Taxus</i> ) along the Hindu Kush-Himalayan region: Exploring the ethnopharmacological relevance among communities of Mongol and Caucasian origins. <i>Journal of Ethnopharmacology</i> , 2013, 147, 190-203.   | 2.0 | 32        |
| 47 | An approach to identify putative hybrids in the “coalescent stochasticity zone”, as exemplified in the <sc>A</sc>frican plant genus <i>S</i>treptocarpus</i> (<sc>G</sc>esneriaceae). <i>New Phytologist</i> , 2013, 198, 284-300.                             | 3.5 | 21        |
| 48 | Geological and ecological factors drive cryptic speciation of yews in a biodiversity hotspot. <i>New Phytologist</i> , 2013, 199, 1093-1108.   | 3.5 | 236       |
| 49 | A multidisciplinary approach reveals hidden taxonomic diversity in the morphologically challenging <i>Taxus wallichiana</i> complex. <i>Taxon</i> , 2013, 62, 1161-1177.   | 0.4 | 18        |
| 50 | Developmental Genetics of the Perianthless Flowers and Bracts of a Paleoherb Species, <i>Saururus chinensis</i> . <i>PLoS ONE</i> , 2013, 8, e53019.   | 1.1 | 4         |
| 51 | Microsatellite markers developed for <i>Corallodiscus lanuginosus</i> (Gesneriaceae) and their cross-species transferability. <i>American Journal of Botany</i> , 2012, 99, e490-e492.   | 0.8 | 0         |
| 52 | <i>Paraboea hekouensis</i> and <i>P. manhaoensis</i>, Two New Species of Gesneriaceae from China. <i>Annales Botanici Fennici</i> , 2012, 49, 179-187.   | 0.0 | 9         |
| 53 | Using Morphological, Molecular and Climatic Data to Delimitate Yews along the Hindu Kush-Himalaya and Adjacent Regions. <i>PLoS ONE</i> , 2012, 7, e46873.   | 1.1 | 45        |
| 54 | <i>Tribounia</i>, a new genus of Gesneriaceae from Thailand. <i>Taxon</i> , 2012, 61, 1286-1295.   | 0.4 | 10        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 55 | The genetic ghost of an invasion past: colonization and extinction revealed by historical hybridization in <i>Senecio</i> . <i>Molecular Ecology</i> , 2012, 21, 369-387.  | 2.0 | 34        |
| 56 | Applying plant DNA barcodes to identify species of <i>Parnassia</i> (Parnassiaceae). <i>Molecular Ecology Resources</i> , 2012, 12, 267-275.   | 2.2 | 52        |
| 57 | DNA barcoding for the discrimination of Eurasian yews ( <i>Taxus</i> L., Taxaceae) and the discovery of cryptic species. <i>Molecular Ecology Resources</i> , 2011, 11, 89-100.  | 2.2 | 154       |
| 58 | New insights into the relationships between <i>Paraboea</i> , <i>Trisepalum</i> , and <i>Phylloboea</i> (Gesneriaceae) and their taxonomic consequences. <i>Taxon</i> , 2011, 60, 1693-1702.   | 0.4 | 18        |
| 59 | Molecular systematics and remodelling of <i>Chirita</i> and associated genera (Gesneriaceae). <i>Taxon</i> , 2011, 60, 767-790.  | 0.4 | 138       |
| 60 | The taxonomy and conservation of <i>Campanula primulifolia</i> (Campanulaceae), a critically endangered species in the Iberian Peninsula. <i>Willdenowia</i> , 2011, 41, 35-42.  | 0.5 | 1         |
| 61 | Phytogeographic aspects of <i>Lysionotus pauciflorus</i> sensu lato (Gesneriaceae) in the China, Japan and Taiwan regions: phylogenetic and morphological relationships and taxonomic consequences. <i>Plant Systematics and Evolution</i> , 2011, 292, 177-188. | 0.3 | 5         |
| 62 | A molecular phylogenetic assessment of the advanced Asiatic and Malesian didymocarpoid Gesneriaceae with focus on non-monophyletic and monotypic genera. <i>Plant Systematics and Evolution</i> , 2011, 292, 223-248.  | 0.3 | 92        |
| 63 | Cross-species amplification and development of new microsatellite loci for <i>Taxus wallichiana</i> (Taxaceae). <i>American Journal of Botany</i> , 2011, 98, e70-3.   | 0.8 | 15        |
| 64 | Extended expression of B-class MADS-box genes in the paleoherb <i>Asarum caudigerum</i> . <i>Planta</i> , 2010, 231, 265-76.   | 1.6 | 4         |
| 65 | A complex case of simple leaves: indeterminate leaves co-express ARP and KNOX1 genes. <i>Development Genes and Evolution</i> , 2010, 220, 25-40.   | 0.4 | 22        |
| 66 | A molecular phylogenetic study of <i>Hemsleya</i> (Cucurbitaceae) based on ITS, rpl16, trnH-psbA, and trnL DNA sequences. <i>Plant Systematics and Evolution</i> , 2010, 285, 23-32.   | 0.3 | 9         |
| 67 | Taxonomic treatments of <i>Camellia</i> (Theaceae) species with secretory structures based on integrated leaf characters. <i>Plant Systematics and Evolution</i> , 2010, 290, 1-20.  | 0.3 | 12        |
| 68 | High nrDNA ITS polymorphism in the ancient extant seed plant Cycas: Incomplete concerted evolution and the origin of pseudogenes. <i>Molecular Phylogenetics and Evolution</i> , 2010, 55, 168-177.  | 1.2 | 83        |
| 69 | <i>GRAMINIFOLIA</i> homolog expression in <i>Streptocarpus rexii</i> is associated with the basal meristems in phyllomorphs, a morphological novelty in Gesneriaceae. <i>Evolution &amp; Development</i> , 2010, 12, 61-73.                                      | 1.1 | 13        |
| 70 | (1948) Proposal to conserve <i>Paraboea</i> against <i>Phylloboea</i> and <i>Trisepalum</i> (Gesneriaceae). <i>Taxon</i> , 2010, 59, 1603-1603.  | 0.4 | 3         |
| 71 | A preliminary phylogeny of the didymocarpoid Gesneriaceae based on three molecular data sets: Incongruence with available tribal classifications. <i>American Journal of Botany</i> , 2009, 96, 989-1010.  | 0.8 | 81        |
| 72 | WUS and STM homologs are linked to the expression of lateral dominance in the acaulescent <i>Streptocarpus rexii</i> (Gesneriaceae). <i>Planta</i> , 2009, 230, 529-542.   | 1.6 | 15        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 73 | Molecular evidence for fragmentation among populations of <i>Taxus wallichiana</i> var. <i>mairei</i> , a highly endangered conifer in China. Canadian Journal of Forest Research, 2009, 39, 755-764.                                | 0.8 | 14        |
| 74 | Taxonomic Notes on <i>Parnassia</i> Section <i>Saxifragastrum</i> ( <i>Parnassiaceae</i> ) from China. Annales Botanici Fennici, 2009, 46, 559-565.  | 0.0 | 2         |
| 75 | < i> <i>Chirita grandibracteata</i> </i>(Gesneriaceae), a New Species from Yunnan, China. Annales Botanici Fennici, 2009, 46, 125-129.   | 0.0 | 13        |
| 76 | A new species of < i> <i>Paraboea</i> </i>(Gesneriaceae) from a karst cave in Guangxi, China, and observations on variations in flower and inflorescence architecture. Botanical Journal of the Linnean Society, 2008, 158, 681-688. | 0.8 | 18        |
| 77 | New< i> <i>Streptocarpus</i> </i>species (Gesneriaceae) from South Africa. Botanical Journal of the Linnean Society, 2008, 158, 743-748.   | 0.8 | 5         |
| 78 | Genetic diversity within and among populations of the endangered species <i>Taxus fuana</i> (Taxaceae) from Pakistan and implications for its conservation. Biochemical Systematics and Ecology, 2008, 36, 183-193.                  | 0.6 | 42        |
| 79 | Aspects of Genome Evolution in Gesneriaceae: Patterns of 45SnrDNA Site Evolution based on Evidence from Fluorescent In Situ Hybridization (FISH). International Journal of Plant Sciences, 2008, 169, 667-676.                       | 0.6 | 7         |
| 80 | The impact of pollination syndrome and habitat on gene flow: a comparative study of two < i> <i>Streptocarpus</i> </i> (Gesneriaceae) species. American Journal of Botany, 2007, 94, 1688-1695.                                      | 0.8 | 22        |
| 81 | Variation and Inheritance of Nuclear Ribosomal DNA Clusters in <i>Streptocarpus</i> (Gesneriaceae) and Their Biological and Phylogenetic Implications. International Journal of Plant Sciences, 2007, 168, 455-467.                  | 0.6 | 16        |
| 82 | Morphometric analysis of the <i>Taxus wallichiana</i> complex (Taxaceae) based on herbarium material. Botanical Journal of the Linnean Society, 2007, 155, 307-335.  | 0.8 | 42        |
| 83 | Anisocotyl and meristem initiation in an unorthodox plant, <i>Streptocarpus rexii</i> (Gesneriaceae). Planta, 2007, 225, 653-663.  | 1.6 | 23        |
| 84 | Nuclear and plastid DNA sequences confirm the placement of the enigmatic <i>Canacomyrica monticolain</i> Myricaceae. Taxon, 2006, 55, 349-357.   | 0.4 | 18        |
| 85 | Molecular evidence for repeated hybridization events involved in the origin of the genus Ă— <i>Crepidiastrixeris</i> (Asteraceae) using RAPDs and ITS data. Botanical Journal of the Linnean Society, 2006, 151, 333-343.            | 0.8 | 11        |
| 86 | Phylogeny and Biogeography of <i>Exacum</i> (Gentianaceae): A Disjunctive Distribution in the Indian Ocean Basin Resulted from Long Distance Dispersal and Extensive Radiation. Systematic Biology, 2005, 54, 21-34.                 | 2.7 | 109       |
| 87 | The Role of KNOX Genes in the Evolution of Morphological Novelty in <i>Streptocarpus</i> . Plant Cell, 2005, 17, 430-443.  | 3.1 | 45        |
| 88 | Population Genetic Structure of <i>Titanotrichum oldhamii</i> (Gesneriaceae), a Subtropical Bulbiliferous Plant with Mixed Sexual and Asexual Reproduction. Annals of Botany, 2004, 93, 201-209.                                     | 1.4 | 30        |
| 89 | Aspects of sexual failure in the reproductive processes of a rare bulbiliferous plant, <i>Titanotrichum oldhamii</i> (Gesneriaceae), in subtropical Asia. Sexual Plant Reproduction, 2004, 17, 23-31.                                | 2.2 | 11        |
| 90 | Altered expression of GFLO , the Gesneriaceae homologue of FLORICAULA/LEAFY , is associated with the transition to bulbil formation in <i>Titanotrichum oldhamii</i> . Development Genes and Evolution, 2004, 214, 122-127.          | 0.4 | 36        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 91  | Phylogenetic Position of <i>Titanotrichum oldhamii</i> (Gesneriaceae) Inferred From Four Different Gene Regions. <i>Systematic Botany</i> , 2004, 29, 407-418.  | 0.2 | 55        |
| 92  | Monophyly and relationships of the tribe Exaceae (Gentianaceae) inferred from nuclear ribosomal and chloroplast DNA sequences. <i>Molecular Phylogenetics and Evolution</i> , 2003, 28, 500-517.  | 1.2 | 51        |
| 93  | The phylogenetic position of <i>Aulotandra</i> (Zingiberaceae). <i>Nordic Journal of Botany</i> , 2003, 23, 725-734.  | 0.2 | 7         |
| 94  | Phylogenetic position and generic differentiation of Epithemateae (Gesneriaceae) inferred from plastid DNA sequence data. <i>American Journal of Botany</i> , 2003, 90, 321-329.  | 0.8 | 50        |
| 95  | Species separation of <i>&lt; i&gt;Taxus baccata&lt;/i&gt;</i> , <i>&lt; i&gt;T. canadensis&lt;/i&gt;</i> , and <i>&lt; i&gt;T. cuspidata&lt;/i&gt;</i> (Taxaceae) and origins of their reputed hybrids inferred from RAPD and cpDNA data. <i>American Journal of Botany</i> , 2003, 90, 175-182. | 0.8 | 51        |
| 96  | Phylogenetic Studies in <i>Streptocarpus</i> (Gesneriaceae): Reconstruction of Biogeographic History and Distribution Patterns. <i>Systematics and Geography of Plants</i> , 2001, 71, 545.   | 0.1 | 32        |
| 97  | Diversity of cycloidea -like Genes in Gesneriaceae in Relation to Floral Symmetry. <i>Annals of Botany</i> , 2000, 86, 167-176.   | 1.4 | 108       |
| 98  | Phylogenetic Relationships of <i>Asterella</i> (Ayttoniaceae, Marchantiopsida) Inferred from Chloroplast DNA Sequences. <i>Bryologist</i> , 2000, 103, 625-644.   | 0.1 | 26        |
| 99  | Evolution and Development of Floral Diversity in <i>Streptocarpus</i> and <i>Saintpaulia</i> . <i>Annals of Botany</i> , 1999, 84, 49-60.   | 1.4 | 71        |
| 100 | Origin and relationships of <i>Saintpaulia</i> (Gesneriaceae) based on ribosomal DNA internal transcribed spacer (ITS) sequences. <i>American Journal of Botany</i> , 1997, 84, 956-965.  | 0.8 | 114       |
| 101 | Phylogeny and disjunct distribution: evolution of <i>Saintpaulia</i> (Gesneriaceae). <i>Proceedings of the Royal Society B: Biological Sciences</i> , 1997, 264, 1827-1836.   | 1.2 | 30        |
| 102 | Strange morphogenesis – organ determination in Monophyllaea. <i>Trends in Plant Science</i> , 1997, 2, 327-328.   | 4.3 | 18        |