Yingxian Sun

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/374961/publications.pdf

Version: 2024-02-01

104 papers 2,075 citations

257450 24 h-index 330143 37 g-index

104 all docs

104 docs citations

times ranked

104

3232 citing authors

#	Article	IF	CITATIONS
1	Hyperlipidemia and hypertension have synergistic interaction on ischemic stroke: insights from a general population survey in China. BMC Cardiovascular Disorders, 2022, 22, 47.	1.7	9
2	Serum Spermidine in Relation to Risk of Stroke: A Multilevel Study. Frontiers in Nutrition, 2022, 9, 843616.	3.7	5
3	The effects of transitions in metabolic health and obesity status on incident cardiovascular disease: Insights from a general Chinese population. European Journal of Preventive Cardiology, 2021, 28, 1250-1258.	1.8	25
4	Self-reported snoring, snoring intensity, and incident cardiovascular disease events: insights from a large Asian population-based cohort study. Sleep and Breathing, 2021, 25, 365-366.	1.7	5
5	Predictive value of echocardiographic left atrial size for incident stoke and stroke cause mortality: a population-based study. BMJ Open, 2021, 11, e043595.	1.9	5
6	Combined effect of hypertension and hyperuricemia on ischemic stroke in a rural Chinese population. BMC Public Health, 2021, 21, 776.	2.9	8
7	Echocardiographic left ventricular geometry profiles for prediction of stroke, coronary heart disease and all-cause mortality in the Chinese community: a rural cohort population study. BMC Cardiovascular Disorders, 2021, 21, 238.	1.7	7
8	Corrected QT Interval Is Associated With Stroke but Not Coronary Heart Disease: Insights From a General Chinese Population. Frontiers in Cardiovascular Medicine, 2021, 8, 605774.	2.4	3
9	Associations between aortic regurgitation severity and risk of incident myocardial infarction and stroke among patients with degenerative aortic valve disease: insights from a large Chinese population-based cohort study. BMJ Open, 2021, 11, e046824.	1.9	1
10	Gender discrepancy in the predictive effect of metabolic syndrome and its components on newly onset cardiovascular disease in elderly from rural China. BMC Geriatrics, 2021, 21, 505.	2.7	11
11	It's Time to Add Electrocardiography and Echocardiography to CVD Risk Prediction Models: Results from a Prospective Cohort Study. Risk Management and Healthcare Policy, 2021, Volume 14, 4657-4671.	2.5	3
12	Usefulness of metabolic score for insulin resistance index in estimating the risk of mildly reduced estimate glomerular filtration rate: a cross-sectional study of rural population in China. BMJ Open, 2021, 11, e050907.	1.9	5
13	Cardiometabolic comorbidities and epidemiological features among rural Chinese elderly people. Aging Clinical and Experimental Research, 2020, 32, 1777-1788.	2.9	5
14	Selfâ€reported snoring, snoring intensity and incident coronary heart disease events: Insights from a large Chinese populationâ€based cohort study. Geriatrics and Gerontology International, 2020, 20, 89-90.	1.5	2
15	Lower or higher HDL-C levels are associated with cardiovascular events in the general population in rural China. Lipids in Health and Disease, 2020, 19, 152.	3.0	13
16	PHLDA1 is a new therapeutic target of oxidative stress and ischemia reperfusion-induced myocardial injury. Life Sciences, 2020, 245, 117347.	4.3	20
17	Effects of long- and short-term body mass index changes on incident hypertension are different. Nutrition, 2020, 74, 110755.	2.4	1
18	Using the Atherogenic Index of Plasma to Estimate the Prevalence of Ischemic Stroke within a General Population in a Rural Area of China. BioMed Research International, 2020, 2020, 1-7.	1.9	16

#	Article	IF	Citations
19	Hyperhomocysteinemia accompany with metabolic syndrome increase the risk of left ventricular hypertrophy in rural Chinese. BMC Cardiovascular Disorders, 2020, 20, 44.	1.7	9
20	Prevalence and risk factors of prolonged corrected QT interval in general Chinese population. BMC Cardiovascular Disorders, 2019, 19, 276.	1.7	19
21	Metabolic Profile for Prediction of Ischemic Stroke in Chinese Hypertensive Population. Journal of Stroke and Cerebrovascular Diseases, 2019, 28, 1062-1069.	1.6	21
22	Non-traditional lipid profiles associated with ischemic stroke not hemorrhagic stroke in hypertensive patients: results from an 8.4 years follow-up study. Lipids in Health and Disease, 2019, 18, 9.	3.0	19
23	Gender-specific contribution of cardiometabolic index and lipid accumulation product to left ventricular geometry change in general population of rural China. BMC Cardiovascular Disorders, 2018, 18, 62.	1.7	21
24	Validity of cardiometabolic index, lipid accumulation product, and body adiposity index in predicting the risk of hypertension in Chinese population. Postgraduate Medicine, 2018, 130, 325-333.	2.0	52
25	BAG3 promotes the phenotypic transformation of primary rat vascular smooth muscle cells via TRAIL. International Journal of Molecular Medicine, 2018, 41, 2917-2926.	4.0	5
26	Improving diagnosis of left ventricular hypertrophy in Chinese obese individuals: Which indexation criteria of left ventricular mass?. International Journal of Cardiology, 2018, 271, 344.	1.7	0
27	Effect of serum electrolytes within normal ranges on QTc prolongation: a cross-sectional study in a Chinese rural general population. BMC Cardiovascular Disorders, 2018, 18, 175.	1.7	14
28	Body adiposity index, lipid accumulation product, and cardiometabolic index reveal the contribution of adiposity phenotypes in the risk of hyperuricemia among Chinese rural population. Clinical Rheumatology, 2018, 37, 2221-2231.	2.2	21
29	Does sociodemographic characteristics affect ideal cardiovascular health score in Chinese population?. International Journal of Cardiology, 2018, 264, 179.	1.7	0
30	Ideal Cardiovascular Health Score and Fuster-BEWAT Score as Predictors of Left Ventricular Hypertrophy Classification. Journal of the American College of Cardiology, 2018, 72, 581-582.	2.8	4
31	Estimate of ischemic stroke prevalence according to a novel 4-tiered classification of left ventricular hypertrophy: insights from the general Chinese population. Annals of Medicine, 2018, 50, 519-528.	3.8	4
32	The impact of nontraditional lipid profiles on left ventricular geometric abnormalities in general Chinese population. BMC Cardiovascular Disorders, 2018, 18, 88.	1.7	10
33	Effects of Metabolically Healthy and Unhealthy Obesity on Prolongation of Corrected QT Interval. American Journal of Cardiology, 2017, 119, 1199-1204.	1.6	14
34	Advanced glycation end products promote the proliferation and migration of primary rat vascular smooth muscle cells via the upregulation of BAG3. International Journal of Molecular Medicine, 2017, 39, 1242-1254.	4.0	12
35	The association of ideal cardiovascular health and left ventricle hypertrophy in rural population of northeast China. Medicine (United States), 2017, 96, e6050.	1.0	5
36	Sex-specific association between serum uric acid and self-reported snoring in rural China: a cross-sectional study. Sleep and Breathing, 2017, 21, 939-947.	1.7	0

#	Article	IF	CITATIONS
37	Sex differences in association between decreased glomerular filtration rate and prolongation of corrected QT interval in general Chinese population. European Journal of Internal Medicine, 2017, 43, e33-e35.	2.2	3
38	Plasma homocysteine levels associated with a corrected QT interval. BMC Cardiovascular Disorders, 2017, 17, 182.	1.7	7
39	Gender-specific association of decreased estimated glomerular filtration rate and left vertical geometry in the general population from rural Northeast China. BMC Cardiovascular Disorders, 2017, 17, 24.	1.7	3
40	Elevated Homocysteine Concentrations Decrease the Antihypertensive Effect of Angiotensin-Converting Enzyme Inhibitors in Hypertensive Patients. Arteriosclerosis, Thrombosis, and Vascular Biology, 2017, 37, 166-172.	2.4	38
41	The prevalence and predictors of metabolically healthy obesity in obese rural population of China: a cross-sectional study. Psychology, Health and Medicine, 2017, 22, 303-309.	2.4	10
42	Metabolic Abnormalities, But Not Metabolically Healthy Obesity, Are Associated with Left Ventricular Hypertrophy. Heart Lung and Circulation, 2017, 26, 251-257.	0.4	15
43	Relation of four nontraditional lipid profiles to diabetes in rural Chinese H-type hypertension population. Lipids in Health and Disease, 2017, 16, 199.	3.0	5
44	The Prevalence of Hypertension Accompanied by High Homocysteine and its Risk Factors in a Rural Population: A Cross-Sectional Study from Northeast China. International Journal of Environmental Research and Public Health, 2017, 14, 376.	2.6	16
45	Advanced Glycation End Products Inhibit the Proliferation of Human Umbilical Vein Endothelial Cells by Inhibiting Cathepsin D. International Journal of Molecular Sciences, 2017, 18, 436.	4.1	19
46	Impact of hypertension with or without diabetes on left ventricular remodeling in rural Chinese population: a cross-sectional study. BMC Cardiovascular Disorders, 2017, 17, 206.	1.7	19
47	Self-Reported Snoring Is Associated with Dyslipidemia, High Total Cholesterol, and High Low-Density Lipoprotein Cholesterol in Obesity: A Cross-Sectional Study from a Rural Area of China. International Journal of Environmental Research and Public Health, 2017, 14, 86.	2.6	25
48	Association between Serum Uric Acid and Elevated Alanine Aminotransferase in the General Population. International Journal of Environmental Research and Public Health, 2016, 13, 841.	2.6	7
49	Diagnosed but Not Undiagnosed Diabetes Is Associated with Depression in Rural Areas. International Journal of Environmental Research and Public Health, 2016, 13, 1136.	2.6	15
50	The Protective Role of the TOPK/PBK Pathway in Myocardial Ischemia/Reperfusion and H2O2-Induced Injury in H9C2 Cardiomyocytes. International Journal of Molecular Sciences, 2016, 17, 267.	4.1	26
51	Visceral Adiposity Index and Lipid Accumulation Product Index: Two Alternate Body Indices to Identify Chronic Kidney Disease among the Rural Population in Northeast China. International Journal of Environmental Research and Public Health, 2016, 13, 1231.	2.6	49
52	Modified Ideal Cardiovascular Health Status is Associated with Lower Prevalence of Stroke in Rural Northeast China. International Journal of Environmental Research and Public Health, 2016, 13, 207.	2.6	9
53	Anthropometric Indices in Adults: Which Is the Best Indicator to Identify Alanine Aminotransferase Levels?. International Journal of Environmental Research and Public Health, 2016, 13, 226.	2.6	12
54	Association between the Hypertriglyceridemic Waist Phenotype, Prediabetes, and Diabetes Mellitus in Rural Chinese Population: A Cross-Sectional Study. International Journal of Environmental Research and Public Health, 2016, 13, 368.	2.6	19

#	Article	IF	Citations
55	Association between Obese Phenotype and Mildly Reduced eGFR among the General Population from Rural Northeast China. International Journal of Environmental Research and Public Health, 2016, 13, 540.	2.6	6
56	The Relation of Moderate Alcohol Consumption to Hyperuricemia in a Rural General Population. International Journal of Environmental Research and Public Health, 2016, 13, 732.	2.6	21
57	Comprehensive Comparison between Empty Nest and Non-Empty Nest Elderly: A Cross-Sectional Study among Rural Populations in Northeast China. International Journal of Environmental Research and Public Health, 2016, 13, 857.	2.6	37
58	Atherogenic Index of Plasma Predicts Hyperuricemia in Rural Population: A Cross-Sectional Study from Northeast China. International Journal of Environmental Research and Public Health, 2016, 13, 879.	2.6	10
59	The Association of Ideal Cardiovascular Health and Atherogenic Index of Plasma in Rural Population: A Cross-Sectional Study from Northeast China. International Journal of Environmental Research and Public Health, 2016, 13, 1027.	2.6	16
60	The Current Situation of Hypertension among Rural Minimal Assurance Family Participants in Liaoning (China): A Cross-Sectional Study. International Journal of Environmental Research and Public Health, 2016, 13, 1199.	2.6	2
61	Metabolism rather than obesity is associated with ischemic stroke: a cross-sectional study in rural Northeastern China. SpringerPlus, 2016, 5, 1419.	1.2	10
62	Metabolic Syndrome and Serum Liver Enzymes in the General Chinese Population. International Journal of Environmental Research and Public Health, 2016, 13, 223.	2.6	43
63	Atorvastatin inhibits the apoptosis of human umbilical vein endothelial cells induced by angiotensin II via the lysosomal-mitochondrial axis. Apoptosis: an International Journal on Programmed Cell Death, 2016, 21, 977-996.	4.9	28
64	The Association Between Alcohol Consumption and Left Ventricular Ejection Fraction. Medicine (United States), 2016, 95, e3763.	1.0	15
65	A Body Shape Index and Body Roundness Index: Two New Body Indices to Identify left Ventricular Hypertrophy among Rural Populations in Northeast China. Heart Lung and Circulation, 2016, 25, 358-364.	0.4	33
66	Prevalence and Metrics Distribution of Ideal Cardiovascular Health: A Population-based, Cross-sectional Study in Rural China. Heart Lung and Circulation, 2016, 25, 982-992.	0.4	29
67	Metabolically healthy obesity also has risk for hyperuricemia among Chinese general population: A cross-sectional study. Obesity Research and Clinical Practice, 2016, 10, S84-S95.	1.8	29
68	Prevalence of abnormal serum liver enzymes in patients with type 2 diabetes mellitus: a cross-sectional study from China. Postgraduate Medicine, 2016, 128, 770-776.	2.0	15
69	Relation of Heavy Alcohol Consumption to QTc Interval Prolongation. American Journal of Cardiology, 2016, 118, 1201-1206.	1.6	16
70	Geometrical and functional changes of left heart in adults with prehypertension and hypertension: a cross-sectional study from China. BMC Cardiovascular Disorders, 2016, 16, 114.	1.7	19
71	Sex-specific association between serum uric acid and prolonged corrected QT interval. Medicine (United States), 2016, 95, e5568.	1.0	7
72	Hypertriglyceridemic waist phenotype and metabolic abnormalities in hypertensive adults. Medicine (United States), 2016, 95, e5613.	1.0	18

#	Article	IF	CITATIONS
73	The relationship between snoring and left ventricular hypertrophy of China: a cross-sectional study. BMC Cardiovascular Disorders, 2016, 16, 15.	1.7	10
74	Prevalence of left atrial enlargement and its risk factors in general Chinese population. BMC Cardiovascular Disorders, 2016, 16, 53.	1.7	12
75	Daytime sleepiness is associated with hyperhomocysteinemia in rural area of China: A cross-sectional study. European Journal of Internal Medicine, 2016, 35, 73-77.	2.2	3
76	A Body Shape Index and Body Roundness Index: Two new body indices for detecting association between obesity and hyperuricemia in rural area of China. European Journal of Internal Medicine, 2016, 29, 32-36.	2.2	48
77	Alcohol consumption and cardiovascular diseases in rural China. International Journal of Cardiology, 2016, 215, 257-262.	1.7	68
78	Optimal Blood Pressure in Patients after Stroke in Rural Areas of China. Journal of Stroke and Cerebrovascular Diseases, 2016, 25, 270-280.	1.6	3
79	Diabetes mellitus is an independent risk factor for atrial fibrillation in a general Chinese population. Journal of Diabetes Investigation, 2016, 7, 791-796.	2.4	16
80	Prevalence of dyslipidemia and associated factors among the hypertensive population from rural Northeast China. BMC Public Health, 2015, 15, 1152.	2.9	16
81	Prevalence of Diabetes and Impaired Fasting Glucose in Hypertensive Adults in Rural China: Far from Leveling-Off. International Journal of Environmental Research and Public Health, 2015, 12, 14764-14779.	2.6	22
82	Atorvastatin Calcium Inhibits Phenotypic Modulation of PDGF-BB-Induced VSMCs via Down-Regulation the Akt Signaling Pathway. PLoS ONE, 2015, 10, e0122577.	2.5	34
83	Advanced glycation end products promote proliferation and suppress autophagy via reduction of Cathepsin D in rat vascular smooth muscle cells. Molecular and Cellular Biochemistry, 2015, 403, 73-83.	3.1	22
84	Grim status of hypertension in rural China: results from Northeast China Rural Cardiovascular Health Study 2013. Journal of the American Society of Hypertension, 2015, 9, 358-364.	2.3	52
85	Combination of hyperuricemia and metabolic syndrome is an independent and powerful predictor for left ventricular hypertrophy in rural Chinese. Annales D'Endocrinologie, 2015, 76, 264-271.	1.4	12
86	Soybeans or soybean products consumption and depressive symptoms in older residents in rural Northeast China: A cross-sectional study. Journal of Nutrition, Health and Aging, 2015, 19, 884-893.	3.3	15
87	Metabolic syndrome in hypertensive adults from rural Northeast China: an update. BMC Public Health, 2015, 15, 247.	2.9	17
88	Hyperuricemia is independently associated with left ventricular hypertrophy in post-menopausal women but not in pre-menopausal women in rural Northeast China. Gynecological Endocrinology, 2015, 31, 736-741.	1.7	24
89	Selfâ€reported sleep duration is associated with reduced glomerular filtration rate among adults with hypertension: a populationâ€based study from rural northeast <scp>C</scp> hina. Journal of Sleep Research, 2015, 24, 351-358.	3.2	27
90	Comparison of four nontraditional lipid profiles in relation to ischemic stroke among hypertensive Chinese population. International Journal of Cardiology, 2015, 201, 123-125.	1.7	13

#	Article	IF	CITATION
91	Association between elevated serum alanine aminotransferase and cardiometabolic risk factors in rural Chinese population: a cross-sectional study. BMC Cardiovascular Disorders, 2015, 15, 65.	1.7	31
92	Prevalence and risk factors of hypertension among pre- and post-menopausal women: A cross-sectional study in a rural area of northeast China. Maturitas, 2015, 80, 282-287.	2.4	28
93	An update on overweight and obesity in rural Northeast China: from lifestyle risk factors to cardiometabolic comorbidities. BMC Public Health, 2014, 14, 1046.	2.9	42
94	Prehypertension in Rural Northeastern China: Results From the Northeast China Rural Cardiovascular Health Study. Journal of Clinical Hypertension, 2014, 16, 664-670.	2.0	24
95	An update on the prevalence of metabolic syndrome and its associated factors in rural northeast China. BMC Public Health, 2014, 14, 877.	2.9	69
96	The Prevalence and Risk Factors for Depression Symptoms in a Rural Chinese Sample Population. PLoS ONE, 2014, 9, e99692.	2.5	85
97	The association between glomerular filtration rate and stroke in hypertensive patients in rural areas of China. Journal of Hypertension, 2012, 30, 901-907.	0.5	15
98	Decreased Glomerular Filtration Rate Is Associated with Mortality and Cardiovascular Events in Patients with Hypertension: A Prospective Study. PLoS ONE, 2011, 6, e27359.	2.5	16
99	Prevalence of diabetes and impaired fasting glucose in hypertensive adults in rural China. Acta Cardiologica, 2009, 64, 351-356.	0.9	9
100	Prevalence and Associated Factors of Overweight and Obesity in a Chinese Rural Population. Obesity, 2008, 16, 168-171.	3.0	44
101	Pulse Pressure and Mean Arterial Pressure in Relation to Ischemic Stroke Among Patients With Uncontrolled Hypertension in Rural Areas of China. Stroke, 2008, 39, 1932-1937.	2.0	123
102	Prevalence, Awareness, Treatment, and Control of Hypertension in Rural Adults from Liaoning Province, Northeast China. Hypertension Research, 2007, 30, 951-958.	2.7	44
103	Prevalence and Risk Factors of the Rural Adult People Prehypertension Status in Liaoning Province of China. Circulation Journal, 2007, 71, 550-553.	1.6	45
104	The prevalence of prehypertension and hypertension among rural adults in liaoning province of China. Clinical Cardiology, 2007, 30, 183-187.	1.8	61