

Beata Greb-Markiewicz

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3751168/publications.pdf>

Version: 2024-02-01

16
papers

231
citations

1307594

7
h-index

996975

15
g-index

18
all docs

18
docs citations

18
times ranked

255
citing authors

#	ARTICLE	IF	CITATIONS
1	Immobilization of wood-rotting fungi laccases on modified cellulose and acrylic carriers. <i>Process Biochemistry</i> , 2002, 37, 1387-1394.	3.7	77
2	The Significance of the Intrinsically Disordered Regions for the Functions of the bHLH Transcription Factors. <i>International Journal of Molecular Sciences</i> , 2019, 20, 5306.	4.1	29
3	bHLH-PAS Proteins: Their Structure and Intrinsic Disorder. <i>International Journal of Molecular Sciences</i> , 2019, 20, 3653.	4.1	27
4	Sequences that direct subcellular traffic of the <i>Drosophila</i> methoprene-tolerant protein (MET) are located predominantly in the PAS domains. <i>Molecular and Cellular Endocrinology</i> , 2011, 345, 16-26.	3.2	22
5	The variety of complexes formed by EcR and Usp nuclear receptors in the nuclei of living cells. <i>Molecular and Cellular Endocrinology</i> , 2008, 294, 45-51.	3.2	12
6	Mapping of the Sequences Directing Localization of the <i>Drosophila</i> Germ Cell-Expressed Protein (GCE). <i>PLoS ONE</i> , 2015, 10, e0133307.	2.5	12
7	Multiple sequences orchestrate subcellular trafficking of neuronal PAS domain-containing protein 4 (NPAS4). <i>Journal of Biological Chemistry</i> , 2018, 293, 11255-11270.	3.4	10
8	Subcellular Localization Signals of bHLH-PAS Proteins: Their Significance, Current State of Knowledge and Future Perspectives. <i>International Journal of Molecular Sciences</i> , 2019, 20, 4746.	4.1	9
9	The intrinsically disordered region of GCE protein adopts a more fixed structure by interacting with the LBD of the nuclear receptor FTZ-F1. <i>Cell Communication and Signaling</i> , 2020, 18, 180.	6.5	8
10	Intrinsic Disorder of the C-Terminal Domain of <i>Drosophila</i> Methoprene-Tolerant Protein. <i>PLoS ONE</i> , 2016, 11, e0162950.	2.5	8
11	The method utilized to purify the SARS-CoV-2 N protein can affect its molecular properties. <i>International Journal of Biological Macromolecules</i> , 2021, 188, 391-403.	7.5	6
12	The subcellular localization of bHLH transcription factor TCF4 is mediated by multiple nuclear localization and nuclear export signals. <i>Scientific Reports</i> , 2019, 9, 15629.	3.3	5
13	Ligand binding is without effect on complex formation of the ligand binding domain of the ecdysone receptor (EcR). <i>Archives of Insect Biochemistry and Physiology</i> , 2005, 59, 1-11.	1.5	3
14	The Participation of the Intrinsically Disordered Regions of the bHLH-PAS Transcription Factors in Disease Development. <i>International Journal of Molecular Sciences</i> , 2021, 22, 2868.	4.1	2
15	Microbial transformation of isoprenoid systems by means of fungi of <i>Zoopphthora</i> genus – Microbial transformation of compounds with 1 \pm -campholenic system by means of fungus <i>Zoopphthora</i> (<i>Neopandora</i>) sp.. <i>Journal of Basic Microbiology</i> , 1995, 35, 433-439.	3.3	1
16	Structural Analyses of Ordered and Disordered Regions in Ecdysteroid Receptor. , 2015, , 93-117.		0