

# Monika Ryba-StanisÅ,awowska

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/375207/publications.pdf>

Version: 2024-02-01

28  
papers

550  
citations

840585

11  
h-index

642610

23  
g-index

30  
all docs

30  
docs citations

30  
times ranked

1039  
citing authors

#	ARTICLE	IF	CITATIONS
1	Associations of TP53 codon 72 polymorphism with complications and comorbidities in patients with type 1 diabetes. <i>Journal of Molecular Medicine</i> , 2021, 99, 675-683.	1.7	13
2	A new potential mode of cardiorenal protection of KLOTHO gene variability in type 1 diabetic adolescents. <i>Journal of Molecular Medicine</i> , 2020, 98, 955-962.	1.7	4
3	Regulatory T cells: the future of autoimmune disease treatment. <i>Expert Review of Clinical Immunology</i> , 2019, 15, 777-789.	1.3	22
4	Putative loss of CD83 immunosuppressive activity in long-standing complication-free juvenile diabetic patients during disease progression. <i>Immunologic Research</i> , 2019, 67, 70-76.	1.3	4
5	Monocytes of newly diagnosed juvenile DM1 patients are prone to differentiate into regulatory IL-10+ M2 macrophages. <i>Immunologic Research</i> , 2019, 67, 58-69.	1.3	9
6	CCR5- $\Delta 32$ polymorphism is a genetic risk factor associated with dyslipidemia in patients with type 1 diabetes. <i>Cytokine</i> , 2019, 114, 81-85.	1.4	11
7	New mechanisms of CCR5- $\Delta 32$ carriers' advantage "Impact on progenitor cells and renal function. <i>International Journal of Biochemistry and Cell Biology</i> , 2019, 108, 92-97.	1.2	1
8	The KL-VS polymorphism of KLOTHO gene is protective against retinopathy incidence in patients with type 1 diabetes. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2018, 1864, 758-763.	1.8	16
9	Circulating CD34+ and CD34+VEGFR2+ progenitor cells are associated with KLOTHO KL-VS polymorphism. <i>Microvascular Research</i> , 2018, 119, 1-6.	1.1	2
10	Sex-related association of serum uric acid with inflammation, kidney function and blood pressure in type 1 diabetic patients. <i>Pediatric Diabetes</i> , 2018, 19, 1014-1019.	1.2	6
11	Estrogen receptor $\beta$ gene polymorphism and vascular complications in girls with type 1 diabetes mellitus. <i>Molecular and Cellular Biochemistry</i> , 2018, 437, 153-161.	1.4	9
12	IL-33 improves the suppressive potential of regulatory T cells in patients with type 1 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2017, 128, 67-73.	1.1	13
13	CCR5- $\Delta 32$ gene polymorphism is related to celiac disease and autoimmune thyroiditis coincidence in patients with type 1 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 615-618.	1.2	8
14	CCR5- $\Delta 32$ gene polymorphism is associated with retinopathy in patients with type 1 diabetes. <i>Molecular and Cellular Endocrinology</i> , 2017, 439, 256-260.	1.6	10
15	IL-33 Effect on Quantitative Changes of CD4+CD25highFOXP3+Regulatory T Cells in Children with Type 1 Diabetes. <i>Mediators of Inflammation</i> , 2016, 2016, 1-7.	1.4	17
16	Th9 and Th22 immune response in young patients with type 1 diabetes. <i>Immunologic Research</i> , 2016, 64, 730-735.	1.3	26
17	Elevated levels of peripheral blood CD <sup>bright</sup> CD <sup>16</sup> + and CD <sup>dim</sup> CD <sup>16</sup> + monocytes may contribute to the development of retinopathy in patients with juvenile onset type 1 diabetes. <i>Apmis</i> , 2015, 123, 793-799.	0.9	12
18	Quantitative and functional characteristics of endothelial progenitor cells in newly diagnosed hypertensive patients. <i>Journal of Human Hypertension</i> , 2015, 29, 324-330.	1.0	10

#	ARTICLE	IF	CITATIONS
19	Different pathways of macrophage activation and polarization. Postepy Higieny I Medycyny Doswiadczalnej, 2015, 69, 496-502.	0.1	120
20	IVS1 $\hat{r}$ 397T<b>&gt;</b>C Estrogen Receptor<b><i> $\hat{\pm}$ </i></b>Polymorphism Is Associated with Low-Grade Systemic Inflammatory Response in Type 1 Diabetic Girls. Mediators of Inflammation, 2014, 2014, 1-8.	1.4	11
21	Enhanced Apoptosis of Monocytes from Complication-Free Juvenile-Onset Diabetes Mellitus Type 1 May Be Ameliorated by TNF $\hat{\pm}$ Inhibitors. Mediators of Inflammation, 2014, 2014, 1-11.	1.4	11
22	Effector and regulatory T cell subsets in diabetes-associated inflammation. Is there a connection with ST2/IL-33 axis? Perspective. Autoimmunity, 2014, 47, 361-371.	1.2	1
23	Elevated Levels of Serum IL-12 and IL-18 are Associated with Lower Frequencies of CD4+CD25 <sup>high</sup> FOXP3+ Regulatory T cells in Young Patients with Type 1 Diabetes. Inflammation, 2014, 37, 1513-1520.	1.7	47
24	Loss of the balance between CD4+Foxp3+ regulatory T cells and CD4+IL17A+ Th17 cells in patients with type 1 diabetes. Human Immunology, 2013, 74, 701-707.	1.2	59
25	The Serum IL-6 Profile and Treg/Th17 Peripheral Cell Populations in Patients with Type 1 Diabetes. Mediators of Inflammation, 2013, 2013, 1-7.	1.4	41
26	Anti-TNF rescue CD4+Foxp3+ regulatory T cells in patients with type 1 diabetes from effects mediated by TNF. Cytokine, 2011, 55, 353-361.	1.4	34
27	The association of the IVS1-397T>C estrogen receptor $\hat{\pm}$ polymorphism with the regulatory conditions in longstanding type 1 diabetic girls. Molecular Immunology, 2011, 49, 324-328.	1.0	7
28	Lower Frequency of CD62L <sup>high</sup> and Higher Frequency of TNFR2 <sup>++</sup> Tregs Are Associated with Inflammatory Conditions in Type 1 Diabetic Patients. Mediators of Inflammation, 2011, 2011, 1-7.	1.4	20