

# Eric A Schmelz

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/3754253/eric-a-schmelz-publications-by-citations.pdf>

**Version:** 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

104  
papers

8,441  
citations

51  
h-index

91  
g-index

119  
ext. papers

9,894  
ext. citations

6.2  
avg, IF

5.77  
L-index

#	Paper	IF	Citations
104	Airborne signals prime plants against insect herbivore attack. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2004</b> , 101, 1781-5	11.5	627
103	ABA is an essential signal for plant resistance to pathogens affecting JA biosynthesis and the activation of defenses in Arabidopsis. <i>Plant Cell</i> , <b>2007</b> , 19, 1665-81	11.6	621
102	Fragments of ATP synthase mediate plant perception of insect attack. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2006</b> , 103, 8894-9	11.5	300
101	Simultaneous analysis of phytohormones, phytotoxins, and volatile organic compounds in plants. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2003</b> , 100, 10552-7	11.5	280
100	Quantification, correlations and manipulations of wound-induced changes in jasmonic acid and nicotine in <i>Nicotiana sylvestris</i> . <i>Planta</i> , <b>1997</b> , 201, 397-404	4.7	261
99	Circadian regulation of the PhCCD1 carotenoid cleavage dioxygenase controls emission of beta-ionone, a fragrance volatile of petunia flowers. <i>Plant Physiology</i> , <b>2004</b> , 136, 3504-14	6.6	221
98	The use of vapor phase extraction in metabolic profiling of phytohormones and other metabolites. <i>Plant Journal</i> , <b>2004</b> , 39, 790-808	6.9	208
97	Phytohormone-based activity mapping of insect herbivore-produced elicitors. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2009</b> , 106, 653-7	11.5	205
96	Wound-induced changes in root and shoot jasmonic acid pools correlate with induced nicotine synthesis in <i>Nicotiana sylvestris</i> spegazzini and comes. <i>Journal of Chemical Ecology</i> , <b>1994</b> , 20, 2139-57	2.7	203
95	tasselseed1 is a lipoxygenase affecting jasmonic acid signaling in sex determination of maize. <i>Science</i> , <b>2009</b> , 323, 262-5	33.3	198
94	Identification of loci affecting flavour volatile emissions in tomato fruits. <i>Journal of Experimental Botany</i> , <b>2006</b> , 57, 887-96	7	194
93	Disulfoxy fatty acids from the American bird grasshopper <i>Schistocerca americana</i> , elicitors of plant volatiles. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2007</b> , 104, 12976-81	11.5	182
92	Identity, regulation, and activity of inducible diterpenoid phytoalexins in maize. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2011</b> , 108, 5455-60	11.5	179
91	Novel acidic sesquiterpenoids constitute a dominant class of pathogen-induced phytoalexins in maize. <i>Plant Physiology</i> , <b>2011</b> , 156, 2082-97	6.6	156
90	Quantitative relationships between induced jasmonic acid levels and volatile emission in <i>Zea mays</i> during <i>Spodoptera exigua</i> herbivory. <i>Planta</i> , <b>2003</b> , 216, 665-73	4.7	155
89	Biosynthesis, elicitation and roles of monocot terpenoid phytoalexins. <i>Plant Journal</i> , <b>2014</b> , 79, 659-78	6.9	151
88	The influence of intact-plant and excised-leaf bioassay designs on volicitin- and jasmonic acid-induced sesquiterpene volatile release in <i>Zea mays</i> . <i>Planta</i> , <b>2001</b> , 214, 171-9	4.7	148

87	Plant elicitor peptides are conserved signals regulating direct and indirect antiherbivore defense. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2013</b> , 110, 5707-12	11.5	137
86	Susceptible to intolerance--a range of hormonal actions in a susceptible Arabidopsis pathogen response. <i>Plant Journal</i> , <b>2003</b> , 33, 245-57	6.9	133
85	XopD SUMO protease affects host transcription, promotes pathogen growth, and delays symptom development in xanthomonas-infected tomato leaves. <i>Plant Cell</i> , <b>2008</b> , 20, 1915-29	11.6	132
84	ZmPep1, an ortholog of Arabidopsis elicitor peptide 1, regulates maize innate immunity and enhances disease resistance. <i>Plant Physiology</i> , <b>2011</b> , 155, 1325-38	6.6	129
83	Attraction of Spodoptera frugiperda larvae to volatiles from herbivore-damaged maize seedlings. <i>Journal of Chemical Ecology</i> , <b>2006</b> , 32, 1911-24	2.7	124
82	Simultaneous quantification of jasmonic acid and salicylic acid in plants by vapor-phase extraction and gas chromatography-chemical ionization-mass spectrometry. <i>Analytical Biochemistry</i> , <b>2003</b> , 312, 242-50	3.1	121
81	Ethylene-regulated floral volatile synthesis in petunia corollas. <i>Plant Physiology</i> , <b>2005</b> , 138, 255-66	6.6	120
80	Synergistic interactions between volicitin, jasmonic acid and ethylene mediate insect-induced volatile emission in Zea mays. <i>Physiologia Plantarum</i> , <b>2003</b> , 117, 403-412	4.6	117
79	Multiple hormones act sequentially to mediate a susceptible tomato pathogen defense response. <i>Plant Physiology</i> , <b>2003</b> , 133, 1181-9	6.6	117
78	Nitrogen deficiency increases volicitin-induced volatile emission, jasmonic acid accumulation, and ethylene sensitivity in maize. <i>Plant Physiology</i> , <b>2003</b> , 133, 295-306	6.6	116
77	A 13-lipoxygenase, TomloxC, is essential for synthesis of C5 flavour volatiles in tomato. <i>Journal of Experimental Botany</i> , <b>2014</b> , 65, 419-28	7	111
76	Differential volatile emissions and salicylic acid levels from tobacco plants in response to different strains of Pseudomonas syringae. <i>Planta</i> , <b>2003</b> , 217, 767-75	4.7	106
75	Cowpea chloroplastic ATP synthase is the source of multiple plant defense elicitors during insect herbivory. <i>Plant Physiology</i> , <b>2007</b> , 144, 793-805	6.6	103
74	Functional analysis of a tomato salicylic acid methyl transferase and its role in synthesis of the flavor volatile methyl salicylate. <i>Plant Journal</i> , <b>2010</b> , 62, 113-23	6.9	98
73	Dynamic Maize Responses to Aphid Feeding Are Revealed by a Time Series of Transcriptomic and Metabolomic Assays. <i>Plant Physiology</i> , <b>2015</b> , 169, 1727-43	6.6	94
72	Accumulation of terpenoid phytoalexins in maize roots is associated with drought tolerance. <i>Plant, Cell and Environment</i> , <b>2015</b> , 38, 2195-207	8.4	88
71	Maize death acids, 9-lipoxygenase-derived cyclopente(a)nones, display activity as cytotoxic phytoalexins and transcriptional mediators. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2015</b> , 112, 11407-12	11.5	83
70	Identification of genes in the phenylalanine metabolic pathway by ectopic expression of a MYB transcription factor in tomato fruit. <i>Plant Cell</i> , <b>2011</b> , 23, 2738-53	11.6	82

69	Effects of elevated [CO <sub>2</sub> ] on maize defence against mycotoxigenic <i>Fusarium verticillioides</i> . <i>Plant, Cell and Environment</i> , <b>2014</b> , 37, 2691-706	8.4	81
68	<i>Physcomitrella patens</i> activates reinforcement of the cell wall, programmed cell death and accumulation of evolutionary conserved defence signals, such as salicylic acid and 12-oxo-phytodienoic acid, but not jasmonic acid, upon <i>Botrytis cinerea</i> infection. <i>Molecular Plant Pathology</i> , <b>2012</b> , 13, 960-74	5.7	78
67	Immunological "Memory" in the Induced Accumulation of Nicotine in Wild Tobacco. <i>Ecology</i> , <b>1996</b> , 77, 236-246	4.6	74
66	Effects of octadecanoid metabolites and inhibitors on induced nicotine accumulation in <i>Nicotiana sylvestris</i> . <i>Journal of Chemical Ecology</i> , <b>1996</b> , 22, 61-74	2.7	72
65	Allocation of nitrogen to an inducible defense and seed production in <i>Nicotiana attenuata</i> . <i>Oecologia</i> , <b>1998</b> , 115, 541-552	2.9	71
64	<i>Pythium</i> infection activates conserved plant defense responses in mosses. <i>Planta</i> , <b>2009</b> , 230, 569-79	4.7	69
63	Coronatine and salicylic acid: the battle between <i>Arabidopsis</i> and <i>Pseudomonas</i> for phytohormone control. <i>Molecular Plant Pathology</i> , <b>2005</b> , 6, 79-83	5.7	69
62	Systemic acquired tolerance to virulent bacterial pathogens in tomato. <i>Plant Physiology</i> , <b>2005</b> , 138, 1481-1490	6.0	65
61	The novel monocot-specific 9-lipoxygenase ZmLOX12 is required to mount an effective jasmonate-mediated defense against <i>Fusarium verticillioides</i> in maize. <i>Molecular Plant-Microbe Interactions</i> , <b>2014</b> , 27, 1263-76	3.6	61
60	Interactions between <i>Spinacia oleracea</i> and <i>Bradysia impatiens</i> : a role for phytoecdysteroids. <i>Archives of Insect Biochemistry and Physiology</i> , <b>2002</b> , 51, 204-21	2.3	61
59	The attraction of <i>Spodoptera frugiperda</i> neonates to cowpea seedlings is mediated by volatiles induced by conspecific herbivory and the elicitor inceptin. <i>Journal of Chemical Ecology</i> , <b>2008</b> , 34, 291-300	2.7	60
58	Homologous RXLR effectors from <i>Hyaloperonospora arabidopsidis</i> and <i>Phytophthora sojae</i> suppress immunity in distantly related plants. <i>Plant Journal</i> , <b>2012</b> , 72, 882-93	6.9	59
57	Discovery, Biosynthesis and Stress-Related Accumulation of Dolabradiene-Derived Defenses in Maize. <i>Plant Physiology</i> , <b>2018</b> , 176, 2677-2690	6.6	55
56	Fungal-induced protein hyperacetylation in maize identified by acetylome profiling. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2018</b> , 115, 210-215	11.5	55
55	Rapidly induced chemical defenses in maize stems and their effects on short-term growth of <i>Ostrinia nubilalis</i> . <i>Journal of Chemical Ecology</i> , <b>2011</b> , 37, 984-91	2.7	55
54	Biosynthesis and function of terpenoid defense compounds in maize ( <i>Zea mays</i> ). <i>Planta</i> , <b>2019</b> , 249, 21-30	4.7	52
53	The effects of climate change associated abiotic stresses on maize phytochemical defenses. <i>Phytochemistry Reviews</i> , <b>2018</b> , 17, 37-49	7.7	51
52	An apoplastic peptide activates salicylic acid signalling in maize. <i>Nature Plants</i> , <b>2018</b> , 4, 172-180	11.5	50

51	Cotton plant, <i>Gossypium hirsutum</i> L., defense in response to nitrogen fertilization. <i>Journal of Chemical Ecology</i> , <b>2008</b> , 34, 1553-64	2.7	49
50	Insect-Induced Synthesis of Phytoecdysteroids in Spinach, <i>Spinacia oleracea</i> . <i>Journal of Chemical Ecology</i> , <b>1999</b> , 25, 1739-1757	2.7	48
49	Impacts of insect oral secretions on defoliation-induced plant defense. <i>Current Opinion in Insect Science</i> , <b>2015</b> , 9, 7-15	5.1	47
48	The maize viviparous15 locus encodes the molybdopterin synthase small subunit. <i>Plant Journal</i> , <b>2006</b> , 45, 264-74	6.9	45
47	Cell wall invertase-deficient miniature1 kernels have altered phytohormone levels. <i>Phytochemistry</i> , <b>2008</b> , 69, 692-9	4	39
46	European corn borer ( <i>Ostrinia nubilalis</i> ) induced responses enhance susceptibility in maize. <i>PLoS ONE</i> , <b>2013</b> , 8, e73394	3.7	39
45	Selinene Volatiles Are Essential Precursors for Maize Defense Promoting Fungal Pathogen Resistance. <i>Plant Physiology</i> , <b>2017</b> , 175, 1455-1468	6.6	38
44	An amino acid substitution inhibits specialist herbivore production of an antagonist effector and recovers insect-induced plant defenses. <i>Plant Physiology</i> , <b>2012</b> , 160, 1468-78	6.6	38
43	The maize Viviparous10/Viviparous13 locus encodes the Cnx1 gene required for molybdenum cofactor biosynthesis. <i>Plant Journal</i> , <b>2006</b> , 45, 250-63	6.9	38
42	Multiple genes recruited from hormone pathways partition maize diterpenoid defences. <i>Nature Plants</i> , <b>2019</b> , 5, 1043-1056	11.5	36
41	Development of a lesion-mimic phenotype in a transgenic wheat line overexpressing genes for pathogenesis-related (PR) proteins is dependent on salicylic acid concentration. <i>Molecular Plant-Microbe Interactions</i> , <b>2003</b> , 16, 916-25	3.6	35
40	Insect-induced daidzein, formononetin and their conjugates in soybean leaves. <i>Metabolites</i> , <b>2014</b> , 4, 532-46	3.6	33
39	A receptor-like protein mediates plant immune responses to herbivore-associated molecular patterns. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2020</b> , 117, 31510-31518	11.5	33
38	Damage-Induced Accumulation of Phytoecdysteroids in Spinach: A Rapid Root Response Involving the Octadecanoic Acid Pathway. <i>Journal of Chemical Ecology</i> , <b>1998</b> , 24, 339-360	2.7	32
37	Interactive Effects of Elevated [CO <sub>2</sub> ] and Drought on the Maize Phytochemical Defense Response against Mycotoxigenic <i>Fusarium verticillioides</i> . <i>PLoS ONE</i> , <b>2016</b> , 11, e0159270	3.7	29
36	Accumulation of 5-hydroxynorvaline in maize ( <i>Zea mays</i> ) leaves is induced by insect feeding and abiotic stress. <i>Journal of Experimental Botany</i> , <b>2015</b> , 66, 593-602	7	28
35	Commercial hybrids and mutant genotypes reveal complex protective roles for inducible terpenoid defenses in maize. <i>Journal of Experimental Botany</i> , <b>2018</b> , 69, 1693-1705	7	26
34	Spatial patterns of aflatoxin levels in relation to ear-feeding insect damage in pre-harvest corn. <i>Toxins</i> , <b>2011</b> , 3, 920-31	4.9	26

33	Activation of Shikimate, Phenylpropanoid, Oxylipins, and Auxin Pathways in <i>Pectobacterium carotovorum</i> Elicitors-Treated Moss. <i>Frontiers in Plant Science</i> , <b>2016</b> , 7, 328	6.2	26
32	Ethylene signaling regulates natural variation in the abundance of antifungal acetylated diferuloylsucroses and <i>Fusarium graminearum</i> resistance in maize seedling roots. <i>New Phytologist</i> , <b>2019</b> , 221, 2096-2111	9.8	26
31	Effects of soldier-derived terpenes on soldier caste differentiation in the termite <i>Reticulitermes flavipes</i> . <i>Journal of Chemical Ecology</i> , <b>2009</b> , 35, 256-64	2.7	25
30	Synthesis of caeliferins, elicitors of plant immune responses: accessing lipophilic natural products via cross metathesis. <i>Organic Letters</i> , <b>2011</b> , 13, 5900-3	6.2	23
29	Genetic elucidation of interconnected antibiotic pathways mediating maize innate immunity. <i>Nature Plants</i> , <b>2020</b> , 6, 1375-1388	11.5	23
28	Tissue-specific PhBPBT expression is differentially regulated in response to endogenous ethylene. <i>Journal of Experimental Botany</i> , <b>2008</b> , 59, 609-18	7	20
27	Soldier caste influences on candidate primer pheromone levels and juvenile hormone-dependent caste differentiation in workers of the termite <i>Reticulitermes flavipes</i> . <i>Journal of Insect Physiology</i> , <b>2011</b> , 57, 771-7	2.4	19
26	Phytohormones mediate volatile emissions during the interaction of compatible and incompatible pathogens: the role of ethylene in <i>Pseudomonas syringae</i> infected tobacco. <i>Journal of Chemical Ecology</i> , <b>2005</b> , 31, 439-59	2.7	19
25	Phytoecdysteroid Turnover in Spinach: Long-term Stability Supports a Plant Defense Hypothesis. <i>Journal of Chemical Ecology</i> , <b>2000</b> , 26, 2883-2896	2.7	19
24	Functional Characterization of Two Class II Diterpene Synthases Indicates Additional Specialized Diterpenoid Pathways in Maize (). <i>Frontiers in Plant Science</i> , <b>2018</b> , 9, 1542	6.2	18
23	Head-group acylation of monogalactosyldiacylglycerol is a common stress response, and the acyl-galactose acyl composition varies with the plant species and applied stress. <i>Physiologia Plantarum</i> , <b>2014</b> , 150, 517-28	4.6	15
22	Phenolic Compounds Accumulate Specifically in Maternally-Derived Tissues of Developing Maize Kernels. <i>Cereal Chemistry</i> , <b>2007</b> , 84, 350-356	2.4	15
21	Fungal and herbivore elicitation of the novel maize sesquiterpenoid, zealexin A4, is attenuated by elevated CO. <i>Planta</i> , <b>2018</b> , 247, 863-873	4.7	13
20	The Arabidopsis MAP kinase kinase 7: A crosstalk point between auxin signaling and defense responses?. <i>Plant Signaling and Behavior</i> , <b>2008</b> , 3, 272-4	2.5	13
19	Inducible De Novo Biosynthesis of Isoflavonoids in Soybean Leaves by <i>Spodoptera litura</i> Derived Elicitors: Tracer Techniques Aided by High Resolution LCMS. <i>Journal of Chemical Ecology</i> , <b>2016</b> , 42, 1226-1236	2.7	12
18	A maize death acid, 10-oxo-11-phytoenoic acid, is the predominant cyclopentenone signal present during multiple stress and developmental conditions. <i>Plant Signaling and Behavior</i> , <b>2016</b> , 11, e1120395	2.5	11
17	A fragmentation study of isoflavones by IT-TOF-MS using biosynthesized isotopes. <i>Bioscience, Biotechnology and Biochemistry</i> , <b>2018</b> , 82, 1309-1315	2.1	8
16	Biosynthetic pathway of aliphatic formates via a Baeyer-Villiger oxidation in mechanism present in astigmatid mites. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2017</b> , 114, 2616-2621	11.5	6

15	A receptor for herbivore-associated molecular patterns mediates plant immunity		6
14	Biosynthesis and antifungal activity of fungus-induced O-methylated flavonoids in maize. <i>Plant Physiology</i> , <b>2021</b> ,	6.6	4
13	Survey of Sensitivity to Fatty Acid-Amino Acid Conjugates in the Solanaceae. <i>Journal of Chemical Ecology</i> , <b>2020</b> , 46, 330-343	2.7	3
12	Influence of brown stink bug feeding, planting date and sampling time on common smut infection of maize. <i>Insect Science</i> , <b>2014</b> , 21, 564-71	3.6	3
11	Comparative analyses of responses to exogenous and endogenous antiherbivore elicitors enable a forward genetics approach to identify maize gene candidates mediating sensitivity to herbivore-associated molecular patterns. <i>Plant Journal</i> , <b>2021</b> , 108, 1295-1316	6.9	3
10	Seed Treatment with Live or Dead <i>Fusarium verticillioides</i> Equivalently Reduces the Severity of Subsequent Stalk Rot. <i>Journal of Phytopathology</i> , <b>2014</b> , 162, 201-204	1.8	2
9	Evaluation of spatial and temporal patterns of insect damage and aflatoxin level in the pre-harvest corn fields to improve management tactics. <i>Insect Science</i> , <b>2014</b> , 21, 572-83	3.6	2
8	Plant height heterosis is quantitatively associated with expression levels of plastid ribosomal proteins. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2021</b> , 118,	11.5	2
7	Plant trait heterosis is quantitatively associated with expression heterosis of the plastid ribosomal proteins		2
6	Getting back to the grass roots: harnessing specialized metabolites for improved crop stress resilience. <i>Current Opinion in Biotechnology</i> , <b>2021</b> , 70, 174-186	11.4	2
5	Genetic elucidation of complex biochemical traits mediating maize innate immunity		1
4	A sorghum genome-wide association study (GWAS) identifies a WRKY transcription factor as a candidate gene underlying sugarcane aphid ( <i>Melanaphis sacchari</i> ) resistance.. <i>Planta</i> , <b>2022</b> , 255, 37	4.7	0
3	Induces Maize-Derived Ethylene to Promote Virulence by Engaging Fungal G-Protein Signaling. <i>Molecular Plant-Microbe Interactions</i> , <b>2021</b> , 34, 1157-1166	3.6	0
2	Synthesis and Determination of Absolute Configuration of Zealexin A1, a Sesquiterpenoid Phytoalexin from <i>Zea mays</i> . <i>European Journal of Organic Chemistry</i> , <b>2021</b> , 2021, 1174-1178	3.2	0
1	Efficient synthesis of zealexin B1, a maize sesquiterpenoid phytoalexin, via Suzuki-Miyaura coupling. <i>Tetrahedron Letters</i> , <b>2022</b> , 91, 153641		2