Anna Perri

List of Publications by Citations

Source: https://exaly.com/author-pdf/3761108/anna-perri-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

53 969 19 30 g-index

54 1,082 4.4 3.68 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
53	Missense mutation in the transcription factor NKX2-5: a novel molecular event in the pathogenesis of thyroid dysgenesis. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2006 , 91, 1428-33	5.6	132
52	Congenital hypothyroidism with impaired thyroid response to thyrotropin (TSH) and absent circulating thyroglobulin: evidence for a new inactivating mutation of the TSH receptor gene. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2000 , 85, 1001-8	5.6	65
51	Low prevalence of thyrotropin receptor mutations in a large series of subjects with sporadic and familial nonautoimmune subclinical hypothyroidism. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2004 , 89, 5787-93	5.6	55
50	Congenital Hypothyroidism with Impaired Thyroid Response to Thyrotropin (TSH) and Absent Circulating Thyroglobulin: Evidence for a New Inactivating Mutation of the TSH Receptor Gene. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2000 , 85, 1001-1008	5.6	51
49	Activating Thyrotropin Receptor Mutations Are Present in Nonadenomatous Hyperfunctioning Nodules of Toxic or Autonomous Multinodular Goiter. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2000 , 85, 2270-2274	5.6	49
48	High doses of hydroxytyrosol induce apoptosis in papillary and follicular thyroid cancer cells. <i>Journal of Endocrinological Investigation</i> , 2017 , 40, 153-162	5.2	42
47	Benign nonfunctioning thyroid adenomas are characterized by a defective targeting to cell membrane or a reduced expression of the sodium iodide symporter protein. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2002 , 87, 352-7	5.6	41
46	Sporadic nonautoimmune congenital hyperthyroidism due to a strong activating mutation of the thyrotropin receptor gene. <i>Thyroid</i> , 2000 , 10, 859-63	6.2	40
45	Epigallocatechin gallate inhibits growth and epithelial-to-mesenchymal transition in human thyroid carcinoma cell lines. <i>Journal of Cellular Physiology</i> , 2013 , 228, 2054-62	7	38
44	Genetic analysis of the PAX8 gene in children with congenital hypothyroidism and dysgenetic or eutopic thyroid glands: identification of a novel sequence variant. <i>Clinical Endocrinology</i> , 2007 , 67, 34-4	10 ^{3.4}	36
43	Thyroid resistance to TSH complicated by autoimmune thyroiditis. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 4543-6	5.6	32
42	Adiponectin is expressed and secreted by renal tubular epithelial cells. <i>Journal of Nephrology</i> , 2013 , 26, 1049-54	4.8	29
41	Rapamycin-induced autophagy protects proximal tubular renal cells against proteinuric damage through the transcriptional activation of the nerve growth factor receptor NGFR. <i>Autophagy</i> , 2018 , 14, 1028-1042	10.2	26
40	Identification of TSH receptor mutations in three families with resistance to TSH. <i>Clinical Endocrinology</i> , 2007 , 67, 712-8	3.4	26
39	Oxidative Balance and Inflammation in Hemodialysis Patients: Biomarkers of Cardiovascular Risk?. Oxidative Medicine and Cellular Longevity, 2019 , 2019, 8567275	6.7	25
38	Genetic analysis of TTF-2 gene in children with congenital hypothyroidism and cleft palate, congenital hypothyroidism, or isolated cleft palate. <i>Thyroid</i> , 2004 , 14, 584-8	6.2	22
37	Functioning and Nonfunctioning Thyroid Adenomas Involve Different Molecular Pathogenetic Mechanisms. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1999 , 84, 4155-4158	5.6	20

(2016-2003)

Congenital hypothyroidism due to a new deletion in the sodium/iodide symporter protein. <i>Clinical Endocrinology</i> , 2003 , 59, 500-6	3.4	19
Functioning and nonfunctioning thyroid adenomas involve different molecular pathogenetic mechanisms. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1999 , 84, 4155-8	5.6	19
Exposure to nerve growth factor worsens nephrotoxic effect induced by Cyclosporine A in HK-2 cells. <i>PLoS ONE</i> , 2013 , 8, e80113	3.7	17
TSH receptor and Gs(alpha) genetic analysis in children with Down's syndrome and subclinical hypothyroidism. <i>Journal of Endocrinological Investigation</i> , 2003 , 26, 997-1000	5.2	17
T3 enhances thyroid cancer cell proliferation through TRII/Oct-1-mediated cyclin D1 activation. <i>Molecular and Cellular Endocrinology</i> , 2014 , 382, 205-217	4.4	16
Beneficial effects of iodized salt prophylaxis on thyroid volume in an iodine deficient area of southern Italy. <i>Clinical Endocrinology</i> , 2009 , 71, 124-9	3.4	16
Adiponectin secreted by tubular renal cells during LPS exposure worsens the cellular inflammatory damage. <i>Journal of Nephrology</i> , 2016 , 29, 185-194	4.8	15
Olive leaf extract counteracts epithelial to mesenchymal transition process induced by peritoneal dialysis, through the inhibition of TGFI signaling. <i>Cell Biology and Toxicology</i> , 2019 , 35, 95-109	7.4	11
Soluble Klotho levels in adult renal transplant recipients are modulated by recombinant human erythropoietin. <i>Journal of Nephrology</i> , 2014 , 27, 577-85	4.8	11
Thyroid Resistance to TSH Complicated by Autoimmune Thyroiditis		10
Early cytomegalovirus-specific T-cell response and estimated glomerular filtration rate identify patients at high risk of infection after renal transplantation. <i>Transplant Infectious Disease</i> , 2016 , 18, 1919	-2071	10
Nerve growth factor exposure promotes tubular epithelial-mesenchymal transition via TGF-II signaling activation. <i>Growth Factors</i> , 2015 , 33, 169-80	1.6	9
Efficacy of eculizumab in severe ADAMTS13-deficient thrombotic thrombocytopenic purpura (TTP)		
refractory to standard therapies. <i>Transfusion and Apheresis Science</i> , 2018 , 57, 247-249	2.4	9
An Olive Leaf Extract Rich in Polyphenols Promotes Apoptosis in Cervical Cancer Cells by Upregulating p21 Gene Expression. <i>Nutrition and Cancer</i> , 2019 , 71, 320-333	2.4	8
An Olive Leaf Extract Rich in Polyphenols Promotes Apoptosis in Cervical Cancer Cells by	·	
An Olive Leaf Extract Rich in Polyphenols Promotes Apoptosis in Cervical Cancer Cells by Upregulating p21 Gene Expression. <i>Nutrition and Cancer</i> , 2019 , 71, 320-333 Erectile Dysfunction after Kidney Transplantation. <i>Journal of Clinical Medicine</i> , 2020 , 9, Monitoring the effects of iodine prophylaxis in the adult population of southern Italy with deficient	2.8	8
An Olive Leaf Extract Rich in Polyphenols Promotes Apoptosis in Cervical Cancer Cells by Upregulating p21 Gene Expression. <i>Nutrition and Cancer</i> , 2019 , 71, 320-333 Erectile Dysfunction after Kidney Transplantation. <i>Journal of Clinical Medicine</i> , 2020 , 9, Monitoring the effects of iodine prophylaxis in the adult population of southern Italy with deficient and sufficient iodine intake levels: a cross-sectional, epidemiological study. <i>British Journal of Nutrition</i> , 2017 , 117, 170-175	2.8	8
	Functioning and nonfunctioning thyroid adenomas involve different molecular pathogenetic mechanisms. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1999, 84, 4155-8 Exposure to nerve growth factor worsens nephrotoxic effect induced by Cyclosporine A in HK-2 cells. <i>PLoS ONE</i> , 2013, 8, e80113 TSH receptor and Gs(alpha) genetic analysis in children with Down's syndrome and subclinical hypothyroidism. <i>Journal of Endocrinological Investigation</i> , 2003, 26, 997-1000 T3 enhances thyroid cancer cell proliferation through TRil/Oct-1-mediated cyclin D1 activation. <i>Molecular and Cellular Endocrinology</i> , 2014, 382, 205-217 Beneficial effects of iodized salt prophylaxis on thyroid volume in an iodine deficient area of southern Italy. <i>Clinical Endocrinology</i> , 2009, 71, 124-9 Adiponectin secreted by tubular renal cells during LPS exposure worsens the cellular inflammatory damage. <i>Journal of Nephrology</i> , 2016, 29, 185-194 Olive leaf extract counteracts epithelial to mesenchymal transition process induced by peritoneal dialysis, through the inhibition of TGFil signaling. <i>Cell Biology and Toxicology</i> , 2019, 35, 95-109 Soluble Klotho levels in adult renal transplant recipients are modulated by recombinant human erythropoietin. <i>Journal of Nephrology</i> , 2014, 27, 577-85 Thyroid Resistance to TSH Complicated by Autoimmune Thyroiditis Early cytomegalovirus-specific T-cell response and estimated glomerular filtration rate identify patients at high risk of infection after renal transplantation. <i>Transplant Infectious Disease</i> , 2016, 18, 191 Nerve growth factor exposure promotes tubular epithelial-mesenchymal transition via TGF-II signaling activation. <i>Growth Factors</i> , 2015, 33, 169-80	Functioning and nonfunctioning thyroid adenomas involve different molecular pathogenetic mechanisms. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1999, 84, 4155-8 Exposure to nerve growth factor worsens nephrotoxic effect induced by Cyclosporine A in HK-2 cells. <i>PLoS ONE</i> , 2013, 8, e80113 37 TSH receptor and Gs(alpha) genetic analysis in children with Down's syndrome and subclinical hypothyroidism. <i>Journal of Endocrinological Investigation</i> , 2003, 26, 997-1000 52 T3 enhances thyroid cancer cell proliferation through TR®/Oct-1-mediated cyclin D1 activation. <i>Molecular and Cellular Endocrinology</i> , 2014, 382, 205-217 44 Beneficial effects of iodized salt prophylaxis on thyroid volume in an iodine deficient area of southern Italy. <i>Clinical Endocrinology</i> , 2009, 71, 124-9 Adiponectin secreted by tubular renal cells during LPS exposure worsens the cellular inflammatory damage. <i>Journal of Nephrology</i> , 2016, 29, 185-194 Olive leaf extract counteracts epithelial to mesenchymal transition process induced by peritoneal dialysis, through the inhibition of TGF® signaling. <i>Cell Biology and Toxicology</i> , 2019, 35, 95-109 74 Soluble Klotho levels in adult renal transplant recipients are modulated by recombinant human erythropoietin. <i>Journal of Nephrology</i> , 2014, 27, 577-85 Thyroid Resistance to TSH Complicated by Autoimmune Thyroiditis Early cytomegalovirus-specific T-cell response and estimated glomerular filtration rate identify patients at high risk of infection after renal transplantation. <i>Transplant Infectious Disease</i> , 2016, 18, 191-201 Nerve growth factor exposure promotes tubular epithelial-mesenchymal transition via TGF-® signaling activation. <i>Growth Factors</i> , 2015, 33, 169-80

18	Active compounds extracted from extra virgin olive oil counteract mesothelial-to-mesenchymal transition of peritoneal mesothelium cells exposed to conventional peritoneal dialysate: in vitro and in vivo evidences. <i>Journal of Nephrology</i> , 2017 , 30, 841-850	4.8	5
17	High nerve growth factor blood concentration in renal transplantation: a new prognostic marker?. <i>Transplantation Proceedings</i> , 2013 , 45, 2654-6	1.1	4
16	The Risky Health Behaviours of Male Adolescents in the Southern Italian Region: Implications for Sexual and Reproductive Disease. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	3
15	Proinflammatory profile of visceral adipose tissue and oxidative stress in severe obese patients carrying the variant rs4612666 C of NLRP3 gene. <i>Minerva Endocrinology</i> , 2021 , 46, 309-316	2.5	3
14	Rapamycin promotes autophagy cell death of Kaposi's sarcoma cells through P75NTR activation. <i>Experimental Dermatology</i> , 2021 ,	4	3
13	Sequencing of the entire coding region of the receptor associated protein (RAP) in patients with primary hypothyroidism of unknown origin. <i>Journal of Endocrinological Investigation</i> , 2007 , 30, 839-43	5.2	2
12	Mean Platelet Volume as a Marker of Vasculogenic Erectile Dysfunction and Future Cardiovascular Risk. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	2
11	Olive leaf extract counteracts cell proliferation and cyst growth in an in vitro model of autosomal dominant polycystic kidney disease. <i>Food and Function</i> , 2018 , 9, 5925-5935	6.1	2
10	Lexicon of pandemics: a semantic analysis of the Spanish flu and the Covid-19 timeframe terminology. <i>Journal of Documentation</i> , 2021 , ahead-of-print,	1.3	1
9	NFKB1 promoter polymorphism: A new predictive marker of cytomegalovirus infection after kidney transplantation. <i>Transplant Infectious Disease</i> , 2019 , 21, e13027	2.7	1
8	Analysis on chronic pain management: Focus on the Italian network. <i>International Journal of Health Planning and Management</i> , 2021 , 36, 151-157	2.2	1
7	Identification of a New Complement Factor H Mutation in a Patient With Pregnancy-Related Acute Kidney Injury. <i>Kidney International Reports</i> , 2020 , 5, 1603-1607	4.1	O
6	Oleuropein Counteracts Both the Proliferation and Migration of Intra- and Extragonadal Seminoma Cells. <i>Nutrients</i> , 2022 , 14, 2323	6.7	O
5	FP028NERVE GROWTH FACTOR EXPOSURE PROMOTES TUBULAR EPITHELIAL-MESENCHYMAL-TRANSITION VIA TGF® SIGNALING ACTIVATION. <i>Nephrology Dialysis Transplantation</i> , 2015 , 30, iii74-iii74	4.3	
4	SP035IDENTIFICATION AND FUNCTIONAL STUDY OF A HETEROZYGOUS MISSENSE MUTATION IN UMOD GENE IN AN ITALIAN FAMILY AFFECTED BY MEDULLARY CYSTIC KIDNEY DISEASE TYPE II. Nephrology Dialysis Transplantation, 2015 , 30, iii392-iii392	4.3	
3	SP797TREATMENT OF EARLY SUBCLINICAL REJECTION IS ASSOCIATED TO GRAFT SURVIVAL IMPROVEMENT: 10 YEARS SINGLE CENTER EXPERIENCE. <i>Nephrology Dialysis Transplantation</i> , 2015 , 30, iii641-iii641	4.3	
2	FP024ADIPONECTIN SECRETED BY TUBULAR RENAL CELLS MEDIATES THE CELLULAR INFLAMMATORY DAMAGE LPS-INDUCED IN AN AUTOCRINE DEPENDENT-MANNER. <i>Nephrology Dialysis Transplantation</i> , 2015 , 30, iii73-iii73	4.3	
1	Effects of nutraceuticals on sexual satisfaction and lower urinary tract symptoms in a cohort of young-old men. <i>Phytotherapy Research</i> , 2018 , 32, 284-289	6.7	