

Paulo Belmonte-de-Abreu

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3761663/paulo-belmonte-de-abreu-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

103
papers

3,168
citations

31
h-index

52
g-index

113
ext. papers

3,488
ext. citations

4.2
avg. IF

4.52
L-index

| # | Paper | IF | Citations |
|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 103 | Differential Physical and Mental Benefits of Physiotherapy Program Among Patients With Schizophrenia and Healthy Controls Suggesting Different Physical Characteristics and Needs. <i>Frontiers in Psychiatry</i> , 2021 , 12, 536767 | 5 | |
| 102 | Proinflammatory and anti-inflammatory biomarkers in schizophrenia and influence of simvastatin on the interleukin-6. <i>International Clinical Psychopharmacology</i> , 2019 , 34, 84-88 | 2.2 | 5 |
| 101 | Postural changes in different stages of schizophrenia is associated with inflammation and pain: a cross-sectional observational study. <i>International Journal of Psychiatry in Clinical Practice</i> , 2017 , 21, 104-111 | 2.1 | 5 |
| 100 | Effects of repetitive transcranial magnetic stimulation over supplementary motor area in patients with schizophrenia with obsessive-compulsive-symptoms: A pilot study. <i>Psychiatry Research</i> , 2016 , 242, 34-38 | 9.9 | 8 |
| 99 | Generation of iPS cell lines from schizophrenia patients using a non-integrative method. <i>Stem Cell Research</i> , 2016 , 17, 97-101 | 1.6 | 4 |
| 98 | Trends in elderly psychiatric admissions to the Brazilian public health care system. <i>Revista Brasileira De Psiquiatria</i> , 2016 , 38, 314-317 | 2.6 | 3 |
| 97 | Resveratrol Supplementation in Schizophrenia Patients: A Randomized Clinical Trial Evaluating Serum Glucose and Cardiovascular Risk Factors. <i>Nutrients</i> , 2016 , 8, 73 | 6.7 | 33 |
| 96 | Physical Functional Capacity and C-Reactive Protein in Schizophrenia. <i>Frontiers in Psychiatry</i> , 2016 , 7, 131 | 5 | 5 |
| 95 | Resveratrol Supplementation Did Not Improve Cognition in Patients with Schizophrenia: Results from a Randomized Clinical Trial. <i>Frontiers in Psychiatry</i> , 2016 , 7, 159 | 5 | 30 |
| 94 | Persistence and remission of ADHD during adulthood: a 7-year clinical follow-up study. <i>Psychological Medicine</i> , 2015 , 45, 2045-56 | 6.9 | 54 |
| 93 | The effects of sodium nitroprusside treatment on cognitive deficits in schizophrenia: a pilot study. <i>Journal of Clinical Psychopharmacology</i> , 2015 , 35, 83-5 | 1.7 | 22 |
| 92 | Differences in parental bonding between schizophrenia and bipolar disorder: Evidence of prodromal symptoms?. <i>Schizophrenia Research</i> , 2015 , 165, 134-7 | 3.6 | 4 |
| 91 | Co-expression network of neural-differentiation genes shows specific pattern in schizophrenia. <i>BMC Medical Genomics</i> , 2015 , 8, 23 | 3.7 | 31 |
| 90 | Innate immune response is differentially dysregulated between bipolar disease and schizophrenia. <i>Schizophrenia Research</i> , 2015 , 161, 215-21 | 3.6 | 46 |
| 89 | New-onset panic attacks after deep brain stimulation of the nucleus accumbens in a patient with refractory obsessive-compulsive and bipolar disorders: a case report. <i>Revista Brasileira De Psiquiatria</i> , 2015 , 37, 182-3 | 2.6 | 4 |
| 88 | Further evidence for the association between a polymorphism in the promoter region of SLC6A3/DAT1 and ADHD: findings from a sample of adults. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2014 , 264, 401-8 | 5.1 | 20 |
| 87 | P.3.f.011 C-reactive protein associated to illness severity in schizophrenia. <i>European Neuropsychopharmacology</i> , 2014 , 24, S572 | 1.2 | |

| | | | |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 86 | Differences in eotaxin serum levels patients with recent onset and in chronic stable schizophrenia: a clue for understanding accelerating aging profile. <i>Schizophrenia Research</i> , 2014 , 152, 528-9 | 3.6 | 18 |
| 85 | BDNF: a biomarker for social vulnerability in individuals diagnosed with gender dysphoria. <i>Journal of Psychiatric Research</i> , 2014 , 50, 16-7 | 5.2 | 3 |
| 84 | Sodium nitroprusside treatment of clozapine-refractory schizophrenia. <i>Journal of Clinical Psychopharmacology</i> , 2014 , 34, 761-3 | 1.7 | 12 |
| 83 | Could comorbid bipolar disorder account for a significant share of executive function deficits in adults with attention-deficit hyperactivity disorder?. <i>Bipolar Disorders</i> , 2014 , 16, 270-6 | 3.8 | 12 |
| 82 | Severity but not comorbidities predicts response to methylphenidate in adults with attention-deficit/hyperactivity disorder: results from a naturalistic study. <i>Journal of Clinical Psychopharmacology</i> , 2014 , 34, 212-7 | 1.7 | 13 |
| 81 | Discordant transsexualism in male monozygotic twins: neuroanatomical and psychological differences. <i>Archives of Sexual Behavior</i> , 2014 , 43, 399-405 | 3.5 | 2 |
| 80 | ADHD diagnosis may influence the association between polymorphisms in nicotinic acetylcholine receptor genes and tobacco smoking. <i>NeuroMolecular Medicine</i> , 2014 , 16, 389-97 | 4.6 | 16 |
| 79 | Verbal memory impairment in healthy siblings of patients with schizophrenia. <i>Schizophrenia Research</i> , 2013 , 150, 580-2 | 3.6 | 13 |
| 78 | Rapid therapeutic response to anodal tDCS of right dorsolateral prefrontal cortex in acute mania. <i>Brain Stimulation</i> , 2013 , 6, 701-3 | 5.1 | 20 |
| 77 | Serum concentrations of brain-derived neurotrophic factor in patients with gender identity disorder. <i>Journal of Psychiatric Research</i> , 2013 , 47, 1546-8 | 5.2 | 15 |
| 76 | The role of a mineralocorticoid receptor gene functional polymorphism in the symptom dimensions of persistent ADHD. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2013 , 263, 181-8 | 5.1 | 15 |
| 75 | Rapid improvement of acute schizophrenia symptoms after intravenous sodium nitroprusside: a randomized, double-blind, placebo-controlled trial. <i>JAMA Psychiatry</i> , 2013 , 70, 668-76 | 14.5 | 102 |
| 74 | The CYP1A2 -163C>A polymorphism is associated with clozapine-induced generalized tonic-clonic seizures in Brazilian schizophrenia patients. <i>Psychiatry Research</i> , 2013 , 209, 242-5 | 9.9 | 19 |
| 73 | Cognitive deficits in adults with ADHD go beyond comorbidity effects. <i>Journal of Attention Disorders</i> , 2013 , 17, 483-8 | 3.7 | 22 |
| 72 | rTMS as an add-on treatment for resistant obsessive-compulsive symptoms in patients with schizophrenia: report of three cases. <i>Revista Brasileira De Psiquiatria</i> , 2013 , 35, 210-1 | 2.6 | 1 |
| 71 | The role of BDNF in the pathophysiology and treatment of schizophrenia. <i>Journal of Psychiatric Research</i> , 2012 , 46, 1-11 | 5.2 | 137 |
| 70 | Similarities in serum oxidative stress markers and inflammatory cytokines in patients with overt schizophrenia at early and late stages of chronicity. <i>Journal of Psychiatric Research</i> , 2012 , 46, 819-24 | 5.2 | 118 |
| 69 | The burdened life of adults with ADHD: impairment beyond comorbidity. <i>European Psychiatry</i> , 2012 , 27, 309-13 | 6 | 43 |

| | | | |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 68 | Does age of onset of impairment impact on neuropsychological and personality features of adult ADHD?. <i>Journal of Psychiatric Research</i> , 2012 , 46, 1307-11 | 5.2 | 10 |
| 67 | Gene expression of peripheral blood lymphocytes may discriminate patients with schizophrenia from controls. <i>Psychiatry Research</i> , 2012 , 200, 1018-21 | 9.9 | 14 |
| 66 | Reduced serum non-enzymatic antioxidant defense and increased lipid peroxidation in schizophrenic patients on a hypocaloric diet. <i>Neuroscience Letters</i> , 2012 , 512, 43-7 | 3.3 | 3 |
| 65 | Do genes and environment meet to regulate cerebrospinal fluid dynamics? Relevance for schizophrenia. <i>Frontiers in Cellular Neuroscience</i> , 2012 , 6, 31 | 6.1 | 14 |
| 64 | Is there a role for rare variants in DRD4 gene in the susceptibility for ADHD? Searching for an effect of allelic heterogeneity. <i>Molecular Psychiatry</i> , 2012 , 17, 520-6 | 15.1 | 23 |
| 63 | No significant association between genetic variants in 7 candidate genes and response to methylphenidate treatment in adult patients with ADHD. <i>Journal of Clinical Psychopharmacology</i> , 2012 , 32, 820-3 | 1.7 | 28 |
| 62 | Altered oxygen metabolism associated to neurogenesis of induced pluripotent stem cells derived from a schizophrenic patient. <i>Cell Transplantation</i> , 2012 , 21, 1547-59 | 4 | 112 |
| 61 | The role of a lifetime history of oppositional defiant and conduct disorders in adults with ADHD: implications for clinical practice. <i>CNS Spectrums</i> , 2012 , 17, 94-9 | 1.8 | 5 |
| 60 | Serum brain-derived neurotrophic factor and clozapine daily dose in patients with schizophrenia: a positive correlation. <i>Neuroscience Letters</i> , 2011 , 491, 207-10 | 3.3 | 54 |
| 59 | Increased annexin-V and decreased TNF- β serum levels in chronic-medicated patients with schizophrenia. <i>Neuroscience Letters</i> , 2011 , 502, 143-6 | 3.3 | 26 |
| 58 | ADRA2A polymorphisms and ADHD in adults: possible mediating effect of personality. <i>Psychiatry Research</i> , 2011 , 186, 345-50 | 9.9 | 16 |
| 57 | A pilot double-blind sham-controlled trial of repetitive transcranial magnetic stimulation for patients with refractory schizophrenia treated with clozapine. <i>Psychiatry Research</i> , 2011 , 188, 203-7 | 9.9 | 54 |
| 56 | Long-acting injectable risperidone in partially adherent and nonadherent patients with schizophrenia. <i>Neuropsychiatric Disease and Treatment</i> , 2011 , 7, 391-8 | 3.1 | 6 |
| 55 | Smoking and ADHD: an evaluation of self medication and behavioral disinhibition models based on comorbidity and personality patterns. <i>Journal of Psychiatric Research</i> , 2011 , 45, 829-34 | 5.2 | 25 |
| 54 | Adrenergic α A receptor gene is not associated with methylphenidate response in adults with ADHD. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2011 , 261, 205-11 | 5.1 | 22 |
| 53 | Shortage and underutilization of psychiatric beds in southern Brazil: independent data of Brazilian mental health reform. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2011 , 46, 425-9 | 4.5 | 5 |
| 52 | Serum levels of IL-6, IL-10 and TNF- α ; in patients with bipolar disorder and schizophrenia: differences in pro- and anti-inflammatory balance. <i>Revista Brasileira De Psiquiatria</i> , 2011 , 33, 268-274 | 2.6 | 93 |
| 51 | Improvement of schizophrenia with electroconvulsive therapy and serum brain-derived neurotrophic factor levels: Lack of association in a pilot study. <i>Psychiatry and Clinical Neurosciences</i> , 2010 , 64, 663-5 | 6.2 | 10 |

| | | | |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----|
| 50 | Response to methylphenidate is not influenced by DAT1 polymorphisms in a sample of Brazilian adult patients with ADHD. <i>Journal of Neural Transmission</i> , 2010 , 117, 269-76 | 4.3 | 21 |
| 49 | Influence of serotonin transporter gene polymorphisms on clozapine response in Brazilian schizophrenics. <i>Journal of Psychiatric Research</i> , 2010 , 44, 1158-62 | 5.2 | 25 |
| 48 | Improvement of schizophrenia negative and positive symptoms with amantadine as add-on therapy to antipsychotics: a case series. <i>Revista Brasileira De Psiquiatria</i> , 2010 , 32, 193-4 | 2.6 | 1 |
| 47 | Avaliaçã antropomãrica e bioquãmica em pacientes com esquizofrenia usuãrios de clozapina. <i>Revista De Nutricao</i> , 2009 , 22, 697-705 | 1.8 | 2 |
| 46 | Molecular diversity at the CYP2D6 locus in healthy and schizophrenic southern Brazilians. <i>Pharmacogenomics</i> , 2009 , 10, 1457-66 | 2.6 | 26 |
| 45 | Inattention and hyperactivity dimensions of ADHD are associated with different personality profiles. <i>Psychopathology</i> , 2009 , 42, 108-12 | 3.4 | 46 |
| 44 | Improved stress response in bipolar affective disorder with adjunctive spironolactone (mineralocorticoid receptor antagonist): case series. <i>Journal of Psychopharmacology</i> , 2009 , 23, 985-7 | 4.6 | 26 |
| 43 | Late-onset ADHD in adults: milder, but still dysfunctional. <i>Journal of Psychiatric Research</i> , 2009 , 43, 697-701 | 5.0 | 38 |
| 42 | Serum brain-derived neurotrophic factor in bipolar and unipolar depression: a potential adjunctive tool for differential diagnosis. <i>Journal of Psychiatric Research</i> , 2009 , 43, 1200-4 | 5.2 | 100 |
| 41 | The association of child abuse and neglect with adult disability in schizophrenia and the prominent role of physical neglect. <i>Child Abuse and Neglect</i> , 2009 , 33, 618-24 | 4.3 | 58 |
| 40 | Differentiating attention-deficit/hyperactivity disorder inattentive and combined types: a (1)H-magnetic resonance spectroscopy study of fronto-striato-thalamic regions. <i>Journal of Neural Transmission</i> , 2009 , 116, 623-9 | 4.3 | 31 |
| 39 | Serum brain-derived neurotrophic factor (BDNF) is not associated with response to electroconvulsive therapy (ECT): a pilot study in drug resistant depressed patients. <i>Neuroscience Letters</i> , 2009 , 453, 195-8 | 3.3 | 69 |
| 38 | How Hormones Influence Gene Expression: Implications to the Pathophysiology of Schizophrenia. <i>European Psychiatry</i> , 2009 , 24, 1-1 | 6 | |
| 37 | Improvement of negative and positive symptoms in treatment-refractory schizophrenia: a double-blind, randomized, placebo-controlled trial with memantine as add-on therapy to clozapine. <i>Journal of Clinical Psychiatry</i> , 2009 , 70, 1416-23 | 4.6 | 91 |
| 36 | Aripiprazole for patients with schizophrenia and schizoaffective disorder: an open-label, randomized, study versus haloperidol. <i>CNS Spectrums</i> , 2009 , 14, 93-102 | 1.8 | 12 |
| 35 | Reasons for pretreatment attrition and dropout from methylphenidate in adults with attention-deficit/hyperactivity disorder: the role of comorbidities. <i>Journal of Clinical Psychopharmacology</i> , 2009 , 29, 614-6 | 1.7 | 16 |
| 34 | Association of the gene encoding neurogranin with schizophrenia in males. <i>Journal of Psychiatric Research</i> , 2008 , 42, 125-33 | 5.2 | 43 |
| 33 | Elevated serum thiobarbituric acid reactive substances in clinically symptomatic schizophrenic males. <i>Neuroscience Letters</i> , 2008 , 433, 270-3 | 3.3 | 34 |

| | | | |
|----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----|
| 32 | Decreased serum neurotrophin 3 in chronically medicated schizophrenic males. <i>Neuroscience Letters</i> , 2008 , 440, 197-201 | 3.3 | 20 |
| 31 | Serum levels of brain-derived neurotrophic factor in schizophrenia on a hypocaloric diet. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2008 , 32, 1595-8 | 5.5 | 25 |
| 30 | Elevated serum superoxide dismutase and thiobarbituric acid reactive substances in different phases of bipolar disorder and in schizophrenia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2008 , 32, 1677-81 | 5.5 | 158 |
| 29 | G-protein gene 825C>T polymorphism is associated with response to clozapine in Brazilian schizophrenics. <i>Pharmacogenomics</i> , 2008 , 9, 1429-36 | 2.6 | 33 |
| 28 | Smoking is associated with lower performance in WAIS-R Block Design scores in adults with ADHD. <i>Nicotine and Tobacco Research</i> , 2008 , 10, 683-8 | 4.9 | 12 |
| 27 | Naturalistic pharmacogenetic study of treatment resistance to typical neuroleptics in European-Brazilian schizophrenics. <i>Pharmacogenetics and Genomics</i> , 2008 , 18, 599-609 | 1.9 | 34 |
| 26 | Serum levels of brain-derived neurotrophic factor and thiobarbituric acid reactive substances in chronically medicated schizophrenic patients: a positive correlation. <i>Revista Brasileira De Psiquiatria</i> , 2008 , 30, 337-40 | 2.6 | 27 |
| 25 | A twelve-week randomized controlled study of the cognitive-behavioral Integrated Psychological Therapy program: positive effect on the social functioning of schizophrenic patients. <i>Revista Brasileira De Psiquiatria</i> , 2007 , 29, 140-7 | 2.6 | 19 |
| 24 | Transtorno de oposio e desafio e transtorno de conduta: os desfechos no TDAH em adultos. <i>Jornal Brasileiro De Psiquiatria</i> , 2007 , 56, 34-38 | 0.5 | 2 |
| 23 | Ganho de peso e alteraes metablicas em esquizofrenia. <i>Revista De Psiquiatria Clinica</i> , 2007 , 34, 184-188 | 0.8 | 4 |
| 22 | Family-based and case-control studies reveal no association of lipocalin-type prostaglandin D2 synthase with schizophrenia. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2007 , 144B, 642-6 | 3.5 | 3 |
| 21 | Transthyretin: no association between serum levels or gene variants and schizophrenia. <i>Journal of Psychiatric Research</i> , 2007 , 41, 667-72 | 5.2 | 8 |
| 20 | The role of comorbid major depressive disorder in the clinical presentation of adult ADHD. <i>Journal of Psychiatric Research</i> , 2007 , 41, 991-6 | 5.2 | 73 |
| 19 | Serotonin transporter gene polymorphism and the phenotypic heterogeneity of adult ADHD. <i>Journal of Neural Transmission</i> , 2007 , 114, 1631-6 | 4.3 | 20 |
| 18 | Increased serum adenosine deaminase activity in schizophrenic receiving antipsychotic treatment. <i>Neuroscience Letters</i> , 2007 , 414, 61-4 | 3.3 | 22 |
| 17 | Serum levels of brain-derived neurotrophic factor in patients with schizophrenia and bipolar disorder. <i>Neuroscience Letters</i> , 2007 , 420, 45-8 | 3.3 | 123 |
| 16 | Repeated transcranial magnetic stimulation effect over treatment-resistant auditory hallucinations in a female patient with schizophrenia. <i>Revista Brasileira De Psiquiatria</i> , 2007 , 29, 291-2 | 2.6 | 4 |
| 15 | Is age-at-onset criterion relevant for the response to methylphenidate in attention-deficit/hyperactivity disorder?. <i>Journal of Clinical Psychiatry</i> , 2007 , 68, 1109-16 | 4.6 | 31 |

| | | | |
|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----|
| 14 | Evaluation of neuropsychological performance of HIV-infected patients with minor motor cognitive dysfunction treated with highly active antiretroviral therapy. <i>Infection</i> , 2006 , 34, 357-60 | 5.8 | 10 |
| 13 | Elevated serum superoxide dismutase and thiobarbituric acid reactive substances in schizophrenia: a study of patients treated with haloperidol or clozapine. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2006 , 30, 512-5 | 5.5 | 91 |
| 12 | Lack of gender effects on subtype outcomes in adults with attention-deficit/hyperactivity disorder: support for the validity of subtypes. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2006 , 256, 311-9 | 5.1 | 54 |
| 11 | Efficacy of milnacipran in treating anxiety symptoms in schizophrenic patients receiving clozapine: a case series study. <i>Revista Brasileira De Psiquiatria</i> , 2006 , 28, 339-40 | 2.6 | 3 |
| 10 | Increased dyslipidemia in schizophrenic outpatients using new generation antipsychotics. <i>Revista Brasileira De Psiquiatria</i> , 2006 , 28, 301-304 | 2.6 | 9 |
| 9 | Evidências da eficácia da eletroconvulsoterapia na prática psiquiátrica. <i>Revista De Psiquiatria Do Rio Grande Do Sul</i> , 2005 , 27, 302-310 | | 2 |
| 8 | NR4A2 and schizophrenia: lack of association in a Portuguese/Brazilian study. <i>American Journal of Medical Genetics Part A</i> , 2004 , 128B, 41-5 | | 11 |
| 7 | The etiology of schizophrenia and the origin of language: overview of a theory. <i>Comprehensive Psychiatry</i> , 2003 , 44, 7-14 | 7.3 | 46 |
| 6 | The adenosine antagonist theophylline impairs p50 auditory sensory gating in normal subjects. <i>Neuropsychopharmacology</i> , 2002 , 27, 629-37 | 8.7 | 28 |
| 5 | Therapeutic benefit of adjunctive dipyridamole in schizophrenia is probably due to adenosine-glutamate interactions. <i>Journal of Clinical Pharmacy and Therapeutics</i> , 2001 , 26, 155-6 | 2.2 | 11 |
| 4 | Increased serum S100B protein in schizophrenia: a study in medication-free patients. <i>Journal of Psychiatric Research</i> , 2001 , 35, 11-4 | 5.2 | 120 |
| 3 | Neurodevelopmental risk factors in schizophrenia. <i>Brazilian Journal of Medical and Biological Research</i> , 2001 , 34, 155-63 | 2.8 | 9 |
| 2 | Allopurinol augmentation for poorly responsive schizophrenia. <i>International Clinical Psychopharmacology</i> , 2001 , 16, 235-7 | 2.2 | 40 |
| 1 | Allopurinol for refractory aggression and self-inflicted behaviour. <i>Journal of Psychopharmacology</i> , 2000 , 14, 81-3 | 4.6 | 30 |