

# Kirsten Wedervang-Resell

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3763229/publications.pdf>

Version: 2024-02-01

16  
papers

189  
citations

1163117

8  
h-index

1199594

12  
g-index

18  
all docs

18  
docs citations

18  
times ranked

245  
citing authors

#	ARTICLE	IF	CITATIONS
1	Lower circulating neuron-specific enolase concentrations in adults and adolescents with severe mental illness. <i>Psychological Medicine</i> , 2023, 53, 1479-1488.	4.5	6
2	Intracranial and subcortical volumes in adolescents with <sc>early-onset</sc> psychosis: A multisite <sc>mega-analysis</sc> from the <sc>ENIGMA</sc> consortium. <i>Human Brain Mapping</i> , 2022, 43, 373-384.	3.6	27
3	Plasma Levels of the Cytokines B Cell-Activating Factor (BAFF) and A Proliferation-Inducing Ligand (APRIL) in Schizophrenia, Bipolar, and Major Depressive Disorder: A Cross Sectional, Multisite Study. <i>Schizophrenia Bulletin</i> , 2022, 48, 37-46.	4.3	10
4	Increased circulating IL-18 levels in severe mental disorders indicate systemic inflammasome activation. <i>Brain, Behavior, and Immunity</i> , 2022, 99, 299-306.	4.1	33
5	Cognitive impairment profile in adolescent early-onset psychosis using the MATRICS Battery: Age and sex effects.. <i>Neuropsychology</i> , 2021, 35, 300-309.	1.3	6
6	Lower plasma total tau in adolescent psychosis: Involvement of the orbitofrontal cortex. <i>Journal of Psychiatric Research</i> , 2021, 144, 255-261.	3.1	3
7	Aberrant Default Mode Connectivity in Adolescents with Early-Onset Psychosis: A resting state fMRI study. <i>NeuroImage: Clinical</i> , 2021, 33, 102881.	2.7	12
8	Negative Symptom Domains Are Associated With Verbal Learning in Adolescents With Early Onset Psychosis. <i>Frontiers in Psychiatry</i> , 2021, 12, 825681.	2.6	6
9	Lipid alterations in adolescents with early-onset psychosis may be independent of antipsychotic medication. <i>Schizophrenia Research</i> , 2020, 216, 295-301.	2.0	10
10	Increased interleukin 18 activity in adolescents with early-onset psychosis is associated with cortisol and depressive symptoms. <i>Psychoneuroendocrinology</i> , 2020, 112, 104513.	2.7	21
11	Atherogenic Lipid Ratios Related to Myeloperoxidase and C-Reactive Protein Levels in Psychotic Disorders. <i>Frontiers in Psychiatry</i> , 2020, 11, 672.	2.6	15
12	Reduced levels of circulating adhesion molecules in adolescents with early-onset psychosis. <i>NPJ Schizophrenia</i> , 2020, 6, 20.	3.6	10
13	Trajectory and early predictors of apathy development in first-episode psychosis and healthy controls: a 10-year follow-up study. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2020, 270, 709-722.	3.2	9
14	Negative and disorganized symptoms mediate the relationship between verbal learning and global functioning in adolescents with early-onset psychosis. <i>European Child and Adolescent Psychiatry</i> , 2020, 29, 1693-1703.	4.7	7
15	Higher vitamin B12 levels in neurodevelopmental disorders than in healthy controls and schizophrenia. <i>FASEB Journal</i> , 2020, 34, 8114-8124.	0.5	11
16	T8. ATHEROGENIC LIPID RATIOS RELATED TO MYELOPEROXIDASE AND C-REACTIVE PROTEIN LEVELS IN PSYCHOTIC DISORDERS. <i>Schizophrenia Bulletin</i> , 2019, 45, S206-S206.	4.3	3