

Adriana Maria Calvo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3772589/publications.pdf>

Version: 2024-02-01

28
papers

573
citations

516561

16
h-index

610775

24
g-index

32
all docs

32
docs citations

32
times ranked

519
citing authors

#	ARTICLE	IF	CITATIONS
1	Epinephrine Concentration (1:100,000 or 1:200,000) Does Not Affect the Clinical Efficacy of 4% Articaine for Lower Third Molar Removal: A Double-Blind, Randomized, Crossover Study. <i>Journal of Oral and Maxillofacial Surgery</i> , 2007, 65, 2445-2452.	0.5	59
2	Articaine and mepivacaine efficacy in postoperative analgesia for lower third molar removal: a double-blind, randomized, crossover study. <i>Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontics</i> , 2006, 102, 169-174.	1.6	58
3	A comparison of the clinical anesthetic efficacy of 4% articaine and 0.5% bupivacaine (both with) Tj ETQq1 1 0.784314 rgBT /Overlock Oral Radiology and Endodontics, 2008, 106, 19-28.	1.6	55
4	Timing of Alveolar Bone Grafting Determines Different Outcomes in Patients With Unilateral Cleft Palate. <i>Journal of Craniofacial Surgery</i> , 2012, 23, 1283-1286.	0.3	49
5	Analgesic and anti-inflammatory dose-response relationship of 7.5 and 15mg meloxicam after lower third molar removal: a double-blind, randomized, crossover study. <i>International Journal of Oral and Maxillofacial Surgery</i> , 2007, 36, 26-31.	0.7	35
6	Are antibiotics necessary after lower third molar removal?. <i>Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology</i> , 2012, 114, S199-S208.	0.2	32
7	Efficacy and Safety of 2% and 4% Articaine for Lower Third Molar Surgery. <i>Journal of Dental Research</i> , 2015, 94, 166S-173S.	2.5	30
8	The selective and non-selective cyclooxygenase inhibitors valdecoxib and piroxicam induce the same postoperative analgesia and control of trismus and swelling after lower third molar removal. <i>Brazilian Journal of Medical and Biological Research</i> , 2007, 40, 1133-1140.	0.7	29
9	Increase in Age Is Associated With Worse Outcomes in Alveolar Bone Grafting in Patients With Bilateral Complete Cleft Palate. <i>Journal of Craniofacial Surgery</i> , 2014, 25, 380-382.	0.3	24
10	Quantification of piroxicam and 5-hydroxy-piroxicam in human plasma and saliva using liquid chromatography-tandem mass spectrometry following oral administration. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2016, 120, 212-220.	1.4	21
11	Efficacy of piroxicam for postoperative pain after lower third molar surgery associated with CYP2C8 and CYP2C9. <i>Journal of Pain Research</i> , 2017, Volume 10, 1581-1589.	0.8	19
12	Comparison of oral versus sublingual piroxicam during postoperative pain management after lower third molar extraction. <i>International Journal of Oral and Maxillofacial Surgery</i> , 2011, 40, 292-297.	0.7	18
13	Sublingual ketorolac and sublingual piroxicam are equally effective for postoperative pain, trismus, and swelling management in lower third molar removal. <i>Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology</i> , 2012, 114, 27-34.	0.2	17
14	Ideal Versus Late Secondary Alveolar Bone Graft Surgery. <i>Cleft Palate-Craniofacial Journal</i> , 2018, 55, 369-374.	0.5	17
15	Losartan and isoproterenol promote alterations in the local renin-angiotensin system of rat salivary glands. <i>PLoS ONE</i> , 2019, 14, e0217030.	1.1	17
16	Efficacy of oral diclofenac with or without codeine for pain control after invasive bilateral third molar extractions. <i>International Journal of Oral and Maxillofacial Surgery</i> , 2017, 46, 621-627.	0.7	16
17	Effective method for the detection of piroxicam in human plasma using HPLC. <i>Brazilian Oral Research</i> , 2016, 30, .	0.6	12
18	Efficacy of Ketoprofen With or Without Omeprazole for Pain And Inflammation Control After Third Molar Removal. <i>Brazilian Dental Journal</i> , 2018, 29, 140-149.	0.5	10

#	ARTICLE	IF	CITATIONS
19	Simultaneous separation of naproxen and 6-O-desmethylnaproxen metabolite in saliva samples by liquid chromatography-tandem mass spectrometry: Pharmacokinetic study of naproxen alone and associated with esomeprazole. PLoS ONE, 2020, 15, e0236297.	1.1	10
20	The changing pattern of analgesic and anti-inflammatory drug use in cleft lip and palate repair. Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontics, 2006, 102, e16-e20.	1.6	9
21	Evolution of Postoperative Edema in Alveolar Graft Performed with Bone Morphogenetic Protein (rhBMP-2). Cleft Palate-Craniofacial Journal, 2015, 52, 168-175.	0.5	8
22	Efficacy of naproxen with or without esomeprazole for pain and inflammation in patients after bilateral third molar extractions: A double blinded crossover study. Medicina Oral, Patologia Oral Y Cirugia Bucal, 2016, 22, 0-0.	0.7	8
23	CYP450 polymorphisms and clinical pharmacogenetics of ibuprofen after lower third molar extraction. European Journal of Clinical Pharmacology, 2021, 77, 697-707.	0.8	6
24	Pharmacogenetic and Pharmacokinetic Assays from Saliva Samples Can Guarantee Personalized Drug Prescription. Brazilian Dental Journal, 2021, 32, 3-8.	0.5	6
25	Dental Pulp Fibroblasts Response after Stimulation with HEMA and Adhesive System. Brazilian Dental Journal, 2018, 29, 419-426.	0.5	3
26	Multifocal Analysis of Acute Pain After Third Molar Removal. Frontiers in Pharmacology, 2021, 12, 643874.	1.6	3
27	Molecular Response of Pulp Fibroblasts after Stimulation with Pulp Capping Materials. Brazilian Dental Journal, 2020, 31, 244-251.	0.5	2
28	Impacto do Tabagismo Passivo na Resposta PressÃ3rica Ã Epinefrina e Felipressina em Ratos Hipertensos 1K1C Tratados ou nÃ£o com Atenolol. Arquivos Brasileiros De Cardiologia, 2019, 114, 295-303.	0.3	0