

# Adriana Maria Calvo

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/3772589/adriana-maria-calvo-publications-by-year.pdf>

**Version:** 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

32  
papers

440  
citations

12  
h-index

20  
g-index

32  
ext. papers

510  
ext. citations

2.3  
avg, IF

2.77  
L-index

#	Paper	IF	Citations
32	Multifocal Analysis of Acute Pain After Third Molar Removal. <i>Frontiers in Pharmacology</i> , <b>2021</b> , 12, 643874	5.6	0
31	CYP450 polymorphisms and clinical pharmacogenetics of ibuprofen after lower third molar extraction. <i>European Journal of Clinical Pharmacology</i> , <b>2021</b> , 77, 697-707	2.8	3
30	Pharmacogenetic and Pharmacokinetic Assays from Saliva Samples Can Guarantee Personalized Drug Prescription. <i>Brazilian Dental Journal</i> , <b>2021</b> , 32, 3-8	1.9	0
29	Passive Cigarette Smoking Impact on Blood Pressure Response to Epinephrine and Felypressin in 1K1C Hypertensive Rats Treated or not with Atenolol. <i>Arquivos Brasileiros De Cardiologia</i> , <b>2020</b> , 114, 295-303	1.2	
28	Molecular Response of Pulp Fibroblasts after Stimulation with Pulp Capping Materials. <i>Brazilian Dental Journal</i> , <b>2020</b> , 31, 244-251	1.9	0
27	Simultaneous separation of naproxen and 6-O-desmethylnaproxen metabolite in saliva samples by liquid chromatography-tandem mass spectrometry: Pharmacokinetic study of naproxen alone and associated with esomeprazole. <i>PLoS ONE</i> , <b>2020</b> , 15, e0236297	3.7	1
26	Simultaneous separation of naproxen and 6-O-desmethylnaproxen metabolite in saliva samples by liquid chromatography-tandem mass spectrometry: Pharmacokinetic study of naproxen alone and associated with esomeprazole <b>2020</b> , 15, e0236297		
25	Simultaneous separation of naproxen and 6-O-desmethylnaproxen metabolite in saliva samples by liquid chromatography-tandem mass spectrometry: Pharmacokinetic study of naproxen alone and associated with esomeprazole <b>2020</b> , 15, e0236297		
24	Simultaneous separation of naproxen and 6-O-desmethylnaproxen metabolite in saliva samples by liquid chromatography-tandem mass spectrometry: Pharmacokinetic study of naproxen alone and associated with esomeprazole <b>2020</b> , 15, e0236297		
23	Simultaneous separation of naproxen and 6-O-desmethylnaproxen metabolite in saliva samples by liquid chromatography-tandem mass spectrometry: Pharmacokinetic study of naproxen alone and associated with esomeprazole <b>2020</b> , 15, e0236297		
22	Losartan and isoproterenol promote alterations in the local renin-angiotensin system of rat salivary glands. <i>PLoS ONE</i> , <b>2019</b> , 14, e0217030	3.7	10
21	Ideal Versus Late Secondary Alveolar Bone Graft Surgery: A Bone-Thickness Cone-Beam Computed Tomographic Assessment. <i>Cleft Palate-Craniofacial Journal</i> , <b>2018</b> , 55, 369-374	1.9	12
20	Efficacy of Ketoprofen With or Without Omeprazole for Pain And Inflammation Control After Third Molar Removal. <i>Brazilian Dental Journal</i> , <b>2018</b> , 29, 140-149	1.9	8
19	Dental Pulp Fibroblasts Response after Stimulation with HEMA and Adhesive System. <i>Brazilian Dental Journal</i> , <b>2018</b> , 29, 419-426	1.9	2
18	Efficacy of oral diclofenac with or without codeine for pain control after invasive bilateral third molar extractions. <i>International Journal of Oral and Maxillofacial Surgery</i> , <b>2017</b> , 46, 621-627	2.9	12
17	Efficacy of piroxicam for postoperative pain after lower third molar surgery associated with and. <i>Journal of Pain Research</i> , <b>2017</b> , 10, 1581-1589	2.9	16
16	Efficacy of naproxen with or without esomeprazole for pain and inflammation in patients after bilateral third molar extractions: A double blinded crossover study. <i>Medicina Oral, Patologia Oral Y Cirugia Bucal</i> , <b>2017</b> , 22, e122-e131	2.6	6

15	Quantification of piroxicam and 5-Hydroxy-piroxicam in human plasma and saliva using liquid chromatography-tandem mass spectrometry following oral administration. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2016</b> , 120, 212-20	3.5	10
14	Effective method for the detection of piroxicam in human plasma using HPLC. <i>Brazilian Oral Research</i> , <b>2016</b> , 30,	2.6	5
13	Efficacy and Safety of 2% and 4% Articaine for Lower Third Molar Surgery. <i>Journal of Dental Research</i> , <b>2015</b> , 94, 166S-73S	8.1	27
12	Evolution of Postoperative Edema in Alveolar Graft Performed With Bone Morphogenetic Protein (rhBMP-2). <i>Cleft Palate-Craniofacial Journal</i> , <b>2015</b> , 52, e168-75	1.9	6
11	Increase in age is associated with worse outcomes in alveolar bone grafting in patients with bilateral complete cleft palate. <i>Journal of Craniofacial Surgery</i> , <b>2014</b> , 25, 380-2	1.2	17
10	Sublingual ketorolac and sublingual piroxicam are equally effective for postoperative pain, trismus, and swelling management in lower third molar removal. <i>Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology</i> , <b>2012</b> , 114, 27-34	2	15
9	Are antibiotics necessary after lower third molar removal?. <i>Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology</i> , <b>2012</b> , 114, S199-208	2	28
8	Timing of alveolar bone grafting determines different outcomes in patients with unilateral cleft palate. <i>Journal of Craniofacial Surgery</i> , <b>2012</b> , 23, 1283-6	1.2	37
7	Comparison of oral versus sublingual piroxicam during postoperative pain management after lower third molar extraction. <i>International Journal of Oral and Maxillofacial Surgery</i> , <b>2011</b> , 40, 292-7	2.9	15
6	A comparison of the clinical anesthetic efficacy of 4% articaine and 0.5% bupivacaine (both with 1:200,000 epinephrine) for lower third molar removal. <i>Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontics</i> , <b>2008</b> , 106, 19-28		50
5	Epinephrine concentration (1:100,000 or 1:200,000) does not affect the clinical efficacy of 4% articaine for lower third molar removal: a double-blind, randomized, crossover study. <i>Journal of Oral and Maxillofacial Surgery</i> , <b>2007</b> , 65, 2445-52	1.8	55
4	The selective and non-selective cyclooxygenase inhibitors valdecoxib and piroxicam induce the same postoperative analgesia and control of trismus and swelling after lower third molar removal. <i>Brazilian Journal of Medical and Biological Research</i> , <b>2007</b> , 40, 1133-40	2.8	19
3	Analgesic and anti-inflammatory dose-response relationship of 7.5 and 15 mg meloxicam after lower third molar removal: a double-blind, randomized, crossover study. <i>International Journal of Oral and Maxillofacial Surgery</i> , <b>2007</b> , 36, 26-31	2.9	29
2	Articaine and mepivacaine efficacy in postoperative analgesia for lower third molar removal: a double-blind, randomized, crossover study. <i>Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontics</i> , <b>2006</b> , 102, 169-74		50
1	The changing pattern of analgesic and anti-inflammatory drug use in cleft lip and palate repair. <i>Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontics</i> , <b>2006</b> , 102, e16-20		7