

Monica Garcia-Alvarez

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3772597/publications.pdf>

Version: 2024-02-01

44
papers

697
citations

623188

14
h-index

610482

24
g-index

46
all docs

46
docs citations

46
times ranked

1454
citing authors

#	ARTICLE	IF	CITATIONS
1	Meta-analysis: implications of interleukin-28B polymorphisms in spontaneous and treatment-related clearance for patients with hepatitis C. <i>BMC Medicine</i> , 2013, 11, 6.	2.3	80
2	Relationship of vitamin D status with advanced liver fibrosis and response to hepatitis C virus therapy: A meta-analysis. <i>Hepatology</i> , 2014, 60, 1541-1550.	3.6	68
3	Nosocomial Spread of a <i>Staphylococcus hominis</i> subsp. <i>novobiosepticus</i> Strain Causing Sepsis in a Neonatal Intensive Care Unit. <i>Journal of Clinical Microbiology</i> , 2005, 43, 4877-4879.	1.8	57
4	Resistance of high fitness hepatitis C virus to lethal mutagenesis. <i>Virology</i> , 2018, 523, 100-109.	1.1	30
5	Vitamin D deficiency is associated with severity of liver disease in HIV/HCV coinfecting patients. <i>Journal of Infection</i> , 2014, 68, 176-184.	1.7	28
6	European Mitochondrial DNA Haplogroups and Metabolic Disorders in HIV/HCV-Coinfected Patients on Highly Active Antiretroviral Therapy. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2011, 58, 371-378.	0.9	22
7	Mx1, OAS1 and OAS2 polymorphisms are associated with the severity of liver disease in HIV/HCV-coinfecting patients: A cross-sectional study. <i>Scientific Reports</i> , 2017, 7, 41516.	1.6	22
8	Mitochondrial Haplogroups Are Associated With Clinical Pattern of AIDS Progression in HIV-Infected Patients. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2013, 63, 178-183.	0.9	21
9	Soluble markers of inflammation are associated with Framingham scores in HIV-infected patients on suppressive antiretroviral therapy. <i>Journal of Infection</i> , 2011, 63, 382-390.	1.7	19
10	Relationship between ITPA polymorphisms and hemolytic anemia in HCV-infected patients after ribavirin-based therapy: a meta-analysis. <i>Journal of Translational Medicine</i> , 2015, 13, 320.	1.8	19
11	Surveillance of transmitted drug resistance to integrase inhibitors in Spain: implications for clinical practice. <i>Journal of Antimicrobial Chemotherapy</i> , 2019, 74, 1693-1700.	1.3	19
12	European mitochondrial haplogroups are associated with CD4+ T cell recovery in HIV-infected patients on combination antiretroviral therapy. <i>Journal of Antimicrobial Chemotherapy</i> , 2013, 68, 2349-2357.	1.3	17
13	Mitochondrial DNA haplogroups are associated with severe sepsis and mortality in patients who underwent major surgery. <i>Journal of Infection</i> , 2015, 70, 20-29.	1.7	17
14	NS3 Resistance-Associated Variants (RAVs) in Patients Infected with HCV Genotype 1a in Spain. <i>PLoS ONE</i> , 2016, 11, e0163197.	1.1	16
15	HLA-E variants are associated with sustained virological response in HIV/hepatitis C virus-coinfecting patients on hepatitis C virus therapy. <i>Aids</i> , 2013, 27, 1231-1238.	1.0	15
16	High plasma fractalkine (CX3CL1) levels are associated with severe liver disease in HIV/HCV co-infected patients with HCV genotype 1. <i>Cytokine</i> , 2011, 54, 244-248.	1.4	14
17	IL28B polymorphisms are associated with severity of liver disease in human immunodeficiency virus (HIV) patients coinfecting with hepatitis C virus. <i>Journal of Infection</i> , 2013, 66, 170-178.	1.7	13
18	CXCL9, CXCL10 and CXCL11 polymorphisms are associated with sustained virologic response in HIV/HCV-coinfecting patients. <i>Journal of Clinical Virology</i> , 2014, 61, 423-429.	1.6	13

#	ARTICLE	IF	CITATIONS
19	European mitochondrial DNA haplogroups and liver fibrosis in HIV and hepatitis C virus coinfected patients. <i>Aids</i> , 2011, 25, 1619-1926.	1.0	12
20	Analysis of IL28B alleles with virologic response patterns and plasma cytokine levels in HIV/HCV-coinfected patients. <i>Aids</i> , 2013, 27, 163-173.	1.0	12
21	Association of adiponectin (<i>ADIPOQ</i>) rs2241766 polymorphism and dyslipidemia in HIV/HCV-coinfected patients. <i>European Journal of Clinical Investigation</i> , 2014, 44, 453-462.	1.7	12
22	<i>IL7RA</i> polymorphisms predict the CD4+ recovery in HIV patients on cART. <i>European Journal of Clinical Investigation</i> , 2015, 45, 1192-1199.	1.7	12
23	Impact of patatin-like phospholipase domain-containing 3 gene polymorphism (rs738409) on severity of liver disease in HIV/hepatitis C virus-coinfected patients. <i>Aids</i> , 2016, 30, 465-470.	1.0	12
24	The algorithm used for the interpretation of doravirine transmitted drug resistance strongly influences clinical practice and guideline recommendations. <i>Journal of Antimicrobial Chemotherapy</i> , 2020, 75, 1294-1300.	1.3	12
25	Bacterial DNA Translocation and Liver Disease Severity Among HIV-Infected Patients With Chronic Hepatitis C. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2012, 61, 552-556.	0.9	11
26	PPAR γ 2 Pro12Ala polymorphism was associated with favorable cardiometabolic risk profile in HIV/HCV coinfected patients: a cross-sectional study. <i>Journal of Translational Medicine</i> , 2014, 12, 235.	1.8	11
27	Single Nucleotide Polymorphisms of CXCL9-11 Chemokines Are Associated With Liver Fibrosis in HIV/HCV-Coinfected Patients. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2015, 68, 386-395.	0.9	11
28	First evidence of a pro-inflammatory response to severe infection with influenza virus H1N1. <i>Critical Care</i> , 2010, 14, 115.	2.5	10
29	Association between IL7R polymorphisms and severe liver disease in HIV/HCV coinfected patients: a cross-sectional study. <i>Journal of Translational Medicine</i> , 2015, 13, 206.	1.8	10
30	Analytical performance of four molecular platforms used for HIV-1, HBV and HCV viral load determinations. <i>Expert Review of Molecular Diagnostics</i> , 2019, 19, 941-949.	1.5	10
31	SLC30A8 rs13266634 polymorphism is related to a favorable cardiometabolic lipid profile in HIV/hepatitis C virus-coinfected patients. <i>Aids</i> , 2014, 28, 1325-1332.	1.0	9
32	ACSM4 Polymorphisms Are Associated With Rapid AIDS Progression in HIV-Infected Patients. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2014, 65, 27-32.	0.9	8
33	Optimal vitamin D plasma levels are associated with lower bacterial DNA translocation in HIV/hepatitis c virus coinfected patients. <i>Aids</i> , 2016, 30, 1069-1074.	1.0	7
34	TLR3 polymorphisms are associated with virologic response to hepatitis C virus (HCV) treatment in HIV/HCV coinfected patients. <i>Journal of Clinical Virology</i> , 2015, 65, 62-67.	1.6	6
35	Toll-like receptor 8 (TLR8) polymorphisms are associated with non-progression of chronic hepatitis C in HIV/HCV coinfected patients. <i>Infection, Genetics and Evolution</i> , 2015, 36, 339-344.	1.0	6
36	PPAR γ 2 Pro12Ala Polymorphism Is Associated With Sustained Virological Response in HIV/HCV-Coinfected Patients Under HCV Therapy. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2014, 67, 113-119.	0.9	5

#	ARTICLE	IF	CITATIONS
37	rs7903146 Polymorphism at Transcription Factor 7 Like 2 Gene Is Associated with Total Cholesterol and Lipoprotein Profile in HIV/Hepatitis C Virus-Coinfected Patients. <i>AIDS Research and Human Retroviruses</i> , 2015, 31, 326-334.	0.5	5
38	IL15 polymorphism is associated with advanced fibrosis, inflammation-related biomarkers and virological response in human immunodeficiency virus/hepatitis C virus coinfection. <i>Liver International</i> , 2016, 36, 1258-1266.	1.9	5
39	FTOrs9939609 polymorphism is associated with metabolic disturbances and response to HCV therapy in HIV/HCV-coinfected patients. <i>BMC Medicine</i> , 2014, 12, 198.	2.3	4
40	Serum levels of adipokines in HIV/HCV co-infected patients and their association with insulin resistance and liver disease severity. <i>Journal of Infection</i> , 2010, 61, 499-501.	1.7	3
41	Relationship between European Mitochondrial Haplogroups and Chronic Renal Allograft Rejection in Patients with Kidney Transplant. <i>International Journal of Medical Sciences</i> , 2014, 11, 1129-1132.	1.1	3
42	Sustained Virologic Response Decreases Serum Markers of Angiogenesis, Inflammation, and Fibrosis in HIV/HCV-Coinfected Patients on Hepatitis C Virus Therapy. <i>AIDS Patient Care and STDs</i> , 2011, 25, 131-133.	1.1	2
43	Reply. <i>Hepatology</i> , 2015, 62, 1643-1643.	3.6	2
44	Short Communication: CXCL12 rs1029153 Polymorphism Is Associated with the Sustained Virological Response in HIV/Hepatitis C Virus-Coinfected Patients on Hepatitis C Virus Therapy. <i>AIDS Research and Human Retroviruses</i> , 2016, 32, 226-231.	0.5	0