

# Knut Madslie

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3777646/publications.pdf>

Version: 2024-02-01

28  
papers

595  
citations

567281

15  
h-index

610901

24  
g-index

28  
all docs

28  
docs citations

28  
times ranked

661  
citing authors

#	ARTICLE	IF	CITATIONS
1	Capture, Anesthesia, and Disturbance of Free-Ranging Brown Bears ( <i>Ursus arctos</i> ) during Hibernation. PLoS ONE, 2012, 7, e40520.	2.5	69
2	Studies in bank voles reveal strain differences between chronic wasting disease prions from Norway and North America. Proceedings of the National Academy of Sciences of the United States of America, 2020, 117, 31417-31426.	7.1	57
3	Fennoscandian distribution of an important parasite of cervids, the deer ked ( <i>Lipoptena cervi</i> ), revisited. Parasitology Research, 2010, 107, 117-125.	1.6	42
4	Bartonella Infections in Deer Keds ( <i>Lipoptena cervi</i> ) and Moose ( <i>Alces alces</i> ) in Norway. Applied and Environmental Microbiology, 2013, 79, 322-327.	3.1	41
5	HAIR-LOSS EPIZOOTIC IN MOOSE ( <i>ALCES ALCES</i> ) ASSOCIATED WITH MASSIVE DEER KED ( <i>LIPOPTENA CERVI</i> ) INFESTATION. Journal of Wildlife Diseases, 2011, 47, 893-906.	0.8	38
6	What does the fox say? Monitoring antimicrobial resistance in the environment using wild red foxes as an indicator. PLoS ONE, 2018, 13, e0198019.	2.5	30
7	Adaptive selection of a prion strain conformer corresponding to established North American CWD during propagation of novel emergent Norwegian strains in mice expressing elk or deer prion protein. PLoS Pathogens, 2021, 17, e1009748.	4.7	30
8	Chronic wasting disease associated with prion protein gene ( <i>PRNP</i> ) variation in Norwegian wild reindeer ( <i>Rangifer tarandus</i> ). Prion, 2020, 14, 1-10.	1.8	28
9	<i>Echinococcus multilocularis</i> adaptation of a worm egg isolation procedure coupled with a multiplex PCR assay to carry out large-scale screening of red foxes ( <i>Vulpes vulpes</i> ) in Norway. Parasitology Research, 2009, 104, 509-514.	1.6	26
10	Geographical variation in host use of a blood-feeding ectoparasitic fly: implications for population invasiveness. Oecologia, 2011, 166, 985-995.	2.0	25
11	The demographic pattern of infection with chronic wasting disease in reindeer at an early epidemic stage. Ecosphere, 2019, 10, e02931.	2.2	25
12	Humanized Transgenic Mice Are Resistant to Chronic Wasting Disease Prions From Norwegian Reindeer and Moose. Journal of Infectious Diseases, 2022, 226, 933-937.	4.0	25
13	Factors affecting deer ked ( <i>Lipoptena cervi</i> ) prevalence and infestation intensity in moose ( <i>Alces alces</i> ) in Norway. Parasites and Vectors, 2012, 5, 251.	2.5	22
14	Cervids as sentinel species for tick-borne encephalitis virus in Norway – A serological study. Zoonoses and Public Health, 2020, 67, 342-351.	2.2	19
15	Gastrointestinal parasites in an isolated Norwegian population of wild red deer ( <i>Cervus elaphus</i> ). Acta Veterinaria Scandinavica, 2014, 56, 59.	1.6	18
16	Phenology of deer ked ( <i>Lipoptena cervi</i> ) host-seeking flight activity and its relationship with prevailing autumn weather. Parasites and Vectors, 2016, 9, 95.	2.5	14
17	Seroprevalence of Hepatitis E Virus in Moose ( <i>Alces alces</i> ), Reindeer ( <i>Rangifer tarandus</i> ), Red Deer ( <i>Cervus elaphus</i> ), Roe Deer ( <i>Capreolus capreolus</i> ), and Muskoxen ( <i>Ovibos moschatus</i> ) from Norway. Viruses, 2021, 13, 224.	3.3	14
18	Chronic wasting disease in Norway – A survey of prion protein gene variation among cervids. Transboundary and Emerging Diseases, 2022, 69, .	3.0	13

#	ARTICLE	IF	CITATIONS
19	Gammaherpesvirus in Cervid Species from Norway: Characterization of a New Virus in Wild and Semi-Domesticated Eurasian Tundra Reindeer ( <i>Rangifer tarandus tarandus</i> ). <i>Viruses</i> , 2020, 12, 876.	3.3	10
20	Comparison of anticoagulant rodenticide concentrations in liver and feces from apparently healthy red foxes. <i>Journal of Veterinary Diagnostic Investigation</i> , 2020, 32, 560-564.	1.1	10
21	<i>Bartonella</i> spp. detection in ticks, <i>Culicoides</i> biting midges and wild cervids from Norway. <i>Transboundary and Emerging Diseases</i> , 2021, 68, 941-951.	3.0	9
22	Long-Term Safety of Intraperitoneal Radio Transmitter Implants in Brown Bears ( <i>Ursus arctos</i> ). <i>Frontiers in Veterinary Science</i> , 2018, 5, 252.	2.2	7
23	Hair Cortisol Concentration and Body Mass in Moose ( <i>Alces alces</i> ) Infested with Deer Keds ( <i>Lipoptena</i> ) Tj ETQq1 1 0.784314,rgBT /Over 0.8	0.8	0
24	First detection of highly pathogenic avian influenza virus in Norway. <i>BMC Veterinary Research</i> , 2021, 17, 218.	1.9	6
25	High winter loads of Oestrid larvae and <i>Elaphostrongylus rangiferi</i> are associated with emaciation in wild reindeer calves. <i>International Journal for Parasitology: Parasites and Wildlife</i> , 2021, 15, 214-224.	1.5	3
26	Yew ( <i>Taxus</i> ) intoxication in free-ranging cervids. <i>PLoS ONE</i> , 2017, 12, e0188961.	2.5	3
27	Distribution, prevalence and intensity of moose nose bot fly ( <i>Cephenemyia ulrichii</i> ) larvae in moose ( <i>Alces alces</i> ) from Norway. <i>International Journal for Parasitology: Parasites and Wildlife</i> , 2021, 15, 120-126.	1.5	2
28	PREVALENCE OF ANTICOAGULANT RODENTICIDES IN FECES OF WILD RED FOXES ( ) IN NORWAY. <i>Journal of Wildlife Diseases</i> , 2019, 55, 834-843.	0.8	2