

David M Wellsted

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3782153/david-m-wellsted-publications-by-year.pdf>
Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

71 papers	2,107 citations	27 h-index	44 g-index
74 ext. papers	2,429 ext. citations	4.8 avg, IF	4.69 L-index

#	Paper	IF	Citations
71	Evaluation of a whole system approach to diet and healthy weight in the east of Scotland: Study protocol.. <i>PLoS ONE</i> , 2022 , 17, e0265667	3.7	0
70	Commentary on the article: "Maintenance of wellness in patients with obsessive-compulsive disorder who discontinue medication after exposure/response prevention augmentation A randomized clinical trial" Foa EB et al., <i>JAMA Psychiatry</i> . 2022;79(3):193-200 (1).. <i>Comprehensive Psychiatry</i> , 2022 , 116, 152323	7.3	
69	Contributions of treatment centre and patient characteristics to patient-reported experience of haemodialysis: a national cross-sectional study. <i>BMJ Open</i> , 2021 , 11, e044984	3	0
68	A multicenter feasibility randomized controlled trial to assess the impact of incremental versus conventional initiation of hemodialysis on residual kidney function. <i>Kidney International</i> , 2021 ,	9.9	7
67	Facing the "new normal": How adjusting to the easing of COVID-19 lockdown restrictions exposes mental health inequalities. <i>Journal of Psychiatric Research</i> , 2021 , 141, 276-286	5.2	2
66	Antidepressant Usage in Haemodialysis Patients: Evidence of Sub-Optimal Practice Patterns. <i>Journal of Renal Care</i> , 2020 , 46, 124-132	1.6	4
65	Rationale and design of "Can Very Low Dose Rivaroxaban (VLDR) in addition to dual antiplatelet therapy improve thrombotic status in acute coronary syndrome (VaLiDate-R)" study : A randomised trial modulating endogenous fibrinolysis in patients with acute coronary syndrome. <i>Journal of Thrombosis and Thrombolysis</i> , 2020 , 49, 192-198	5.1	2
64	Impact of incremental versus conventional initiation of haemodialysis on residual kidney function: study protocol for a multicentre feasibility randomised controlled trial. <i>BMJ Open</i> , 2020 , 10, e035919	3	6
63	Apixaban enhances endogenous fibrinolysis in patients with atrial fibrillation. <i>Europace</i> , 2019 , 21, 1297-1306	3.9	9
62	Comparison of characteristics of centers practicing incremental vs. conventional approaches to hemodialysis delivery - postdialysis recovery time and patient survival. <i>Hemodialysis International</i> , 2019 , 23, 288-296	1.7	6
61	The emergence of Exercise Addiction, Body Dysmorphic Disorder, and other image-related psychopathological correlates in fitness settings: A cross sectional study. <i>PLoS ONE</i> , 2019 , 14, e0213060	3.7	43
60	How do patients from South Asian backgrounds experience life on haemodialysis in the UK? A multicentre qualitative study. <i>BMJ Open</i> , 2019 , 9, e024739	3	5
59	Initiating haemodialysis twice-weekly as part of an incremental programme may protect residual kidney function. <i>Nephrology Dialysis Transplantation</i> , 2019 , 34, 1017-1025	4.3	12
58	Impaired endogenous fibrinolysis in ST-segment elevation myocardial infarction patients undergoing primary percutaneous coronary intervention is a predictor of recurrent cardiovascular events: the RISK PPCI study. <i>European Heart Journal</i> , 2019 , 40, 295-305	9.5	32
57	Effect of P2Y inhibitors on thrombus stability and endogenous fibrinolysis. <i>Thrombosis Research</i> , 2019 , 173, 102-108	8.2	8
56	Postdialysis recovery time is extended in patients with greater self-reported depression screening questionnaire scores. <i>Hemodialysis International</i> , 2018 , 22, 369-376	1.7	10
55	Morphine Analgesia Pre-PPCI Is Associated with Prothrombotic State, Reduced Spontaneous Reperfusion and Greater Infarct Size. <i>Thrombosis and Haemostasis</i> , 2018 , 118, 601-612	7	28

54	Self-reported depression symptoms in haemodialysis patients: Bi-factor structures of two common measures and their association with clinical factors. <i>General Hospital Psychiatry</i> , 2018 , 54, 31-36	5.6	6
53	Depression Symptoms in Haemodialysis Patients Predict All-Cause Mortality but Not Kidney Transplantation: A Cause-Specific Outcome Analysis. <i>Annals of Behavioral Medicine</i> , 2018 , 52, 1-8	4.5	25
52	Sertraline Versus Placebo in Patients with Major Depressive Disorder Undergoing Hemodialysis: A Randomized, Controlled Feasibility Trial. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2017 , 12, 280-286	6.9	42
51	Measuring Fatigue Using the Multidimensional Fatigue Inventory-20: A Questionable Factor Structure in Haemodialysis Patients. <i>Nephron</i> , 2017 , 136, 121-126	3.3	10
50	Percutaneous coronary intervention with drug-eluting stent versus coronary artery bypass grafting: A meta-analysis of patients with left main coronary artery disease. <i>International Journal of Cardiology</i> , 2017 , 249, 101-106	3.2	1
49	Impaired thrombolytic status predicts adverse cardiac events in patients undergoing primary percutaneous coronary intervention. <i>Thrombosis and Haemostasis</i> , 2017 , 117, 457-470	7	14
48	Feasibility study of a randomised controlled trial to investigate the effectiveness of using a humanoid robot to improve the social skills of children with autism spectrum disorder (Kaspar RCT): a study protocol. <i>BMJ Open</i> , 2017 , 7, e017376	3	23
47	C reactive protein and depressive symptoms in hemodialysis patients: A questionable association. <i>Hemodialysis International</i> , 2017 , 21, 542-548	1.7	3
46	Exploring Cognitive Concomitants of Mental Fatigue in Patients with Coronary Artery Disease. <i>Neuropsychobiology</i> , 2017 , 76, 151-160	4	7
45	Evidence of improved fluid management in patients receiving haemodialysis following a self-affirmation theory-based intervention: A randomised controlled trial. <i>Psychology and Health</i> , 2016 , 31, 100-14	2.9	24
44	Wordless intervention for people with epilepsy and learning disabilities (WIELD): a randomised controlled feasibility trial. <i>BMJ Open</i> , 2016 , 6, e012993	3	5
43	"Sometimes, it just stops me from doing anything": A qualitative exploration of epilepsy management in people with intellectual disabilities and their carers. <i>Epilepsy and Behavior</i> , 2016 , 64, 133-139	3.2	7
42	Predicting residual kidney function in hemodialysis patients using serum B ₂ -microglobulin and K ₂ -microglobulin. <i>Kidney International</i> , 2016 , 89, 1090-1098	9.9	39
41	Relative effects of different non-vitamin K antagonist oral anticoagulants on global thrombotic status in atrial fibrillation. <i>Platelets</i> , 2016 , 27, 687-693	3.6	12
40	A study of sertraline in dialysis (ASSertID): a protocol for a pilot randomised controlled trial of drug treatment for depression in patients undergoing haemodialysis. <i>BMC Nephrology</i> , 2015 , 16, 172	2.7	13
39	Medication beliefs are associated with phosphate binder non-adherence in hyperphosphatemic haemodialysis patients. <i>British Journal of Health Psychology</i> , 2015 , 20, 563-78	8.3	19
38	Simvastatin improves the sexual health-related quality of life in men aged 40 years and over with erectile dysfunction: additional data from the erectile dysfunction and statin trial. <i>BMC Urology</i> , 2014 , 14, 24	2.2	8
37	Disease-specific predictive formulas for energy expenditure in the dialysis population. <i>Journal of Renal Nutrition</i> , 2014 , 24, 243-51	3	23

36	Can simvastatin improve erectile function and health-related quality of life in men aged ≥40 years with erectile dysfunction? Results of the Erectile Dysfunction and Statins Trial [ISRCTN66772971]. <i>BJU International</i> , 2013 , 111, 324-33	5.6	24
35	Distinct depression symptom trajectories over the first year of dialysis: associations with illness perceptions. <i>Annals of Behavioral Medicine</i> , 2013 , 45, 78-88	4.5	34
34	The size, burden and cost of disorders of the brain in the UK. <i>Journal of Psychopharmacology</i> , 2013 , 27, 761-70	4.6	109
33	How robust is the Burpise questionPin predicting short-term mortality risk in haemodialysis patients?. <i>Nephron Clinical Practice</i> , 2013 , 123, 185-93		23
32	A systematic review of the literature on counselling and psychotherapy for the prevention of suicide: 1. Quantitative outcome and process studies. <i>Counselling and Psychotherapy Research</i> , 2013 , 13, 164-183	1.3	10
31	Perfusion CT assessment of the colon and rectum: feasibility of quantification of bowel wall perfusion and vascularization. <i>European Journal of Radiology</i> , 2012 , 81, 821-4	4.7	13
30	Quality of life and survival in patients with advanced kidney failure managed conservatively or by dialysis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2012 , 7, 2002-9	6.9	135
29	The factor structure of the Revised Illness Perception Questionnaire (IPQ-R) in end-stage renal disease patients. <i>Psychology, Health and Medicine</i> , 2012 , 17, 578-88	2.1	24
28	First direct comparison of platelet reactivity and thrombolytic status between Japanese and Western volunteers: possible relationship to the "Japanese paradox". <i>International Journal of Cardiology</i> , 2011 , 152, 43-8	3.2	40
27	A confirmatory factor analysis of the Beck Depression Inventory-II in end-stage renal disease patients. <i>Journal of Psychosomatic Research</i> , 2011 , 71, 148-53	4.1	38
26	Can simvastatin improve erectile function and health-related quality of life in men aged >40 years with erectile dysfunction? Rationale and design of the Erectile Dysfunction and Statins (EDS) Trial [ISRCTN66772971](1). <i>BJU International</i> , 2011 , 108, 1850-4	5.6	6
25	Commercial software upgrades may significantly alter Perfusion CT parameter values in colorectal cancer. <i>European Radiology</i> , 2011 , 21, 744-9	8	27
24	An association between depressive symptoms and survival in incident dialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2011 , 26, 1628-34	4.3	84
23	Illness representations and concurrent depression symptoms in haemodialysis patients. <i>Journal of Health Psychology</i> , 2011 , 16, 1127-37	3.1	29
22	Does antifungal prophylaxis with daily oral fluconazole reduce the risk of fungal peritonitis in peritoneal dialysis patients? The Pan Thames Renal Audit. <i>Blood Purification</i> , 2011 , 32, 181-5	3.1	10
21	Choosing not to take phosphate binders: the role of dialysis patientsPmedication beliefs. <i>Nephron Clinical Practice</i> , 2011 , 119, c205-13		27
20	Illness perceptions predict survival in haemodialysis patients. <i>American Journal of Nephrology</i> , 2011 , 33, 358-63	4.6	46
19	Identifying depression in South asian patients with end-stage renal disease: considerations for practice. <i>Nephron Extra</i> , 2011 , 1, 262-71		6

18	Quantitative helical dynamic contrast enhanced computed tomography assessment of the spatial variation in whole tumour blood volume with radiotherapy in lung cancer. <i>Lung Cancer</i> , 2010 , 69, 71-6	5.9	15
17	Illness representations are associated with fluid nonadherence among hemodialysis patients. <i>Journal of Psychosomatic Research</i> , 2010 , 68, 203-12	4.1	59
16	A pilot study of the epilepsy risk awareness checklist (ERAC) in people with epilepsy and learning disabilities. <i>Seizure: the Journal of the British Epilepsy Association</i> , 2010 , 19, 592-6	3.2	6
15	Magnetic resonance imaging assessment of squamous cell carcinoma of the anal canal before and after chemoradiation: can MRI predict for eventual clinical outcome?. <i>International Journal of Radiation Oncology Biology Physics</i> , 2010 , 78, 715-21	4	48
14	Depression in end-stage renal disease: current advances and research. <i>Seminars in Dialysis</i> , 2010 , 23, 74-82	5	42
13	A phase I trial of radioimmunotherapy with 131I-A5B7 anti-CEA antibody in combination with combretastatin-A4-phosphate in advanced gastrointestinal carcinomas. <i>Clinical Cancer Research</i> , 2009 , 15, 4484-92	12.9	66
12	An association between residual renal function and depression symptoms in haemodialysis patients. <i>Nephron Clinical Practice</i> , 2009 , 113, c117-24		12
11	Residual renal function improves outcome in incremental haemodialysis despite reduced dialysis dose. <i>Nephrology Dialysis Transplantation</i> , 2009 , 24, 2502-10	4.3	119
10	Long-term outcomes in online hemodiafiltration and high-flux hemodialysis: a comparative analysis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2009 , 4, 1944-53	6.9	100
9	Influence of N-butylscopolamine on SUV in FDG PET of the bowel. <i>Annals of Nuclear Medicine</i> , 2009 , 23, 471-8	2.5	2
8	Can perfusion CT assessment of primary colorectal adenocarcinoma blood flow at staging predict for subsequent metastatic disease? A pilot study. <i>European Radiology</i> , 2009 , 19, 79-89	8	67
7	Assessment of the spatial pattern of colorectal tumour perfusion estimated at perfusion CT using two-dimensional fractal analysis. <i>European Radiology</i> , 2009 , 19, 1358-65	8	49
6	Auditory memory for timbre. <i>Journal of Experimental Psychology: Human Perception and Performance</i> , 2009 , 35, 855-75	2.6	18
5	Depression on dialysis. <i>Nephron Clinical Practice</i> , 2008 , 108, c256-64		116
4	Quantitative assessment of colorectal cancer tumor vascular parameters by using perfusion CT: influence of tumor region of interest. <i>Radiology</i> , 2008 , 247, 726-32	20.5	76
3	Colorectal tumor vascularity: quantitative assessment with multidetector CT--do tumor perfusion measurements reflect angiogenesis?. <i>Radiology</i> , 2008 , 249, 510-7	20.5	113
2	Screening for depression while patients dialyse: an evaluation. <i>Nephrology Dialysis Transplantation</i> , 2008 , 23, 2653-9	4.3	59
1	Facing Loss and Finding Hope in Narrating Together: Accounts of Parenthood Following the Death of a Child to Muscular Dystrophy. <i>Illness Crisis and Loss</i> , 105413732110137	0.6	

