

David M Wellsted

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3782153/publications.pdf>

Version: 2024-02-01

72
papers

2,760
citations

159358

30
h-index

189595

50
g-index

74
all docs

74
docs citations

74
times ranked

3598
citing authors

#	ARTICLE	IF	CITATIONS
1	Quality of Life and Survival in Patients with Advanced Kidney Failure Managed Conservatively or by Dialysis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2012, 7, 2002-2009.	2.2	169
2	The size, burden and cost of disorders of the brain in the UK. <i>Journal of Psychopharmacology</i> , 2013, 27, 761-770.	2.0	163
3	Depression on Dialysis. <i>Nephron Clinical Practice</i> , 2008, 108, c256-c264.	2.3	148
4	Long-Term Outcomes in Online Hemodiafiltration and High-Flux Hemodialysis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2009, 4, 1944-1953.	2.2	142
5	Residual renal function improves outcome in incremental haemodialysis despite reduced dialysis dose. <i>Nephrology Dialysis Transplantation</i> , 2009, 24, 2502-2510.	0.4	140
6	Colorectal Tumor Vascularity: Quantitative Assessment with Multidetector CT—Do Tumor Perfusion Measurements Reflect Angiogenesis?. <i>Radiology</i> , 2008, 249, 510-517.	3.6	128
7	An association between depressive symptoms and survival in incident dialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2011, 26, 1628-1634.	0.4	98
8	Can perfusion CT assessment of primary colorectal adenocarcinoma blood flow at staging predict for subsequent metastatic disease? A pilot study. <i>European Radiology</i> , 2009, 19, 79-89.	2.3	82
9	Quantitative Assessment of Colorectal Cancer Tumor Vascular Parameters by Using Perfusion CT: Influence of Tumor Region of Interest. <i>Radiology</i> , 2008, 247, 726-732.	3.6	81
10	Illness representations are associated with fluid nonadherence among hemodialysis patients. <i>Journal of Psychosomatic Research</i> , 2010, 68, 203-212.	1.2	71
11	The emergence of Exercise Addiction, Body Dysmorphic Disorder, and other image-related psychopathological correlates in fitness settings: A cross sectional study. <i>PLoS ONE</i> , 2019, 14, e0213060.	1.1	71
12	A Phase I Trial of Radioimmunotherapy with 131I-A5B7 Anti-CEA Antibody in Combination with Combretastatin-A4-Phosphate in Advanced Gastrointestinal Carcinomas. <i>Clinical Cancer Research</i> , 2009, 15, 4484-4492.	3.2	68
13	Screening for depression while patients dialyse: an evaluation. <i>Nephrology Dialysis Transplantation</i> , 2008, 23, 2653-2659.	0.4	66
14	Magnetic Resonance Imaging Assessment of Squamous Cell Carcinoma of the Anal Canal Before and After Chemoradiation: Can MRI Predict for Eventual Clinical Outcome?. <i>International Journal of Radiation Oncology Biology Physics</i> , 2010, 78, 715-721.	0.4	62
15	Sertraline Versus Placebo in Patients with Major Depressive Disorder Undergoing Hemodialysis: A Randomized, Controlled Feasibility Trial. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2017, 12, 280-286.	2.2	62
16	Impaired endogenous fibrinolysis in ST-segment elevation myocardial infarction patients undergoing primary percutaneous coronary intervention is a predictor of recurrent cardiovascular events: the RISK PPCI study. <i>European Heart Journal</i> , 2019, 40, 295-305.	1.0	61
17	Assessment of the spatial pattern of colorectal tumour perfusion estimated at perfusion CT using two-dimensional fractal analysis. <i>European Radiology</i> , 2009, 19, 1358-1365.	2.3	59
18	Depression in End-Stage Renal Disease: Current Advances and Research. <i>Seminars in Dialysis</i> , 2010, 23, 74-82.	0.7	54

#	ARTICLE	IF	CITATIONS
19	Illness Perceptions Predict Survival in Haemodialysis Patients. American Journal of Nephrology, 2011, 33, 358-363.	1.4	51
20	Predicting residual kidney function in hemodialysis patients using serum β_2 -trace protein and β_2 -microglobulin. Kidney International, 2016, 89, 1090-1098.	2.6	51
21	First direct comparison of platelet reactivity and thrombolytic status between Japanese and Western volunteers: Possible relationship to the "Japanese paradox". International Journal of Cardiology, 2011, 152, 43-48.	0.8	47
22	Distinct Depression Symptom Trajectories over the First Year of Dialysis: Associations with Illness Perceptions. Annals of Behavioral Medicine, 2013, 45, 78-88.	1.7	42
23	A confirmatory factor analysis of the beck depression inventory-II in end-stage renal disease patients. Journal of Psychosomatic Research, 2011, 71, 148-153.	1.2	40
24	Depression Symptoms in Haemodialysis Patients Predict All-Cause Mortality but Not Kidney Transplantation: A Cause-Specific Outcome Analysis. Annals of Behavioral Medicine, 2018, 52, 1-8.	1.7	37
25	Choosing Not to Take Phosphate Binders: The Role of Dialysis Patients' Medication Beliefs. Nephron Clinical Practice, 2011, 119, 205-213.	2.3	34
26	Morphine Analgesia Pre-PPCI Is Associated with Prothrombotic State, Reduced Spontaneous Reperfusion and Greater Infarct Size. Thrombosis and Haemostasis, 2018, 118, 601-612.	1.8	34
27	Commercial software upgrades may significantly alter Perfusion CT parameter values in colorectal cancer. European Radiology, 2011, 21, 744-749.	2.3	33
28	Illness representations and concurrent depression symptoms in haemodialysis patients. Journal of Health Psychology, 2011, 16, 1127-1137.	1.3	33
29	Feasibility study of a randomised controlled trial to investigate the effectiveness of using a humanoid robot to improve the social skills of children with autism spectrum disorder (Kaspar RCT): a study protocol. BMJ Open, 2017, 7, e017376.	0.8	33
30	How Robust Is the "Surprise Question" in Predicting Short-Term Mortality Risk in Haemodialysis Patients. Nephron Clinical Practice, 2013, 123, 185-193.	2.3	32
31	A multicenter feasibility randomized controlled trial to assess the impact of incremental versus conventional initiation of hemodialysis on residual kidney function. Kidney International, 2022, 101, 615-625.	2.6	31
32	Medication beliefs are associated with phosphate binder non-adherence in hyperphosphatemic haemodialysis patients. British Journal of Health Psychology, 2015, 20, 563-578.	1.9	30
33	The factor structure of the revised illness perception questionnaire (IPQ-R) in end-stage renal disease patients. Psychology, Health and Medicine, 2012, 17, 578-588.	1.3	29
34	Disease-Specific Predictive Formulas for Energy Expenditure in the Dialysis Population. , 2014, 24, 243-251.		29
35	Evidence of improved fluid management in patients receiving haemodialysis following a self-affirmation theory-based intervention: A randomised controlled trial. Psychology and Health, 2016, 31, 100-114.	1.2	28
36	Can simvastatin improve erectile function and health-related quality of life in men aged ≥ 40 years with erectile dysfunction? Results of the Erectile Dysfunction and Statins Trial [scpi>ISRCTN</scpi>66772971]. BJU International, 2013, 111, 324-333.	1.3	26

#	ARTICLE	IF	CITATIONS
37	Postdialysis recovery time is extended in patients with greater self-reported depression screening questionnaire scores. <i>Hemodialysis International</i> , 2018, 22, 369-376.	0.4	23
38	Auditory memory for timbre.. <i>Journal of Experimental Psychology: Human Perception and Performance</i> , 2009, 35, 855-875.	0.7	21
39	Apixaban enhances endogenous fibrinolysis in patients with atrial fibrillation. <i>Europace</i> , 2019, 21, 1297-1306.	0.7	20
40	Facing the "new normal": How adjusting to the easing of COVID-19 lockdown restrictions exposes mental health inequalities. <i>Journal of Psychiatric Research</i> , 2021, 141, 276-286.	1.5	19
41	Perfusion CT assessment of the colon and rectum: Feasibility of quantification of bowel wall perfusion and vascularization. <i>European Journal of Radiology</i> , 2012, 81, 821-824.	1.2	18
42	A systematic review of the literature on counselling and psychotherapy for the prevention of suicide: 1. Quantitative outcome and process studies. <i>Counselling and Psychotherapy Research</i> , 2013, 13, 164-183.	1.7	17
43	Impaired thrombolytic status predicts adverse cardiac events in patients undergoing primary percutaneous coronary intervention. <i>Thrombosis and Haemostasis</i> , 2017, 117, 457-470.	1.8	17
44	Initiating haemodialysis twice-weekly as part of an incremental programme may protect residual kidney function. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, 1017-1025.	0.4	17
45	Quantitative helical dynamic contrast enhanced computed tomography assessment of the spatial variation in whole tumour blood volume with radiotherapy in lung cancer. <i>Lung Cancer</i> , 2010, 69, 71-76.	0.9	16
46	A study of sertraline in dialysis (ASSertID): a protocol for a pilot randomised controlled trial of drug treatment for depression in patients undergoing haemodialysis. <i>BMC Nephrology</i> , 2015, 16, 172.	0.8	16
47	Relative effects of different non-vitamin K antagonist oral anticoagulants on global thrombotic status in atrial fibrillation. <i>Platelets</i> , 2016, 27, 687-693.	1.1	16
48	Effect of P2Y12 inhibitors on thrombus stability and endogenous fibrinolysis. <i>Thrombosis Research</i> , 2019, 173, 102-108.	0.8	16
49	An Association between Residual Renal Function and Depression Symptoms in Haemodialysis Patients. <i>Nephron Clinical Practice</i> , 2009, 113, c117-c124.	2.3	15
50	Measuring Fatigue Using the Multidimensional Fatigue Inventory-20: A Questionable Factor Structure in Haemodialysis Patients. <i>Nephron</i> , 2017, 136, 121-126.	0.9	14
51	Does Antifungal Prophylaxis with Daily Oral Fluconazole Reduce the Risk of Fungal Peritonitis in Peritoneal Dialysis Patients? <i>The Pan Thames Renal Audit</i> . <i>Blood Purification</i> , 2011, 32, 181-185.	0.9	13
52	Comparison of characteristics of centers practicing incremental vs. conventional approaches to hemodialysis delivery " postdialysis recovery time and patient survival. <i>Hemodialysis International</i> , 2019, 23, 288-296.	0.4	12
53	A pilot study of the epilepsy risk awareness checklist (ERAC) in people with epilepsy and learning disabilities. <i>Seizure: the Journal of the British Epilepsy Association</i> , 2010, 19, 592-596.	0.9	11
54	Self-reported depression symptoms in haemodialysis patients: Bi-factor structures of two common measures and their association with clinical factors. <i>General Hospital Psychiatry</i> , 2018, 54, 31-36.	1.2	11

#	ARTICLE	IF	CITATIONS
55	Simvastatin improves the sexual health-related quality of life in men aged 40 years and over with erectile dysfunction: additional data from the erectile dysfunction and statin trial. <i>BMC Urology</i> , 2014, 14, 24.	0.6	10
56	“Sometimes, it just stops me from doing anything” A qualitative exploration of epilepsy management in people with intellectual disabilities and their carers. <i>Epilepsy and Behavior</i> , 2016, 64, 133-139.	0.9	10
57	How do patients from South Asian backgrounds experience life on haemodialysis in the UK? A multicentre qualitative study. <i>BMJ Open</i> , 2019, 9, e024739.	0.8	10
58	Antidepressant Usage in Haemodialysis Patients: Evidence of Suboptimal Practice Patterns. <i>Journal of Renal Care</i> , 2020, 46, 124-132.	0.6	10
59	Identifying Depression in South Asian Patients with End-Stage Renal Disease: Considerations for Practice. <i>Nephron Extra</i> , 2011, 1, 262-271.	1.1	8
60	Exploring Cognitive Concomitants of Mental Fatigue in Patients with Coronary Artery Disease. <i>Neuropsychobiology</i> , 2017, 76, 151-160.	0.9	8
61	Can simvastatin improve erectile function and health-related quality of life in men aged >40 years with erectile dysfunction? Rationale and design of the Erectile Dysfunction and Statins (EDS) Trial [ISRCTN66772971]. <i>BJU International</i> , 2011, 108, 1850-1854.	1.3	7
62	Wordless intervention for people with epilepsy and learning disabilities (WIELD): a randomised controlled feasibility trial. <i>BMJ Open</i> , 2016, 6, e012993.	0.8	7
63	Impact of incremental versus conventional initiation of haemodialysis on residual kidney function: study protocol for a multicentre feasibility randomised controlled trial. <i>BMJ Open</i> , 2020, 10, e035919.	0.8	7
64	Rationale and design of “Can Very Low Dose Rivaroxaban (VLDR) in addition to dual antiplatelet therapy improve thrombotic status in acute coronary syndrome (VaLiDate-R)” study. <i>Journal of Thrombosis and Thrombolysis</i> , 2020, 49, 192-198.	1.0	6
65	Evaluation of a whole system approach to diet and healthy weight in the east of Scotland: Study protocol. <i>PLoS ONE</i> , 2022, 17, e0265667.	1.1	6
66	C reactive protein and depressive symptoms in hemodialysis patients: A questionable association. <i>Hemodialysis International</i> , 2017, 21, 542-548.	0.4	4
67	Influence of N-butylscopolamine on SUV in FDG PET of the bowel. <i>Annals of Nuclear Medicine</i> , 2009, 23, 471-478.	1.2	3
68	Contributions of treatment centre and patient characteristics to patient-reported experience of haemodialysis: a national cross-sectional study. <i>BMJ Open</i> , 2021, 11, e044984.	0.8	3
69	Percutaneous coronary intervention with drug-eluting stent versus coronary artery bypass grafting: A meta-analysis of patients with left main coronary artery disease. <i>International Journal of Cardiology</i> , 2017, 249, 101-106.	0.8	2
70	Facing Loss and Finding Hope in Narrating Together: Accounts of Parenthood Following the Death of a Child to Muscular Dystrophy. <i>Illness Crisis and Loss</i> , 2022, 30, 689-708.	0.4	1
71	Measuring patients’ experience with renal services in the UK: development and validation of the Kidney PREM. <i>Nephrology Dialysis Transplantation</i> , 2022, , .	0.4	1
72	Commentary on the article: “Maintenance of wellness in patients with obsessive-compulsive disorder who discontinue medication after exposure/response prevention augmentation A randomized clinical trial” Foa EB et al., <i>JAMA Psychiatry</i> . 2022;79(3):193-200 (1). <i>Comprehensive Psychiatry</i> , 2022, 116, 152323.	1.5	0