

Marek Krzanowski

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3782473/publications.pdf>

Version: 2024-02-01

15
papers

473
citations

933447

10
h-index

1058476

14
g-index

15
all docs

15
docs citations

15
times ranked

500
citing authors

#	ARTICLE	IF	CITATIONS
1	Directional Atherectomy Followed by a Paclitaxel-Coated Balloon to Inhibit Restenosis and Maintain Vessel Patency. <i>Circulation: Cardiovascular Interventions</i> , 2017, 10, .	3.9	180
2	Diagnosis and treatment of pelvic congestion syndrome: UIP consensus document. <i>International Angiology</i> , 2019, 38, 265-283.	0.9	86
3	Imaging of all three coronary arteries by transthoracic echocardiography. an illustrated guide. <i>Cardiovascular Ultrasound</i> , 2003, 1, 16.	1.6	52
4	Value of Transthoracic Echocardiography for the Detection of High-Grade Coronary Artery Stenosis: Prospective Evaluation in 50 Consecutive Patients Scheduled for Coronary Angiography. <i>Journal of the American Society of Echocardiography</i> , 2000, 13, 1091-1099.	2.8	34
5	Express LD Vascular Stent in the Treatment of Iliac Artery Lesions: 24-Month Results From the MELODIE Trial. <i>Journal of Endovascular Therapy</i> , 2010, 17, 633-641.	1.5	23
6	PRELUDE Prospective Study of the Serranator Device in the Treatment of Atherosclerotic Lesions in the Superficial Femoral and Popliteal Arteries. <i>Journal of Endovascular Therapy</i> , 2019, 26, 18-25.	1.5	18
7	Myocardial injury after endovascular revascularization in critical limb ischemia predicts 1-year mortality: a prospective observational cohort study. <i>Clinical Research in Cardiology</i> , 2018, 107, 319-328.	3.3	15
8	Posture commonly and considerably modifies stenosis of left common iliac and left renal veins in women diagnosed with pelvic venous disorder. <i>Journal of Vascular Surgery: Venous and Lymphatic Disorders</i> , 2019, 7, 845-852.e2.	1.6	15
9	Transthoracic, harmonic mode, contrast enhanced color Doppler echocardiography in detection of restenosis after percutaneous coronary interventions. Prospective evaluation verified by coronary angiography. <i>European Journal of Echocardiography</i> , 2004, 5, 51-64.	2.3	11
10	Involvement of CD8+ T cell subsets in early response to vascular injury in patients with peripheral artery disease in vivo. <i>Clinical Immunology</i> , 2018, 194, 26-33.	3.2	11
11	Reduced coronary flow reserve in Anderson-Fabry disease measured by transthoracic Doppler echocardiography. <i>Cardiovascular Ultrasound</i> , 2005, 3, 11.	1.6	10
12	Coronary flow reserve assessment. <i>European Heart Journal</i> , 2005, 26, 849-849.	2.2	8
13	Costs of Real-Life Endovascular Treatment of Critical Limb Ischemia: Report from Polandâ€”A European Union Country with a Low-Budget Health Care System. <i>Annals of Vascular Surgery</i> , 2016, 31, 111-123.	0.9	8
14	Percutaneous Transluminal Angioplasty in Patients with Peripheral Arterial Disease Does Not Affect Circulating Monocyte Subpopulations. <i>BioMed Research International</i> , 2016, 2016, 1-9.	1.9	2
15	Regarding â€œLower extremity bypass for critical limb ischemia decreases major adverse limb events with equivalent cardiac risk compared with endovascularâ€”interventionâ€” Journal of Vascular Surgery, 2018, 67, 1637.	1.1	0