

# Sonja A Å elemetjev

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3783508/publications.pdf>

Version: 2024-02-01

15  
papers

183  
citations

1163117

8  
h-index

1125743

13  
g-index

15  
all docs

15  
docs citations

15  
times ranked

233  
citing authors

#	ARTICLE	IF	CITATIONS
1	Enhanced activation of matrix metalloproteinase-9 correlates with the degree of papillary thyroid carcinoma infiltration. <i>Croatian Medical Journal</i> , 2014, 55, 128-137.	0.7	31
2	Coexpressed High Levels of VEGF-C and Active MMP-9 Are Associated With Lymphatic Spreading and Local Invasiveness of Papillary Thyroid Carcinoma. <i>American Journal of Clinical Pathology</i> , 2016, 146, 594-602.	0.7	24
3	Apoptosis and proliferation related molecules (Bcl-2, Bax, p53, PCNA) in papillary microcarcinoma versus papillary carcinoma of the thyroid. <i>Pathology</i> , 2008, 40, 475-480.	0.6	21
4	Changes in the expression pattern of apoptotic molecules (galectin-3, Bcl-2, Bax, survivin) during progression of thyroid malignancy and their clinical significance. <i>Wiener Klinische Wochenschrift</i> , 2015, 127, 337-344.	1.9	18
5	Overexpression of epidermal growth factor receptor and its downstream effector, focal adhesion kinase, correlates with papillary thyroid carcinoma progression. <i>International Journal of Experimental Pathology</i> , 2018, 99, 87-94.	1.3	15
6	MMP-9-1562 C/T single nucleotide polymorphism associates with increased MMP-9 level and activity during papillary thyroid carcinoma progression. <i>Pathology</i> , 2019, 51, 55-61.	0.6	15
7	Concomitant high expression of survivin and vascular endothelial growth factor-C is strongly associated with metastatic status of lymph nodes in papillary thyroid carcinoma. <i>Journal of Cancer Research and Therapeutics</i> , 2018, 14, S114-S119.	0.9	14
8	Evaluation of survivin expression and its prognostic value in papillary thyroid carcinoma. <i>Pathology Research and Practice</i> , 2014, 210, 30-34.	2.3	13
9	Focal adhesion kinase splicing and protein activation in papillary thyroid carcinoma progression. <i>Histochemistry and Cell Biology</i> , 2022, 157, 183-194.	1.7	8
10	Strong Expression of HBME-1 Associates with High-Risk Clinicopathological Factors of Papillary Thyroid Carcinoma. <i>Pathology and Oncology Research</i> , 2015, 21, 735-742.	1.9	6
11	High expression and localization of $\beta$ -catenin and epidermal growth factor receptor identify high risk papillary thyroid carcinoma patients. <i>Experimental and Molecular Pathology</i> , 2018, 105, 181-189.	2.1	6
12	Elevated BANC1 expression levels have different effects on papillary thyroid carcinoma progression depending on the presence of the BRAFV600E mutation. <i>European Journal of Surgical Oncology</i> , 2020, 46, 1835-1842.	1.0	6
13	Novel approach to the measurement of antithyroglobulin antibodies in human serum – application of the quartz crystal microbalance sensors. <i>Talanta</i> , 2021, 223, 121588.	5.5	6
14	Survivin in relation to Bcl-2, Bax and in situ apoptotic cell death in anaplastic thyroid carcinoma. <i>Archives of Biological Sciences</i> , 2011, 63, 955-963.	0.5	0
15	Coexistence of BRAFV600E mutation and EGFR overexpression is highly associated with adverse clinicopathological features of papillary thyroid carcinoma. <i>Archives of Biological Sciences</i> , 2020, 72, 37-44.	0.5	0